

BOEM ENVIRONMENTAL STUDIES PROGRAM: Ongoing Studies

Region: Alaska

Planning Area(s): Beaufort Sea, Chukchi Sea

Title: Distribution and Habitat Use of Fish in the Nearshore Ecosystem of the Beaufort and Chukchi Seas (AK-12-06)

BOEM Information Need(s) to be Addressed: Information is needed on nearshore habitats and fish abundance in the Beaufort and Chukchi Seas to refine the legal definitions of Essential Fish Habitat (EFH) as presented in the Arctic FMP, adopted by NOAA in 2009. Fish in the ecologically fragile nearshore environments are particularly vulnerable to oil spills. The information from this study would be used to better identify and describe EFH in NEPA analyses and in oil spill risk assessments. A better understanding of how fish species respond to habitat variables to improve predictions on distributions under different climate conditions. This project will operate concurrently with other fish sampling efforts (AK-10-06 and AK-11-08) to provide a seamless baseline of forage fish data from the beach to the offshore environment. This study addresses aspects of USGS Recommendations 3.05, 3.07 and 4.01.

Total Cost: \$164,000 plus Joint Funding

Period of Performance: FY 2012-2014

Conducting Organization: NOAA

BOEM Contact: [Catherine Coon](#)

Description:

Background: The Arctic is one of the most rapidly changing ecosystems in the world, yet a large void exists in information on EFH and what species and life stages use these habitats. Information is nonexistent or sparse for fishes in the Arctic, especially in shallow, nearshore waters (shoreline out to 8 m depth). Nearshore habitats are some of the most productive habitats in Alaska and the most at risk to development and oil spills. Many species included in the Arctic Fisheries Management Plan for the Arctic, such as capelin and rainbow smelt, use nearshore habitats at some time in their life but estimates of their abundance and habitat use are poorly documented. Nearshore habitats differ from offshore (>30 m depth) habitats, as do fish assemblages in each area. Recent nearshore research in the Arctic has been limited to the Barrow area, which represents only a small fraction of the nearly 1,700 km of the U.S. Arctic coastline. Prior to major development or transportation activities in the Arctic, more information is needed on fish distribution and habitat use, life history characteristics, food webs, and species at risk to make informed management decisions regarding potential effects from global climate change and human disturbance. In addition, warming conditions in the Arctic will likely result in a reorganization of community structure; new fish species are expected to migrate to the Arctic with unknown consequences to existing stocks and food webs.

Many Arctic fish species are important in the diet of higher-level predators and in Inupiat subsistence fisheries. For example, in the Bering, Beaufort, and Chukchi seas, Arctic cod and

saffron cod occur in the diets of 13 marine mammal species and 20 seabird species. Availability of prey is critical to some Arctic marine mammals such as ice seals, which themselves are important in the diet of polar bears. Larger predators are already under stress by reduced ice cover. The distribution, diversity, and habitat use of nearshore fishes is largely unknown in other areas of the Arctic, especially in the Chukchi Sea. The proposed study would expand fish distribution and habitat use information to the eastern Beaufort Sea and western Chukchi Sea.

Objectives:

- Inventory the distribution and diversity of nearshore fish, their habitat and prey along high priority sites in the Beaufort and Chukchi Seas
- Assess age and diet of fish important as prey species
- Describe oceanographic features of areas with nearshore fish
- Understand how habitat variables like temperature and salinity affect fish species distributions

Methods: Beach and small otter trawl sampling will occur in areas of high importance, defined as locations near oil and gas production, or close to foraging areas for birds or marine mammals, followed by areas with opportunities for research platforms for three sampling seasons. In the summer and fall 2013/2014 two primary areas will be at sites between Barrow and Peard Bay to establish a baseline of fish and habitat use and as reference sites to the Chukchi Lease area.

A random sample of key fish species (e.g., Arctic cod, saffron cod, and capelin) will be collected for age and diet analyses. A sample of select fish species will also be collected, frozen, and archived for later fatty acid and genetic analyses. Habitat will be measured at each sampling site. At seine sites each beach will be profiled according to ShoreZone protocols. Intertidal invertebrates and macroalgae will also be recorded. Additionally, a drop camera will be deployed, depending on water visibility, to search for and identify unusual habitat types and other fish species that may not be captured by seine or trawl.

Models will be generated to predict habitat use by fishes according to habitat variables. Maps will be generated to describe species distribution relative to multiple habitat variables. Information that may lead to NOAA's EFH general distribution will be shared. Relational databases will be built that contain data on species presence and abundance that will expand the current BOEM fish database, be suitable for use in GIS, and complement ShoreZone mapping efforts.

Current Status: Ongoing

Final Report Due: December 2014

Publications Completed: None

Affiliated WWW Sites: <http://www.boem.gov/akstudies/>

Revised Date: July 2012

ESPIS: Environmental Studies Program Information System

All *completed* ESP studies can be found

here: http://www.data.boem.gov/homepg/data_center/other/espis/espisfront.asp