

# **CONOCO LYDONIA CANYON BLOCK 145 No. 1 WELL**

## **Geological and Operational Summary**

Edited by:

Gary M. Edson  
Donald L. Olson  
Andrew J. Petty

U. S. Department of the Interior  
Minerals Management Service  
Gulf of Mexico OCS Region  
Office of Resource Evaluation

New Orleans  
May 2000



# CONTENTS

Abbreviations, vi
Introduction, 1
Operational Summary, 2
Well Velocity Profile, 8
Lithologic Interpretation, 13
Biostratigraphy, 16
Formation Evaluation, 17
Geothermal Gradient, 23
Burial History, 25
Company-submitted Data, 27
Selected References, 28

## ILLUSTRATIONS

- Figure
1. Map of the North Atlantic offshore area showing well Locations, 3
  2. Location plat for the Conoco Block 145 No. 1 well on the OCS Lydonia Canyon NK 19-12 protraction diagram, 4
  3. Daily drilling progress for the Conoco Lydonia Canyon Block 145 No. 1 well, 5
  4. Casing diagram for the Conoco Lydonia Canyon Block 145 No. 1 well, 7
  5. Well velocity profile for the Conoco Lydonia Canyon Block 145 No. 1 well, plotted against depth, 10
  6. Well velocity profile for the Conoco Lydonia Canyon Block 145 No. 1 well, plotted against two-way travel time, 11
  7. Columnar chart of the lithology, biostratigraphy, and paleobathymetry of the Conoco Lydonia Canyon Block 145 No. 1 well, 14
  8. Well temperatures and geothermal gradient for the Conoco Lydonia Canyon Block 145 No. 1 well, 24
  9. Burial diagram for the Conoco Lydonia Canyon Block 145 No. 1 well, 26

## **TABLES**

Table	1. Well statistics, 6
	2. Well velocity data, 8
	3. Well velocity intervals, 9
	4. Biostratigraphy and depositional environments, 16
	5. Well logs, 17
	6. Well log interpretation summary, 18
	7. Sidewall core summary, 19
	8. Conventional core summary, 19
	9. Hydrocarbon shows, 20
	10. Well tests, 21

## ABBREVIATIONS

API	-- American Petroleum Institute
bbbl	-- barrels
BOP	-- Blow out preventer
CNL	-- Compensated neutron log
CPI	-- Carbon Preference Index
COST	-- Continental Offshore Stratigraphic Test
DST	-- drill stem test
EQMW	-- equivalent mud weight
FDC	-- compensated formation density log
FEL	-- from east line
FNL	-- from north line
FSL	-- from south line
FWL	-- from west line
k	-- permeability
KB	-- kelly bushing
LS	-- limestone
m	-- meter (s)
md	-- millidarcy
MYBP	-- million years before present
OCS	-- Outer Continental Shelf
ppf	-- pounds per foot
ppg	-- pounds per gallon
ppm	-- parts per million
psi	-- pounds per square inch
R <sub>o</sub>	-- vitrinite reflectance
SS	-- sandstone
Sw	-- water saturation
TAI	-- thermal alteration index
TD	-- total depth
TIOG	-- threshold of intense oil generation
TOC	-- total organic carbon
UTM	-- Universal Transverse Mercator
φ	-- porosity

## INTRODUCTION

The Conoco Lydonia Canyon (LC) Block 145 No. 1 well was the seventh to be spudded and sixth to be completed of the eight industry wildcat wells drilled on Georges Bank. Spudded on May 13, 1982, this is the northeasternmost of the Georges Bank wells. It is about 11 miles east of the Continental Offshore Stratigraphic Test (COST) G-2 well and is about on trend with the Tenneco LC Block 187 No. 1, Mobil LC Block 273 No. 1, and Shell LC Block 357 No. 1 wells. The geologic trend of the continental shelf is northeast-southwest. The Conoco well was drilled by a semi-submersible rig in 300 feet of water on the shelf about 143 miles east-southeast of Nantucket Island and 20 miles from the shelf edge.

Conoco Inc. was the designated operator for the well, and the company identified two drilling targets in its Exploration Plan, each marked by a strong seismic amplitude anomaly. Conoco inferred that the upper anomaly, at 2.05 seconds, two-way travel time, is associated with a lower Upper Jurassic "...porous shelf-edge calcarenite..." at about 9,100 feet deposited on a paleogeographic high. The deeper target, at 2.40 seconds, was interpreted to be at a depth of about 11,800 feet in Middle Jurassic carbonates. The company stated that dry gas should be the most likely hydrocarbon, trapped by a diagenetic permeability barrier on a southwest-plunging nose.

No specific lithologic or petrophysical properties were identified from well data to account for the seismic anomalies.

There were two gas shows during drilling, of 85 and 121 units, and 62 repeat formation tests were attempted but were not successful. The well bottomed in Middle Jurassic (Callovian to Bathonian) rocks, according to a company-submitted paleontological report. The well was plugged as a dry hole on August 12, 1982.

This report relies on geologic and geophysical data provided to the Minerals Management Service (MMS) by Conoco, according to Outer Continental Shelf (OCS) regulations and lease stipulations. The data were released to the public after the Conoco LC Block 145 lease No. OCS-A-0179 was relinquished on January 15, 1985. Interpretations of the data contained in this report are those of MMS and may differ from those of Conoco. Well depths are measured from kelly bushing unless otherwise stated.

The material contained in this report is from unpublished, undated MMS internal interpretations. No attempt has been made to provide more recent geologic, geochemical, or geophysical interpretations or data, published or unpublished.

This report is initially released on the Minerals Management Service Internet site <http://www.gomr.mms.gov>, and, together with the other Georges Bank well reports, on a single compact disk (CD). At a later date, additional technical data, including well "electric" logs will be added to the CD.

## OPERATIONAL SUMMARY

The Conoco Lydonia Canyon (LC) Block No. 1 well (figures 1 and 2) was drilled by the Keydrill *Aleutian Key* semi-submersible drilling rig to a total depth of 14,500 feet. The well was spudded on May 13, 1982, in 300 feet of water. Daily drilling progress is shown for the well in figure 3 and well statistics are presented in table 1. The geological objectives of the well were an Upper Jurassic reef and Lower Jurassic carbonates.

A 26-inch hole was drilled to 675 feet and then opened to 36 inches. A 30-inch grade B casing was set at 677 feet with 1,390 sacks of class H cement; 940 sacks were mixed with 12-percent gel, and 450 sacks were mixed with 2-percent  $\text{CaCl}_2$ . The casing program is shown in figure 4.

The interval from 650 to 1,350 feet was drilled in 14 hours with a single HTC J-22 bit. Rates of penetration averaged between 70 and 75 feet per hour. This section consisted predominately of soft clays and poorly sorted sand. Background gas averaged between 5 and 10 units with no connection gas. A gelled seawater mud system was used for drilling with weights ranging from 8.7 to 9.7 pounds per gallon. A 20-inch Grade J-55 Vetco 194-pounds-per-foot casing was run to 1,299 feet. The casing was cemented with 2,679 sacks of class H cement but was not tested. The lead slurry was 12 percent gelled and the tail slurry was neat.

The interval from 1,350 feet to 4,898 feet was drilled in 85 hours using four bits. Rates of penetration ranged from 25 to 50 feet per hour in unconsolidated clastics, shales, and limestones. Background gas

ranged from 2 to 5 units. A gelled seawater mud system was used in this section with mud weights between 9.3 and 9.5 pounds per gallon. The grade N-80, 72 lbs/ft, 13 3/8-inch casing was set at 4,860 feet. The casing was cemented with 2,679 sacks of class H cement consisting of 1,860 sacks of 12 percent gelled and 819 sacks neat. The formation was tested to 2,150 psi with a 9.5-pounds-per-gallon mud weight.

The interval from 4,898 feet to the total depth of 14,500 feet was drilled in 768 hours using 24 bits. Rates of penetration ranged from 10 feet per hour in clastics to 25 feet per hour in carbonates. Three conventional cores were taken in this interval, 9,191 to 9,227, 10,622 to 10,652, and 12,419 to 12,448 feet. A small increase in background gas to 85 units between 9,165 to 9,180 feet was noted. A second increase, 121 units, occurred at 13,360 to 13,380 feet. Trip gas increased to 275 units at 13,365 feet. Otherwise, background gas was one to two units through the interval.

A gelled seawater and Drispac mud system was used in drilling the entire well. Mud weight was gradually increased to 10.6 pounds per gallon at 13,300 feet. After the high gas reading was encountered at 13,360 feet, the mud weight was raised to 11.2 pounds per gallon. At 13,750 feet mud weight was again increased to 11.7 pounds per gallon to TD, 14,500 feet.

Well abandonment is shown in figure 4. Open-hole plugs were placed at 9,000 to 9,300 feet and 13,200 to 13,500 feet,







Figure 1. Map of the North Atlantic offshore area showing well locations. The Conoco Lydonia Canyon Block 145 No. 1 well is highlighted in red. Bathymetry is in meters.



**Location Plat**




Figure 2. Location plat for the Conoco Block 145 No. 1 well on the OCS Lydonia Canyon NK 19-12 protraction diagram.




Figure 3. Daily drilling progress for the Conoco Lydonia Canyon Block 145 No. 1 well.

**Table 1. Well statistics**

---


Well Identification:	API No. 61-040-00007 Lease No. OCS-A-0179
Surface location:	Lydonia Canyon NK 19-12 LC Block 145 638 feet from North Line 1,832 feet from West Line  Latitude 40° 49' 58.6439" N Longitude 67° 17' 07.0920" W  UTM coordinates: X = 644,507.12m Y = 4,521,417.12m
Bottomhole location:	61.6 feet N and 70 feet E of surface location
Proposed total depth:	12,000 feet amended to 14,500 feet
Measured depth:	14,500 feet
True vertical depth:	14,398 feet
Kelly bushing elevation:	85 feet
Water depth:	300 feet
Spud date:	May 13, 1982
Reached TD:	August 11, 1982
Off location:	August 25, 1982
Final well status:	Plugged and abandoned

---

Note: All well depths indicated in this report are measured from the kelly bushing (measured depths), unless otherwise indicated. Mean sea level is the datum for the water depth.

using 400 and 450 sacks, respectively, of Class H cement. A cast iron bridge plug was set at 4,800 feet and 200 sacks of Class H cement were squeezed below the plug to 5,000 feet and 50 sacks were spotted above the plug to 4,750 feet. A pressure test was conducted and a surface

plug was placed at 500 to 690 feet with 150 sacks of Class H cement. All three casings were cut 8 feet below mud line. Casing stubs, wellhead, and guide base were recovered and the *Aleutian Key* was off location on August 25, 1982.



## Depth Below KB (feet)

Sea level	85
Seafloor	385
Casings cut	393
Top cement plug	500
30" casing set	677
Bottom cement plug	690
20" casing set	1,299
Top cement plug	4,750
Cement retainer	4,800
13 3/8" casing set	4,860
Bottom of cement	5,000
Top cement plug	9,000
Bottom cement plug	9,300
Top cement plug	13,200
Bottom cement plug	13,500
Total measured depth	14,500

Figure 4. Casing diagram for the Conoco Lydonia Canyon Block 145 No. 1 well.

## WELL VELOCITY PROFILE

Birdwell Division, Seismograph Service Corporation, ran a velocity checkshot survey between 300 and 14,395 feet in the Conoco LC Block 145 No. 1 well. The checkshot data, together with that for the other nine wells drilled on Georges Bank, were given to Velocity Databank, Inc. at their request after all leases had been relinquished or had expired. Velocity Databank calculated interval, average, and

RMS velocities, plotted time-depth curves, and tabulated the data. Table 2 presents well depth, two-way travel time and the calculated velocities for the Conoco LC Block 145 No. 1 well. Figures 5 and 6 show interval velocity, average velocity, and RMS velocity plotted against depth and against two-way travel time, respectively. Well depths are subsea.

**Table 2. Well velocity data**

<b>Depth (feet)</b>	<b>Two-way Travel Time (seconds)</b>	<b>Interval Velocity (feet/sec.)</b>	<b>Average Velocity (feet/sec.)</b>	<b>RMS Velocity (feet/sec.)</b>
300	0.120	5,000	5,000	5,000
915	0.334	5,747	5,490	5,479
1,415	0.506	5,813	5,602	5,592
1,915	0.654	6,756	5,883	5,856
2,415	0.790	7,352	6,160	6,113
2,915	0.914	8,064	6,452	6,378
3,415	1.034	8,333	6,697	6,605
3,915	1.146	8,928	6,947	6,832
4,415	1.244	10,204	7,256	7,098
4,915	1.350	9,433	7,450	7,281
5,415	1.438	11,363	7,747	7,531
5,915	1.524	11,627	8,016	7,762
6,415	1.612	11,363	8,234	7,959
6,915	1.692	12,499	8,484	8,173
7,415	1.776	11,904	8,676	8,350
7,915	1.858	12,195	8,861	8,519
8,415	1.938	12,500	9,040	8,684
8,665	1.974	13,888	9,151	8,779
8,915	2.012	13,157	9,243	8,861
9,165	2.046	14,705	9,360	8,958
9,415	2.078	15,624	9,488	9,061
9,665	2.110	15,625	9,610	9,161
9,915	2.136	19,230	9,784	9,283
10,165	2.178	11,904	9,829	9,334
10,415	2.208	16,666	9,954	9,433

*continued*

**Table 2. Well velocity data--continued**

Depth (feet)	Two-way Travel Time (seconds)	Interval Velocity (feet/sec.)	Average Velocity (feet/sec.)	RMS Velocity (feet/sec.)
10,665	2.232	20,833	10,133	9,556
10,915	2.262	16,666	10,247	9,650
11,165	2.294	15,625	10,341	9,734
11,415	2.318	20,833	10,504	9,849
11,665	2.352	14,705	10,576	9,919
11,915	2.376	20,833	10,729	10,029
12,165	2.404	17,857	10,839	10,120
12,415	2.434	16,666	10,930	10,201
12,665	2.468	14,705	10,991	10,263
12,915	2.490	22,727	11,149	10,373
13,165	2.512	22,727	11,302	10,481
13,415	2.536	20,833	11,429	10,579
13,665	2.554	27,777	11,625	10,700
13,915	2.580	19,230	11,726	10,786
14,165	2.612	15,624	11,782	10,846
14,395	2.632	23,000	11,907	10,938

A lithologic column is also shown in figure 5, together with biostratigraphic subdivision of the section, based on a company-submitted peletonological report.

Five velocity intervals are indicated, which generally correlate with five lithologic intervals penetrated by the well:

**Table 3. Well velocity intervals**

Interval	Depth Range (feet)	Interval Velocity Range (feet/second)	Average Interval Velocity (feet/second)
I	0-1,500	5,000-5,813	5,520
II	1,500-5,200	6,756-10,204	8,438
III	5,200-8,500	11,363-12,500	11,921
IV	8,500-9,800	13,157-15,625	14,600
V	9,800-14,395	11,904-27,777	18,865

**Interval I** The first shot, at 300 feet is at or near the seafloor, and the interval velocity of 5,000 feet per second is for seawater. Two more shots are within this interval and are low velocities (together averaging 5,780 feet per second), appropriate to mudstones and

sediments at shallow depth.

**Interval II** This interval is identified on the basis of intermediate interval velocities, which increase with depth, reflecting transitions from mudstone to siltstone and then to siliciclastics




Figure 5. Well velocity profile for the Conoco Lydonia Canyon Block 145 No.1 well, plotted against depth, with biostratigraphic ages and generalized lithologies. Intervals are explained in text.






Figure 6. Well velocity profile for the Conoco Lydonia Canyon Block 145 No. 1 well, plotted against two-way travel time.

interbedded with limestones. This interval is Upper and Lower Cretaceous.

**Interval III** This interval is identified on the basis of a slightly higher interval velocity trend, which may be a consequence of more abundant limestone interbeds in the lowermost Lower Cretaceous and upper portion of the Upper Jurassic.

**Interval IV** This interval is identified on the basis of moderately high interval velocities in the lower portion of the Upper Jurassic and upper portion of the

Middle Jurassic. Rock types are siliciclastics interbedded with limestones.

**Interval V** This interval is identified on the basis of high and very high interval velocities alternating with moderately low (for this depth range) velocities. As with other Georges Bank wells, the data are appropriate to interbedded siliciclastic lithologies, limestone, dolomite, and anhydrite, (and halite in some other wells). According to a company-submitted report, this interval is Middle Jurassic.

# LITHOLOGIC INTERPRETATION

Taken and adapted from R. Smith, MMS internal report

Well cuttings from the Conoco LC Block 145 No. 1 well were collected at 30-foot intervals from 675 to 14,500 feet (TD). Additional lithologic control was provided by conventional and sidewall cores. The lithologic descriptions of this report are based on examination of drill cutting samples with a binocular microscope, on conventional and sidewall core descriptions, and on the formation evaluation "mud" log. Depths of lithologic units are adjusted with reference to electric logs. All depths are from kelly bushing. Lithologic intervals are shown in figure 7.

The interval from 675 to 2,230 feet consists of unconsolidated, gray to black, slightly calcareous clay with abundant fossil fragments, bedded and disseminated glauconite, trace pyrite, lignite, calcite, and siliciclastic silt and sand grains. A glauconite bed exists at 1,360 feet near the Cretaceous-Tertiary boundary. A Cretaceous-Tertiary glauconite bed is a distinct regional marker also seen in the Mobil LC Block 312 No. 1 and the Exxon CO Block 975 No. 1 wells. Below the glauconite bed is a slightly calcareous, poor to moderately sorted sand, having subrounded to subangular and translucent to clear grains. Within the sand are traces of glauconite, fossil fragments, and pyrite.

The interval from 2,230 to 4,900 feet consists of siltstone, mudstone, limestone, and sandstone. The upper third of the interval is light- to dark-gray, calcareous, variably fossiliferous, sandy siltstone with a trace of glauconite. The middle third is moderately consolidated, slightly calcareous mudstone or claystone with


sparse glauconite, underlain by calcareous, sandy siltstone. The bottom third is gray to buff, silty limestone interbedded with slightly fossiliferous sands. The sands are poorly sorted, with subrounded to rounded and medium to coarse grains. Glauconite, pyrite, and lignite are present in trace amounts.

The section between 4,900 and 6,920 feet is massive, blocky, white to gray, granular to microcrystalline, and soft to firm limestone with some chalky portions, shale breaks, and sandstone interbeds. The upper 100 feet is argillaceous and slightly sandy.

The interval from 6,920 to 8,630 feet consists of silty, variably oolitic, light- to medium-gray, soft to firm, microcrystalline limestone interbedded with light-gray, very fine-grained calcareous sandstone, light-gray siltstone, and dark gray, calcareous shale. Siliciclastic beds contain trace amounts of mica, pyrite, and glauconite.

The interval from 8,630 to 10,810 feet is massive, shaley, oolitic limestone with some silt and shale interbeds. The limestone is white to gray, soft to firm, and microcrystalline; the siliciclastic interbeds have traces of fossil fragments, pyrite, and lignite.

The interval from 10,810 to 12,600 feet is massive, white to gray, firm, coarsely crystalline to microcrystalline, variably oolitic limestone with thin shale partings, becoming sandy in the bottom 300 feet.



EXPLANATION

- |  |                        |  |   |  |                  |
|--|------------------------|--|---|--|------------------|
|  | Shale or Clay          |  | Shaly Sandstone                         |  | Anhydrite        |
|  | Silty Shale            |  | Silty Sandstone                         |  | Halite           |
|  | Sandy Shale            |  | Calcareous Sandstone                    |  | Tuff or Ash      |
|  | Calcareous Shale       |  | Sandstone with Pebbles and / or Fossils |  | Glauconite       |
|  | Shale with Fossils     |  | Limestone                               |  | Fossils          |
|  | Shale with Pyrite      |  | Shaly Limestone                         |  | Igneous          |
|  | Siltstone              |  | Silty Limestone                         |  | Coal             |
|  | Shaly Siltstone        |  | Sandy Limestone                         |  | Unconformity     |
|  | Sandy Siltstone        |  | Limestone with Fossils                  |  | Marker Fossil    |
|  | Calcareous Siltstone   |  | Limestone with Anhydrite                |  | Casing Point     |
|  | Siltstone with Fossils |  | Oolitic Limestone                       |  | Hydrocarbon Show |
|  | Siltstone with Pyrite  |  | Dolomite                                |  |                  |
|  | Sandstone              |  |   |  |                  |

Figure 7. Columnar chart of the lithology, biostratigraphy, and paleobathymetry of the Conoco Lydonia Canyon Block 145 No. 1 well. Lithologic interpretations from examination of cuttings; lithologic breaks picked from well logs. Stage tops and paleobathymetry from Conoco.

Traces of pyrite, lignite, quartz, and anhydrite are also present.

The section between 12,600 and 12,700 feet is a sandstone bed bounded above and below by gradational contacts with the massive carbonates. The sandstone is slightly calcareous, variably shaley, poorly consolidated, and poorly sorted, with subrounded to subangular, and fine to medium grains. The sandstone contains traces of mica and pyrite.

The interval from 12,700 to 14,500 feet (TD) is sandy to shaley limestone with sparse anhydrite in the upper half and with dolomite and abundant anhydrite interbeds in the lower half. The limestone is gray to brown, hard, and microcrystalline. The dolomite is white to brown, slightly calcareous, and microcrystalline. The anhydrite is white to light brown, soft to firm, and blocky to nodular.

### **POTENTIAL RESERVOIR ROCKS**

Data from well log analysis (table 6), analysis of 73 sidewall cores (table 7), and analysis of three conventional cores (table 8) show generally decreasing porosity from 5,000 feet to total depth, 14,500 feet, in rocks penetrated by the Conoco LC Block 145 No. 1 well. According to well log interpretation, most porosities of "potential reservoir intervals" are better

than 20 percent above 7,500 feet and poorer than 10 percent below 9,500 feet.

Among sidewall cores, permeabilities decrease with depth except for specific permeable (and porous) beds, such as a water-saturated sandstone at 13,358 to 13,366 feet, which has porosities of 17.2 to 25.8 percent and permeabilities of 12 to 116 md (based on three sidewall cores).

No optical kerogen nor molecular geochemical analyses were done by MMS for this well. However, an organic geochemical analysis report by Geochem Laboratories, Inc., of Houston, was submitted to MMS by Conoco. A summary table in the report refers to rocks above 7,690 feet as "moderately immature, poor oil and associated gas source potential." Rocks from 7,690 to 9,940 feet are "moderately mature, poor oil and associated gas source character." Rocks from 9,940 to TD of 14,500 feet are "mature, poor oil and associated gas source character."

In summary, the best porosities and permeabilities are found in the shallow, thermally immature section, above 7,500 feet, in the Conoco LC Block 145 No. 1 well. However, rocks within the moderately mature and mature part of the section are organically lean and have poor oil and gas generating potential.

## BIOSTRATIGRAPHY

This biostratigraphic and paleoenvironmental summary is taken from a report done for Conoco by

International Biostratigraphers Inc. for the Conoco LC Block 145 No. 1 well. No MMS paleontological interpretations are available.

**Table 4. Biostratigraphy and depositional environments**

<b>Series</b>	<b>Age</b>	<b>Depth Interval (feet)</b>	<b>Environment of Deposition (depths in feet)</b>
Miocene	Middle Miocene	675-1,360	Outer shelf
Eocene	Middle Eocene	1,360-1,390	
Late Cretaceous	Santonian	1,390-1,660	
	Coniacian	1,660-1,870	
	Turonian	1,870-2,110	
	Cenomanian	2,110-2,350	
Early Cretaceous	Albian	2,350-2,830	Inner shelf
	Aptian	2,830-3,730	Inner shelf (2,830-3,400) Middle shelf (3,400-3,700)
	Barremian	3,730-5,000	Inner shelf to marginal marine
	Hauterivian to Berriasian	5,000-5,440	Middle shelf
Late Jurassic	Tithonian to Late Kimmeridgian	5,440-6,430	
	Early Kimmeridgian to ?Oxfordian	6,430-8,410	Middle shelf (6,430-7,400) Inner shelf (7,400-8,410)
	Oxfordian	8,410-9,600	Inner shelf
Middle Jurassic	Callovian to Bathonian	9,600-14,500	Inner shelf to marginal marine

## FORMATION EVALUATION

Taken and adapted from R. Nichols, MMS internal report

Schlumberger Ltd. ran the following “electric” logs in the Conoco LC Block 145 No. 1 well to provide information

for stratigraphic correlation and for evaluation of formation fluids, porosity, and lithology:

**Table 5. Well logs**

Log Type	Depth Interval (feet below KB)
DISFL/Sonic (dual induction spherically focused log/sonic)	1,300-14,482
CNL/FDC (compensated neutron log/compensated formation density)	1,300-14,485
HDT (high resolution dipmeter)	4,866-14,485
CYBERDIP	9,600-14,485
RFT (repeat formation tester)	9,159-14,460
CYBERLOOK	4,672-14,482
CORIBAND	4,875-9,750

Exploration Logging, Inc. provided a formation evaluation “mud” log, which included a rate of penetration curve, sample description, and a graphic presentation of any hydrocarbon shows encountered (1,350 to 14,500 feet). In addition, a delta chloride log (4,800-14,500 feet), a drilling data pressure log (1,350 to 14,500 feet), and a pressure evaluation log (600 to 14,500 feet) were included.

The electric logs, together with the mud log and other available data, were analyzed in detail to determine the thickness of potential reservoirs, average porosities, and feet of hydrocarbon present. Reservoir rocks with porosities less than 5 percent were disregarded. A combination of logs was used in the analysis, but a detailed lithologic and reservoir property determination from samples, conventional cores, and

sidewall cores, in addition to full consideration of any test results, is necessary to substantiate the following estimates as shown in table 6. The electric logs were of poor quality, probably due to poor hole condition. Substantial hole enlargement occurs from 2,620 to 3,220 feet, 4,120 to 4,390 feet, 8,210 to 10,530 feet, 10,750 to 12,220 feet, and 12,780 to 14,450 feet. The SP was cyclic and of poor character over the entire interval. The hole enlargement also had considerable impact on other log responses. The DISFL exhibited a false “annulus” appearance over much of the section, with suspiciously low  $IL_m$  readings. The porosity logs were undoubtedly affected by the washouts as well.

Seventy-three successful sidewall cores were taken in the well, and their petrophysical properties are summarized in table 7. Sidewall core porosities compare favorably to FDC porosities in many cases. However,

**Table 6. Well log interpretation summary**

Series	Depth Interval (feet)	Potential Reservoir <sup>1</sup> (feet)	Ave $\phi$	SW %	Feet of Hydrocarbon
UK-LK	2,254-2,422	97 (?)	35	NC*	NC*
LK	2,635-2,790	75 (?)	35	"	"
	2,866-3,124	62 (?)	35	"	"
	3,658-4,050	139 (?)	34	"	"
	4,222-4,342	86	29	"	"
	4,432-4,505	73	26	"	"
	4,551-4,570	11	30	"	"
	4,623-4,985	202	33	"	"
	5,235-5,263	28	16	66	"
UJ	5,515-5,538	23	25	NC*	"
	5,937-5,954	15 (?)	25	"	"
	6,632-6,650	15 (?)	24	"	"
	7,052-7,080	28	22	"	"
	7,173-7,184	11	22	"	"
	7,514-7,524	10	20	"	"
	7,992-8,010	16	17	"	"
	8,990-9,010	17	15	58	"
	9,148-9,182	17	10	19	17 <sup>2</sup> (?)
MJ	9,604-9,633	24	8	NC*	NC*
	10,620-10,632	12	9	"	"
	10,658-10,670	12	9	"	"
	13,356-13,402	24	8 <sup>3</sup>	67	"

\*Not calculated

<sup>1</sup>Generally in beds > 10 feet thick and > five percent.

<sup>2</sup>Total gas reading = 85 units (C<sub>1-2,3</sub>), but no show in samples or SWC

<sup>3</sup>Indicated porosity is probably too high; the zone is washed out. Total Gas reading = (C<sub>1</sub> only)

below 9,200 feet, the sidewall core porosities are closer to those recorded by the Sonic Log.

Three conventional cores were taken in the well. A detailed petrologic examination was performed and the results are shown in table 8.

The porosities derived from the sonic and density logs compare favorably to the core

analysis for core No. 1 and the porosities derived from sonic, density, and neutron logs compare favorably to the core analysis for core No. 2. However, all log-derived porosities (6 to 9 percent) are substantially higher than the core analysis indicates for core No. 3. The substantial difference could be caused by increased circulation time for hole conditioning prior to the coring operation. This could have induced deep invasion and erratic



**Table 7. Sidewall core summary**

Depth Interval (feet)	Number of Cores	Lithology	Porosity (%)		Permeability (md)	
			Range	Ave.	Range	Ave.
5,270.0-6,992.0	8	SS	16.9-23.8	21.3	2.2-74.0	31.3
7,028.0-8,997.0	21	SS/LS	8.6-24.5	17.7	< 0.1-45.0	9.7
9,000.0-10,699.0	20	LS/SS	12.5-22.7	16.6	< 0.1-37.0	7.1
11,747.0-12,878.0	5	LS	8.3-13.5	11.0	< 0.1-0.8	0.4
13,208.0-13,972.0	19	SS/LS	13.3-25.8	18.5	2.7-116.0	18.4

**Table 8. Conventional core summary**

Core No.	Depth Interval (feet)	Lithology	Porosity (%)		Permeability (md)		Gas % Bulk
			Range	Ave.	Range	Ave.	
1	9,191.0-9,226.5	LS	1.5-6.1	2.9	< 0.01-0.13	0.04	< 5
2*	10,622.0-10,652.3	LS	2.1-13.4	7.7	< 0.01-172.0	19.0	0.6-9.5
3	12,419.0-12,448.0	SiS	1.6-3.3	2.4	< 0.01-0.23	0.09	< 5

\*Log analysis results show  $\phi = 1-3\%$ ,  $SW = 42\%$ . Total gas reading on the mud log = 0.

washouts. If so, the fluid absorption and increased borehole size may have resulted in higher porosity readings for the sonic, density, and neutron logs.

Results of the HDT survey were recorded on a dipmeter arrow plot from 4,866 to 14,485 feet. Possible structural anomalies may be present at 6,275, 9,475, 12,750, and 13,325 feet. From 6,500 to 10,900 feet, the dip is essentially westward at 2 to 4 degrees. From 10,900 to 12,400 feet, the dip decreases to 1 to 3 degrees; from 12,400 to 13,100 feet, the dips increase in magnitude (10 to 30 degrees) and change to a predominantly eastward direction. Below 13,100 feet, the dips become more erratic in both direction and magnitude. From 14,200 to 14,500 feet, the dips range from 1 to 16 degrees and

are predominantly westward again.

Table 9 lists all shows of hydrocarbon encountered in this well. None of the shows encountered were judged to be significant. Table 10 shows the well test results. Pressures appeared normal, but no formation fluids were recovered.

A normal pressure gradient (approx 9.0 ppg eqmw) was encountered to a depth of 8,800 feet. The estimated pore pressure increased to 9.4 ppg eqmw at 9,200 feet. The increase in pore pressure was accompanied by decreasing  $D_{xc}$  values, slightly increasing chlorides, and an increase in background gas. Between 13,000 and 14,500 feet, the pore pressure was estimated to have increased at a moderate rate from 9.4 ppg to 11.0 ppg. Changes in flowline temperature and magnitude of trip gas accompanied the increasing pore pressure.

**Table 9. Hydrocarbon shows**

Depth Interval	Drilling Break (feet/hour)	Sample Description (Mud Log)	Total gas b.g.	Chromatograph	Cuttings Gas	Conventional Cores					Sidewall cores					Well Log Interpretation			Well Tests
						0	K	O <sub>p</sub>	O <sub>b</sub>	G <sub>b</sub>	0	K	O <sub>p</sub>	O <sub>b</sub>	G <sub>b</sub>	0	SW		
9,148 - 9,182	12' - 42'	LS, dull yel flu, no cut	3 85	C <sub>1-2-3</sub>	1	-	9,166 - 9,171	14	2							9,148 - 9,182	10	19	12 RFT attempts (mud sets)
10,632 - 10,660	-	LS, dull yel flu, no cut	2 2	C <sub>1</sub>	1	10,622 - 10,652	2.6 - 13.4	<.01 -172	.6 - 9.5							10,632 - 10,660	1-3	42	None
13,230 - 13,244	-	LS	2 10	C <sub>1-2-3</sub>	1	-	-									13,230 - 13,244	0-1	39	None
13,356 - 13,402	14' - 25'	LS/DOL, dull yel flu, no cut	2 121	C <sub>1</sub>	3	-	-									13,356 - 13,402	8	67	40 RFT attempts (mud sets)

**Table 10. Well tests****Run 1**

<b>Depth</b>	<b>HP</b>	<b>Min. Press.</b>	<b>SIP</b>	<b>Remarks</b>
9,159	5,657	125	159	
9,161	5,660			
9,163	5,661			
9,165	5,663			
9,175	5,689	1,063	4,167	
9,178	5,560	4,147	4,158	
9,180	5,652	3,767	4,179	
9,182	5,664	4,068	4,168	
9,187	5,690	4,088	4,157	
9,192	5,677			
9,194	5,682			
9,195	5,677	310	332	
12,600	7,740			Hole size prevented tool setting
12,603	8,502			"
12,603	7,746			"
13,390	8,261			"
13,392	8,260			"
13,395	8,261			"
13,810	8,501			"
13,812	8,940			"
14,190	8,761			"
14,210	8,760			"
14,460	8,739			"

**Run 2**

<b>Depth</b>	<b>HP</b>	<b>Min. Press.</b>	<b>SIP</b>	<b>Remarks</b>
9,178	5,536			Mud Set
9,515		2,397	4,333	
9,564	-			Mud Set
9,615	5,806			"
9,615	-			
9,615		4,322	4,334	
12,600	7,636			Mud Set
13,344.5	-			"
13,356	8,155			"
13,358	8,151			"

*continued*

**Table 10. Well tests, Run 2--continued**

<b>Depth</b>	<b>HP</b>	<b>Min. Press.</b>	<b>SIP</b>	<b>Remarks</b>
13,360	8,156			Mud Set
13,362	8,158			"
13,364	8,156			"
13,364	8,140			"
13,364.5	8,139			"
13,366	8,156			"
13,367	8,134			"
13,367.5	8,144			"
13,368	8,142			"
13,368	8,155			"
13,368	8,163			"
13,370	8,166			"
13,370	8,142			"
13,370	8,159			"
13,370.5	8,157			"
13,371	8,156			"
13,372	8,157			"
13,374	8,148			"
13,374	8,156			"
13,376	8,154			"
13,378	8,160			"
13,379	8,153			"
13,380	8,161			"
13,382	8,160			"
13,384	8,164			"
13,386	8,162			"
13,388	8,164			"
13,390	8,166			"
13,392	8,169			"
13,392	8,150			"
13,394	8,158			"
13,408	-			"
13,410	-			"
13,410	8,182			"
13,412	-			"
13,416	-			"
13,570	8,266			"
13,952	8,503			"

## GEOHERMAL GRADIENT

Figure 8 shows bottomhole temperatures for three logging runs in the Conoco LC Block 145 No.1 well plotted against depth. A temperature of 60 °F is assumed at the seafloor at an indicated depth of 385 feet (300-foot water depth plus 85-foot kelly bushing elevation). Shown also is a straight-line graph between the seafloor

and total-depth temperatures in order to represent an overall geothermal gradient for the well, which is 1.06 °F/100 ft., the lowest gradient among the Georges Bank wells. Gradients for the other wells range from 1.14 °F/100 ft. (COST G-1) to 1.40 °F/100 ft. (COST G-2).




Figure 8. Well temperatures and geothermal gradient for the Conoco Lydonia Canyon Block 145 No. 1 well. Well temperatures from bottomhole temperatures of logging runs. Geothermal gradient based on bottomhole temperature of deepest logging run.

## BURIAL HISTORY

The burial history model for the stratigraphic section penetrated by the Conoco LC Block 145 No. 1 well (figure 9) is based on the biostratigraphic determinations of International Biostratigraphers Inc. (IBI), Houston, (figure 7) and the Cretaceous and Jurassic time scales of Van Hinte (1976a and 1976b). The burial model for this well is similar to those of the COST G-1 and G-2 wells and the other industry wildcat wells on Georges Bank. In general, they show rapid initial rift-stage (Late Triassic) subsidence and equally rapid early drift-stage (Middle Jurassic) subsidence. In the Early Cretaceous Epoch, basinal subsidence progressively decreased until

some time in the Tertiary Period, when subsidence may have increased again moderately. However, using sea level as the zero datum exaggerates apparent Tertiary thickness and subsidence. For the Conoco LC Block 145 No. 1 well, no rift-stage rocks were penetrated, according to IBI. Rapid Oxfordian-Calloviaian burial reflects early drift-stage subsidence. The burial rate is very low in the Late Cretaceous (Santonian) and increases in the Tertiary Period (Miocene and earlier). However, the indicated Tertiary burial rate includes the water-depth effect. In constructing figure 9, no adjustments have been made for sedimentary compaction or for section removed by erosion.




Figure 9. Burial diagram for the Conoco Lydonia Canyon Block 145 No. 1 well. Based on stage tops from Conoco (See Figure 7).



## COMPANY-SUBMITTED DATA

Data and reports were submitted by Conoco Inc. to MMS when the Conoco LC Block 145 No. 1 well was drilled, as required by Federal regulations and lease stipulations. Items of general geological, geophysical, and engineering usefulness are listed below. Items not listed include routine submittals required by regulation and detailed operations information, such as the Exploration Plan, Application for Permit to Drill, daily drilling reports, monthly reports, well location survey, and drilling pressure and temperature data logs. Well "electric" logs are listed in the **Formation Evaluation** chapter. Listed and unlisted company reports and data are available through the Public Information Unit, Minerals Management Service, Gulf

of Mexico OCS Region, 1201 Elmwood Park Boulevard, New Orleans, Louisiana 70123-2394; telephone (504)736-2519 or 1-800-200-GULF, FAX (504)736-2620. Well logs are available on microfilm from the National Geophysical Data Center, 325 Broadway Street, Boulder CO 80303-3337, attn. Ms Robin Warnken; telephone (303)497-6338, FAX (303)497-6513; e-mail rwarnken@NGDC.NOAA.GOV.

At a later date, additional original technical data, including well logs, will be added to the compact disk (CD) version of the Georges Bank well reports. The CD will be available from the Gulf of Mexico OCS Region Public Information Unit.

### Selected company-submitted data

Final Well Report (summaries of data and interpretations for drilling and engineering, formation pressure, geology and shows, evaluation and testing, and data inventory), Exploration Logging of U.S.A., Inc. (Exlog), undated.

Physical formation "mud" log, EXLOG, undated.

Seismic velocity survey (checkshot survey), Birdwell Division, Seismograph Service Corp., Tulsa OK, undated.

Core analysis report (sidewall and conventional cores), Core Laboratories, Inc., Dallas TX, undated.

Biostratigraphy report (biostratigraphy, environments of deposition), International Biostratigraphers, Inc., Houston TX, 11/82.

Geochemical analysis report (visual kerogen, total organic carbon, Rock-Eval, C<sub>1</sub>-C<sub>7</sub> gas chromatograms, C<sub>15+</sub> soxhlet extraction, saturate hydrocarbon analysis), Geochem Laboratories, Inc., Houston, 10/23/82.

Analysis of hydrocarbons in the RFT sample, Conoco, Exploration Research Division, Ponca City OK, 09/82.

## SELECTED REFERENCES

This list is compiled from published and unpublished Minerals Management Service and USGS Conservation Division reports on Georges Bank wells. Not all of the references could be located and verified.

- Albrecht, P., 1970, Etude de constituents organiques des series sedimentaries de Logbaba et Messel. Transformations deagenetiques: Universite de Strasbourg, Memoires du Service de la Charge Geologique d'Alsac et de Lorraine, no. 32, 119 p.
- Amato, R.V. and J.W. Bebout, 1978, Geological and Operational Summary, COST No. GE-1 Well, Southeast Georgia Embayment Area, South Atlantic OCS: U. S. Geological Survey Open-File Report 78-668, 122 p.
- Amato, R. V. and J. W. Bebout (eds.), 1980, Geologic and Operational Summary, COST No. G-1 Well, Georges Bank Area, North Atlantic OCS: U. S. Geological Survey Open-File Report 80-268, 112 p.
- Amato, R.V., and E.K. Simonis (eds.), 1979, Geologic and Operational Summary, COST No. B-3 Well, Baltimore Canyon Trough Area, Mid-Atlantic OCS: U.S. Geological Survey Open-File Report 79-1159, 118 p.
- Amato, R.V. and E.K. Simonis,(eds.), 1980, Geologic and Operational Summary, COST No. G-2 Well, Georges Bank Area, North Atlantic OCS: U. S. Geological Survey Open-File Report 80-269, 116 p.
- BBN-Geomarine Services Co., 1975, COST wellsite G-1, Georges Bank, engineering geology interpretation of high-resolution geophysical data: Houston, Texas, 11 p.
- Ballard, R. D. and E. Uchupi, 1975, Triassic rift structure in Gulf of Maine: American Association of Petroleum Geologists Bulletin, v. 59, no. 7, p. 1041-1072.
- Bayliss, G. S., 1980, Source-rock evaluation reference manual: Houston, Texas, Geochem Laboratories, Inc., 80 p.
- Bebout, J. W., 1980, Observed stratigraphic distribution of spores, pollen, and *incertae sedis* palynomorphs in the Tertiary section of the COST No. B-2 well, Baltimore Canyon, Atlantic Outer Continental Shelf: Palynology, v. 4, p. 181-196.
- Bebout, J. W., 1981, An informal palynologic zonation for the Cretaceous System of the United States Mid-Atlantic (Baltimore Canyon area) Outer Continental Shelf: Palynology, v. 5, p. 159-194.
- Berggren, W.A., D.V. Kent, C.C. Swisher III, and M.P. Aubry, 1995, A revised Cenozoic geochronology and chronostratigraphy; in Geochronology Time Scales and Global Stratigraphic Correlation, SEPM Special Publication no. 54, p. 129-212.
- Bhat, H., N. J. McMillan, J. Aubert, B. Porthault, and M. Surin, 1975, North American and African drift--the record in Mesozoic coastal plain rocks, Nova Scotia and Morocco, in Yorath, C. J., E. R. Parker, and D. J. Glass, (eds.), Canada's Continental Margins and Offshore Petroleum Exploration: Canadian Society of Petroleum Geologists Memoir 4, p. 375-389.
- Brideau, W. W. and W. C. Elsick, (eds.), 1979, Contributions of stratigraphic palynology (v. 2), Mesozoic Palynology: American Association of Stratigraphic Palynologists Contributions Series No. 4.

- Bronnimann, P., 1955, Microfossils *incertae sedis* from the Upper Jurassic and Lower Cretaceous of Cuba: *Micropaleontology*, v. 1, pp. 28, 2 pls., 10 text.
- Bujak, J. P., M. S. Barss, and G. L. Williams, 1977, Offshore east Canada's organic type and color and hydrocarbon potential: *Oil and Gas Journal*, v. 75, no. 15, p. 96-100.
- Bujak, J. P. and M. J. Fisher, 1976, Dinoflagellate cysts from the Upper Triassic of Arctic Canada: *Micropaleontology*, v. 22, p. 44-70, 9 pls.
- Bujak, J. P. and G. L. Williams, 1977, Jurassic palynostratigraphy of offshore eastern Canada, *in* Swain, F. M., (ed.), *Stratigraphic Micropaleontology of Atlantic Basin and Borderlands*: New York, Elsevier Scientific Publishing Co., p. 321-339.
- Bukry, D., 1969, Upper Cretaceous coccoliths from Texas and Europe: *University of Kansas Paleontological Contributions*, Art. 5 (Protista 2), p. 1-9, 50 pls., 1 text.
- Burk, C. A. and C. L. Drake, (eds.), 1974, *Geology of Continental Margins*: New York, Springer-Verlag, 1,009 p.
- Burke, K., 1975, Atlantic evaporites formed by evaporation of water spilled from Pacific, Tethyan, and southern oceans: *Geology*, v. 3, no. 11, p. 613-616.
- Cepek, P. and W. W. Hay, 1970, Zonation of the Upper Cretaceous using calcareous nannoplankton: *Palaontologische Abhandlungen, Abteilung B Palabotanik, Band III, Heft 3/4*, p. 333-340.
- Cita, M. B. and S. Gartner, 1971, Deep Sea Upper Cretaceous from the western North Atlantic: *in* *Proceedings II International Planktonic Conference, Roma, 1970*: Rome, Edizioni Tecnoscienza, v. 1, p. 287-319.
- Clarke, R. F. A. and J. P. Verdier, 1967, An investigation of microplankton assemblages from the chalk of the Isle of Wight, England: *Verhandelingen der Koninklijke Nederlandse Akademie van Wetenschappen, Afdeling Natuurkunde, and Eerste Reeks*, 24, p. 1-96.
- Claypool, G. E., C. M. Lubeck, J. P. Baysinger, and T. G. Ging, 1977, Organic geochemistry, *in* Scholle, P. A., (ed.), *Geological studies on the COST No. B-2 well, U. S. Mid-Atlantic Outer Continental Shelf area*: U. S. Geological Survey Circular 750, p. 46-59.
- Connan, J. 1974, Time-temperature relation in oil genesis: *American Association of Petroleum Geologists Bulletin*, v. 58, no. 12, p. 2516-2521.
- Core Laboratories, Inc., 1976, *Core studies, COST Atlantic well No. G-1, Georges Bank, Offshore Atlantic Ocean*: Dallas, Texas, 153 p.
- Core Laboratories, Inc., 1977a, *Core studies, COST Atlantic well No. G-2, Georges Bank, Offshore Atlantic Ocean*: Dallas, Texas, 298 p.
- Core Laboratories, Inc., 1977b, *Geochemical service report, COST G-2 Atlantic well, Georges Bank, offshore Massachusetts, U. S. A.*: Dallas, Texas, 147 p.
- Council on Environmental Quality, 1974, *OCS oil and gas--An environmental assessment--A report to the President by the Council on Environmental Quality*: Washington, D. C. (U. S. Government Printing Office), Stock No. 4000-00322, v. 1, 214 p.

- Cousminer, H. L., 1984, Canadian dinoflagellate zones (Middle Jurassic to Middle Eocene) in Georges Bank Basin (abstract): Proceedings of the American Association of Stratigraphic Palynologists, Arlington, Virginia, v. 9, p. 238.
- Cousminer, H. L., W. E. Steinkraus, and C. E. Fry, 1982, Biostratigraphy and thermal maturation profile, Exxon 133 No. 1 (OCS-A-0170) well section: Unpublished Report, Minerals Management Service.
- Cousminer, H. L., W. E. Steinkraus, and R. E. Hall, 1984, Biostratigraphic restudy documents Triassic/Jurassic section in Georges Bank COST G-2 well (abstract): Proceedings of the American Association of Petroleum Geologists, Annual Meeting, San Antonio, Texas, v. 68, no. 4, p. 466.
- Davey, R. J., 1979, The stratigraphic distribution of dinocysts in the Portlandian (latest Jurassic) to Barremian (Early Cretaceous) of northwest Europe: American Association of Stratigraphic Palynologists Contributions, Series No. 5B, p. 49-81.
- Davey, R. J. and J. P. Verdier, 1974, Dinoflagellate cysts from the Aptian type sections at Gargas and La Bedoule, France: Paleontology, v. 17, pt. 3, p. 623-653.
- Davies, E. H., 1985, The miospore and dinoflagellate cyst zonation of the Lias of Portugal: Palynology, v. 9, p. 105-132.
- Dorhofer, G. and E. H. Davies, 1980, Evolution of archeopyle and tabulation in Rhaetogonyaulacian dinoflagellate cysts: Royal Ontario Museum, Life Sciences Miscellaneous Publications, p. 1-91, fig. 1-40.
- Dow, W. G., 1974, Application of oil-correlation and source-rock data to exploration in Williston Basin: American Association of Petroleum Geologists Bulletin, v. 58, no. 7, p. 1253-1262.
- Dow, W. G., 1977, Kerogen studies and geological interpretations: Journal of Geochemical Exploration, v. 7, p. 79-99.
- Drake, C. L., J. I. Ewing, and H. Stockard, 1968, The continental margin of the eastern United States: Canadian Journal of Earth Science, v. 5, no. 4, p. 993-1010.
- Drake, C. L., M. Ewing, and G. H. Sutton, 1959, Continental margins and geosynclines--The east coast of North America north of Cape Hatteras, in Aherns, L. H., and others, (eds.), Physics and Chemistry of the Earth, v. 3: New York, Pergamon, p. 110-198.
- Eliuk, L. S., 1978, the Abenaki Formation, Nova Scotia, Canada--A depositional and diagenetic model for a Mesozoic carbonate platform: Bulletin of Canadian Petroleum Geology, v. 26, no. 4, p. 424-514.
- Emery, K. O. and E. Uchipi, 1972, Western North Atlantic Ocean--Topography, rocks, structure, water, life, and sediments: American Association of Petroleum Geologists Memoir 17, 532 p.
- Evitt, W. R., (ed.), 1975, Proceedings of a forum on dinoflagellates: American Association of Stratigraphic Palynologists Contributions, Series No. 4, 76 p.
- Folger, D. W., 1978, Geologic hazards on Georges Bank--an overview: Geological Society of America Abstracts with Programs, v. 10, no. 1, p. 42.
- Fry, C. E., 1979, Geothermal gradient, in Amato, R. V. and E. K. Simonis (eds.), Geologic and Operational Summary, COST No. B-3 well, Baltimore Canyon Trough Area, Mid-Atlantic OCS: U. S. Geological Survey Open-File Report 79-1159, p. 64-65.

- Gartner, S., Jr., 1968, Coccoliths and related calcareous nanofossils from Upper Cretaceous deposits of Texas and Arkansas: University of Kansas Paleontological Contributions, no. 48, Protista, v. 48, Art. 1, p. 1-56.
- GeoChem Laboratories, Inc., 1976, Hydrocarbon source facies analysis, COST Atlantic G-1 well, Georges Bank, offshore Eastern United States: Houston, Texas, 10 p.
- GeoChem Laboratories, Inc., 1977, Hydrocarbon source facies analysis, COST Atlantic G-2 well, Georges Bank, offshore eastern United States: Houston, Texas, 66 p.
- Gibson, T. G., 1970, Late Mesozoic-Cenozoic tectonic aspects of the Atlantic coastal margin: Geological Society of America Bulletin, v. 81, no. 6, p. 1813-1822.
- Gitmez, G. U. and W. A. S. Sarjeant, 1972, Dinoflagellate cysts and acritarchs from the Kimmeridgian (Upper Jurassic) of England, Scotland and France: Bulletin of the British Museum of Natural History: Geology, v. 21, p. 171-257.
- Given, M. M., 1977, Mesozoic and Early Cenozoic geology of offshore Nova Scotia: Bulletin of Canadian Petroleum Geology, v. 25, p. 63-91.
- Gocht, H., 1970, Dinoflagellaten-Zysten aus dem Bathonium des erdolfeldes Aldorf (Northwest-Setuschland): Palaeontographica, Abt. B., v. 129, p. 125-165.
- Gorka, H., 1963, Coccolithophorides, Dinoflagellates, Hystrichosphaerides et microfossiles *incertae sedis* du Cretace superier de Pologne: Acta Palaeontologica Polonica, v. 8, p. 1-82.
- Gradstein, F.M., F.P.Achterberg, J.G. Ogg, J.Hardenbol, P. van Veen, and Z. Huang, 1995, A Triassic, Jurassic, and Cretaceous time scale; *in* Geochronology Time Scales and Stratigraphic Correlation, SEPM Special Publication no. 54, p. 95-126.
- Grose, P. L. and J. S. Mattson, 1977, The Argo Merchant oil spill--A preliminary scientific report: National Oceanic and Atmospheric Administration Environmental Research Laboratories, 129 p.
- Grow, J. A., R. E. Mattick, and J. S. Schlee, 1979, Multichannel seismic depth sections and interval velocities over continental shelf and upper continental slope between Cape Hatteras and Cape Cod, *in* Watkins, J. S., L. Montadert, and P. W. Dickerson, (eds.), Geological and Geophysical Investigations of Continental Margins: American Association of Petroleum Geologists Memoir 29, p. 65-83.
- Harwood, R. J., 1977, Oil and gas generation by laboratory pyrolysis of kerogen: American Association of Petroleum Geologists Bulletin, v. 61, no. 12, p. 2082-2102.
- Hill, M. E., III, 1976, Lower Cretaceous Nanofossils from Texas and Oklahoma: Paleontographica, Abteilung B, 156, Lfg. 4-6, p. 103-179.
- Hunt, J. M., 1967, The origin of petroleum in carbonate rocks: *in* G. V. Chilingar, H. S. Bissell, and R. W. Fairbridge, (eds.), Carbonate Rocks: New York, Elsevier, p. 225-251.
- Hunt, J. M., 1974, Hydrocarbon and kerogen studies, *in* C. C von der Borch and others, Initial Reports of the Deep Sea Drilling Project, v. 22: Washington, D. C., U. S. Government Printing Office, p. 673-675.
- Hunt, J. M., 1978, Characterization of bitumens and coals: American Association of Petroleum Geologists Bulletin, v. 62, no. 2, p. 301-303.
- Hunt, J. M., 1979, Petroleum Geochemistry and Geology: San Francisco, W. H. Freeman Co., p. 273-350.

- Hurtubise, D. O. and J. H. Puffer, 1985, Nepheline normative alkalic dolerite of the Georges Bank Basin, North Atlantic, part of an Early Cretaceous eastern North American alkalic province: Geological Society of America, Northeastern Section, 20th Annual Meeting, 1985, v. 17, no. 1, p. 25.
- Hurtubise, D. O., J. H. Puffer, and H. L. Cousminer, 1987, An offshore Mesozoic igneous sequence, Georges Bank Basin, North Atlantic: Geological Society of America Bulletin, v. 98, no. 4, p. 430-438.
- International Biostratigraphers, Inc., 1976, Biostratigraphy of the COST G-1 Georges Bank test: Houston, Texas, 16 p.
- International Biostratigraphers, Inc., 1977, Biostratigraphy of the COST G-2 Georges Bank test: Houston, Texas, 16 p.
- Jansa, L. F. and J. A. Wade, 1975, Geology of the continental margin off Nova Scotia and Newfoundland, *in* W. J. M van der Linden and J. A. Wade (eds.), Offshore Geology of Eastern Canada: Geological Survey of Canada Paper 74-30, v. 2, p. 51-105.
- Jansa, L. F. and J. Wiedmann, 1982, Mesozoic-Cenozoic development of the eastern North American and northwest African continental margins: a comparison, *in* V. von Rad, K. Hinz, M. Sarnthein, and E. Seibold (eds.), Geology of the Northwest African Continental Margin: Berlin, Springer-Verlag, p. 215-269.
- Jansa, L. F., G. L. Williams, J. A. Wade, and J. P. Bujak, 1978, COST B-2 well (Baltimore Canyon) and its relation to Scotian Basin (abstract): American Association of Petroleum Geologists Bulletin, v. 62, no. 3, p. 526.
- Jones, R. W. and T. A. Edison, 1978, Microscopic observations of kerogen related to geochemical parameters with emphasis on thermal maturation, *in* D. F. Oltz (ed.), Geochemistry: Low Temperature Metamorphism of Kerogen and Clay Minerals: Society of Economic Paleontologists and Mineralogists, Pacific Section, Annual Meeting, Los Angeles, p. 1-12.
- Kent, D. V. and F. M. Gradstein, 1986, A Jurassic to Recent chronology, *in* P. R. Vogt and B. E. Tucholke (eds.), The Geology of North America, vol. M, The Western North Atlantic Region: Geological Society of America, p. 45-50.
- King, L. H. and B. MacLean, 1975, Geology of the Scotian Shelf and adjacent areas: Canadian Geological Survey Paper 74-23, p. 22-53.
- Kinsman, D. J. J., 1975, Rift Valley basins and sedimentary history of trailing continental margins, *in* A. G. Fisher and S. Judson, (eds.), Petroleum and Global Tectonics: Princeton, Princeton University Press, p. 83-126.
- Kjellstrom, G., 1973, Maastrichtian microplankton from the Hollviken borehole No. 1 in Scania, southern Sweden: Sveriges Geologiska Undersokning, Afhandlingar och Uppsatser, v. 7, p. 1-59.
- Landes, K. K. 1967, Eometamorphism and oil and gas in time and space: American Association of Petroleum Geologists Bulletin, v. 51, no. 6, p. 828-841.
- LaPlante, R. E., 1974, Hydrocarbon generation in Gulf Coast tertiary sediments: American Association of Petroleum Geologists Bulletin, v. 58, no. 7, p. 1281-1289.

- Larskaga, Ye. S. and D. V. Zhabreu, 1964, Effects of stratal temperatures and pressures on the composition of dispersed organic matter (from the example of the Mesozoic-Cenozoic deposits of the Western Ciccaspian region): Dokl. Akad. Nauk SSSR, v. 157, no. 4, pp. 135-139.
- Lentin, J. K. and G. L. Williams, 1981, Fossil Dinoflagellates, Index to Genera and Species: Bedford Institute of Oceanography Report Series B1-R-81-12, p. 1-345.
- Louis, M. C. and B. P. Tissot, 1967, Influence de la temperature et de la pression sur la formation des hydrocarbures dans les argiles a kerogen [Influence of temperature and pressure on the generation of hydrocarbons in shales containing kerogen], *in* 7th World Petroleum Congress, Proceedings, (Mexico), v. 2: Chichester, International, John Wiley and Sons, p. 47-60.
- Lowell, J. D., G. J. Genik, T. H. Nelson, and P. M. Tucker, 1975, Petroleum and plate tectonics of the southern Red Sea, *in* A. G. Fisher and S. Judson, (eds.), Petroleum and Global Tectonics: Princeton University Press, Princeton, p. 129-153.
- McIver, N. L., 1972, Cenozoic and Mesozoic stratigraphy of the Nova Scotia shelf: Canadian Journal of Earth Sciences, v. 9, p. 54-70.
- MacLean, B.C., and J.A. Wade, 1992, Petroleum geology of the continental margin south of the islands of St. Pierre and Miquelon, offshore eastern Canada; Bulletin of Canadian Petroleum Geology, v. 40, no. 3, p. 222-253.
- Maher, J. C., 1971, Geologic Framework and Petroleum Potential of the Atlantic Coastal Plain and Continental Shelf: U. S. Geological Survey Professional Paper 659, 98 p.
- Martini, E., 1971, Standard Tertiary and Quaternary calcareous nannoplankton zonation *in* Proceedings II International Planktonic Conference, Roma, 1970: Rome, Edizioni Tecnoscienza, p. 739-785.
- Mattick, R. E., R. Q. Foote, N. L. Weaver, and M. S. Grim, 1974, Structural framework of United States Atlantic Outer Continental Shelf north of Cape Hatteras: American Association of Petroleum Geologists Bulletin, v. 58, no. 6, 1179-1190.
- Miller, R. E., H. E. Lerch, G. E. Claypool, M. A. Smith, D. K. Owings, D. T. Lignon, and S. B. Eisner, 1982, Organic geochemistry of the Georges Bank basin COST Nos. G-1 and G-2 wells, *in* P. A. Scholle and C. R. Wenkam (eds.), Geological Studies of the COST Nos. G-1 and G-2 Wells, Unites States North Atlantic Outer Continental Shelf: U. S. Geological Survey Circular 861, p. 105-142.
- Miller, R. E., R. E. Mattick, and H. E. Lerch, 1981, Petroleum geochemistry and geology of Cenozoic and Mesozoic sedimentary rocks from Georges Bank basin (abstract): American Association of Petroleum Geologists Bulletin, v. 65, no. 9, p. 1667.
- Miller, R. E., D. M. Schultz, G. E. Claypool, H. E. Lerch, D. T. Lignon, C. Gary, and D. K. Owings, 1979, Organic geochemistry, *in* , P. A. Scholle (ed.), Geological Studies of the COST GE-1 Well, United States South Atlantic Outer Continental Shelf Area: U. S. Geological Survey Circular 800, p. 74-92.
- Miller, R. E., D. M. Schultz, G. E. Claypool, M. A. Smith, H. E. Lerch, D. Ligon, D. K. Owings, and C. Gary, 1980, Organic geochemistry, *in* P.A. Scholle (ed.), Geological Studies of the COST No. B-3 Well, United States Mid-Atlantic Continental Slope Area: U. S. Geological Survey Circular 833, p. 85-104.
- Miller, R. E., D. M. Schultz, H. E. Lerch, D. T. Lignon, and P. C. Bowker, 1986, *in* Edson, G. M.(ed.), Shell Wilmington Canyon 586-1 Well, Geological and Operation Summary: Minerals Management Service, OCS Report MMS 86-0099, p. 37-44.

- Miller, R. E., D. M. Schultz, H. E. Lerch, D. T. Lignon, and P. C. Bowker, 1987, *in* Edson, G. M. (ed.), Shell Wilmington Canyon 587-1 Well, Geological and Operation Summary: Minerals Management Service, OCS Report MMS 87-0074, p. 39-46.
- Momper, J. A., 1978, Oil migration limitations suggested by geological and geochemical considerations, *in* Physical and Chemical Constraints on Petroleum Migration: American Association of Petroleum Geologists, Continuing Education Course Note Series No., 8, p. B1-B60.
- Morbey, S. J., 1975, The palynostratigraphy of the Rhaetian Stage Upper Triassic in the Kerdelbachgraben Austria: *Paleontographica Abt. B*, v. 152, p. 1-75, p. 1-19.
- Murray, G. E., 1961, *Geology of the Atlantic and Gulf Coastal Provinces of North America*: New York, Harper, 692 p.
- Orr, W. L., 1974, Changes in sulfur content and isotopic ratios of sulfur during petroleum maturation--study of Big Horn Basin Paleozoic oils: *American Association Petroleum Geologists Bulletin*, v. 58, no. 11, p. 2295-2318.
- Perry, W. J., J. P. Minard, E. G. A. Weed, E. I. Robbins, and E. C. Rhodehamel, 1975, Stratigraphy of the Atlantic continental margin of the United States north of Cape Hatteras--brief survey: *American Association of Petroleum Geologists Bulletin*, v. 59, no. 9, p. 1529-1548.
- Phillipi, G. T., 1957, Identification of oil-source beds by chemical means, *in* 20th International Geological Congress Proceedings: Mexico City (1956), Sec. 3, p. 25-38.
- Phillipi, G.T., 1965, On the depth, time, and mechanism of petroleum generation: *Geochim. Cosmochim. Acta*, v. 29, p. 1021.
- Postuma, J. A., 1971, *Manual of Planktonic Foraminifera*: New York, Elsevier, 420 p.
- Pusey, W. C., III, 1973, The ESR-kerogen method--how to evaluate potential gas and oil source rocks: *World Oil*, v. 176, no. 5, p. 71-75.
- Reinhardt, P., 1966, Zur taxonomie und biostratigraphie des fossilen nannoplanktons aus dem Malm, der Kreide und dem Alttertiar Mitteleuropas [Taxonomy and biostratigraphy of Malm, Cretaceous, and early Tertiary nannoplanktonic faunas of central Europe], *Frieberger Forschungshefte, Reihe C: Geowissenschaften, Mineralogie-Geochemie*, 196 Paleont.: Leipzig, Bergakademie Freiberg, p. 5-61.
- Ricciardi, K. (ed.), 1989, Exxon Lydonia Canyon 133-1 Well, Geological and Operational Summary: Minerals Management Service OCS Report MMS 89-0007, 46 p.
- Riding, J. B., 1984, Dinoflagellate cyst range-top biostratigraphy of the uppermost Triassic to lowermost Cretaceous of northwest Europe: *Palynology*, v.8, p. 195-210.
- Robbins, E. I. and E. C. Rhodehamel, 1976, Geothermal gradients help predict petroleum potential of Scotian Shelf: *Oil & Gas Journal*, v. 74, no. 9, p. 143-145.
- Rona, P. A., 1973, Relations between rates of sediment accumulation on continental shelf, sea-floor spreading, and eustasy inferred from central North Atlantic: *Geological Society of America Bulletin*, v. 84, no. 9, p. 2851-2872.
- Ryan, W. B. F., M. B. Cita, R. L. Miller, D. Hanselman, B. Hecker, and M. Nibbelink, 1978, Bedrock geology in New England submarine canyons: *Oceanologia Acta*, v. 1, no. 2, p. 233-254.



- Sarjeant, W. A. S., 1979, Middle and Upper Jurassic dinoflagellate cysts--the world excluding North America: American Association of Stratigraphic Palynologists Contributions Series no. 5-B, p. 133-157.
- Schlee, J. S., J. C. Behrendt, J. A. Grow, J. M. Robb, R. E. Mattick, P. T. Taylor, and B. J. Lawson, 1976, Regional geologic framework off northeastern United States: American Association of Petroleum Geologists Bulletin, v. 60, no. 6, p. 926-951.
- Schlee, J. S., W. P. Dillon, and J. A. Dillon, 1979, Structure of the continental slope off the eastern United States, *in* L. J. Doyle and O. H. Pilkey, (eds.), Geology of Continental Slopes: Society of Economic Paleontologists and Mineralogists Special Publication 27, p. 95-117.
- Schlee, J.S. and K.D. Klitgord, 1988, Georges Bank basin: a regional synthesis; *in* R.E. Sheridan and J.A. Grow (eds.), The Geology of North America, vol. I-2, The Atlantic Continental Margin, Geological Society of America, p. 243-268.
- Schlee, J. S., R. G. Martin, R. E. Mattick, W. P. Dillon, and M. M. Ball, 1977, Petroleum geology of the U. S. Atlantic--Gulf of Mexico margins, *in* V. S. Cameron (ed.), Exploration and Economics of the Petroleum Industry--New Ideas, Methods, New Developments: Southwestern Legal Foundation: New York, Mathew Bender and Co., v. 15, p. 47-93.
- Schlee, J. S., R. E. Mattick, D. J. Taylor, O. W. Girard, E. C., Rhodehamel, W. J. Perry, and K. C. Bayer, 1975, Sediments, structural framework, petroleum potential, environmental conditions and operation considerations of the United States North Atlantic Outer Continental Shelf: U. S. Geological Survey, Open-File Report 75-353, 179 p.
- Scholle, P. A. and C. R. Wenkam (eds.), 1982, Geological studies of the COST Nos. G-1 and G-2 wells, United States North Atlantic OCS: U. S. Geological Survey Circular 861, 193 p.
- Schultz, L. K. and R. L. Grover, 1974, Geology of Georges Bank Basin: American Association of Petroleum Geologists Bulletin, v. 58, no. 6, p. 1159-1168.
- Schwab, K.W., P. van Gijssel, and M.A. Smith, 1990, Kerogen evolution and microscopy workshop short course, International Symposium on Organic Petrology, Zeist, the Netherlands, January 10 and 11, 1990 (unpublished).
- Shell Canada Limited, 1970a, Well history report, Oneida O-25, 50 p.
- Shell Canada Limited, 1970b, Well history report, Mohawk B-93, 25 p.
- Shell Canada Limited, 1972, Well history report, Mohican I-100, 76 p.
- Sheridan, R. E., 1974a, Conceptual model for the block-fault origin of the North American Atlantic continental margin geosyncline: Geology, v. 2, no. 9, p. 465-468.
- Sheridan, R. E., 1974b, Atlantic continental margin of North America, *in* C. A. Burk and C. L. Drake, (eds.), Geology of Continental Margins: New York, Springer-Verlag, p. 391-407.
- Sheridan, R. E., 1976, Sedimentary basins of the Atlantic margin of North America: Tectonophysics, v. 36, p. 113-132.
- Sherwin, D. F., 1973, Scotian Shelf and Grand Banks, *in* R. G. McCrossan (ed.), Future Petroleum Provinces of Canada--Their Geology and Potential: Canadian Society of Petroleum Geologists Memoir 1, p. 519-559.

- Singh, C., 1971, Lower Cretaceous microfloras of the Peace River area, northwestern Alberta: Research Council of Alberta Bulletin 28, 2 volumes, 542 p.
- Smith, H. A., 1975, Geology of the West Sable structure: Bulletin of Canadian Petroleum Geology, v. 23, no. 1, p. 109-130.
- Smith, M. A., 1979, Geochemical analysis, *in* R. V. Amato and E. K. Simonis (eds.), Geologic and Operational Summary, COST No. B-3 Well, Baltimore Canyon Trough Area, Mid-Atlantic OCS: U. S. Geological Survey Open-File Report 79-1159, p. 81-99.
- Smith, M. A., 1980, Geochemical analysis, *in* R.V. Amato and E.K. Simonis (eds.), Geologic and Operational Summary, COST No. G-2 Well, Georges Bank Area, North Atlantic OCS: U. S. Geological Survey Open-File Report, 80-269, p. 77-99.
- Smith, M. A., 1995, Assessment of U.S. Atlantic hydrocarbon resources using new geochemical technology: U.S. Geological Society of America, Abstracts with programs, 1995 Annual Meeting, New Orleans, LA.
- Smith, M.A., R.V. Amato, M.A. Furbush, D.M. Pert, M.E. Nelson, J. S. Hendrix, L.C. Tamm, G. Wood, Jr., and D.R. Shaw, 1976, Geological and Operational Summary, COST No. B-2 Well, Baltimore Canyon Trough Area, Mid-Atlantic OCS: U. S. Geological Survey Open-File Report 76-774, 79 p.
- Smith, M. A. and D. R. Shaw, 1980, Geochemical analysis, *in* R. V. Amato and J. W. Bebout (eds.), Geologic and Operational Summary, COST No. G-1 well, Georges Bank Area, North Atlantic OCS: U. S. Geological Survey Open-File Report 80-268, p. 81-94.
- Smith, M.A., and P. van Gijzel, 1990, New perspectives on the depositional and thermal history of Georges Bank; *in* W.J.J. Fermont and J.W. Weegink (eds.), Proceedings, International Symposium on Organic Petrology, Zeist, the Netherlands.
- Smith, R. A., J. R. Stack, and R. K. Davis, 1976, An oil spill risk analysis for the Mid-Atlantic Outer Continental Shelf lease area: U. S. Geological Survey Open-File Report 76-451, 24 p.
- Staplin, F. L., 1969, Sedimentary organic matter, organic metamorphism, and oil and gas occurrence: Bulletin of Canadian Petroleum Geology, v. 17, no. 1, p. 47-66.
- Steinkraus, W. E., 1980, Biostratigraphy, *in* R. V. Amato and J. W. Bebout, (eds.), Geologic and Operation Summary, COST No. G-1 Well, Georges Bank, North Atlantic OCS: U. S. Geological Survey Open-File Report 80-268, p. 39-51.
- Stewart, H. B., Jr. and G. F. Jordan, 1964, Underwater sand ridges on Georges Shoal, *in* R. L. Miller (ed.), Papers in Marine Geology, Shepard Commemorative Volume: New York, Macmillan, p. 102-114.
- Tamm, L. C., 1978, Electric log interpretations, *in* R. V. Amato and J. W. Bebout (eds.), Geological and Operational Summary, COST No. GE-1 Well, Southeast Georgia Embayment Area, South Atlantic OCS: U. S. Geological Survey Open-File Report 78-668, 61-75.
- Thierstein, H. R., 1971, Tentative Lower Cretaceous calcareous nannoplankton zonation: *Eclogae Geologicae Helveticae*, v. 64, p. 459-487.
- Tissolt, B. P. and D. H. Welte, 1978, Petroleum Formation and Occurrence, A New Approach to Oil and Gas Exploration: Berlin, Springer-Verlag, p. 123-201.

- Tissot, B., B. Durand, J. Espitalie, and A. Combaz, 1974, Influence of nature and diagenesis of organic matter in formation of petroleum: American Association of Petroleum Geologists Bulletin, v. 58, no. 3, p. 499-506.
- Tschudy, R. H., 1973, *Complexiopollis* Pollen Lineage in Mississippi Embayment Rocks: U. S. Geological Survey Professional Paper 743-C, p. C1-C15.
- Uchupi E. and K. O. Emery, 1967, Structure of continental margin off Atlantic coast of United States: American Association of Petroleum Geologists Bulletin, v. 51, no. 2, p. 223-234.
- U. S. Department of Commerce, 1973, Environmental Conditions within Specified Geographical Regions-- Offshore East and West Coast of the United States and in the Gulf of Mexico: Washington, D. C., National Oceanographic Data Center, National Oceanographic and Atmospheric Administration, 735 p.
- Van Gijzel, P., 1990, Transmittance colour index (TCI) of amorphous organic matter: a new thermal maturity indicator for hydrocarbon source rocks, and its correlation with mean vitrinite reflectance and thermal alteration index (TAI); in W.J.J. Fermont and J.W. Weegink, eds., Proceedings, International Symposium on Organic Petrology, Zeist, the Netherlands.
- Van Hinte, J. E., 1976a, A Jurassic time scale: American Association of Petroleum Geologists Bulletin, v. 60, no. 4, p. 489-497.
- Van Hinte, J. E., 1976b, A Cretaceous time scale: American Association of Petroleum Geologists Bulletin, v. 60, no. 4, p. 498-516.
- Vassoyevich, N. B., Yu. I. Korchagina, N. V. Lopatin, and V. V. Chernyshev, 1969, Glavanaya faza nefteobrazovaniya [Principal phase of oil formation]: Moskovskogo Universiteta Vestnik, Ser. 4, Geologii, v. 24, no. 6, p. 3-27; English translation in International Geology Review, 1970, v. 12, no. 11, p. 1,276-1,296.
- Wade, J.A., 1977, Stratigraphy of Georges Bank Basin-- interpretation from seismic correlation to the western Scotian Shelf: Canadian Journal of Earth Science, v. 14, no. 10, p. 2274-2283.
- Wade, J.A., G.R.Campbell, R.M. Proctor, and G.C. Taylor, 1989, Petroleum Resources of the Scotian Shelf, Geological Survey of Canada Paper 88-19.
- Walper, J. L. and R. E. Miller, 1985, Tectonic evolution of Gulf Coast basins, in B. F. Perkins and G. B. Martin (eds.), Habitat of Oil and Gas, Program and Abstracts, Fourth Annual Research Conference, Gulf Coast Section: Austin, Society of Economic Paleontologists and Mineralogists Foundation, Earth Enterprises, p. 25-42.
- Waples, D. W., 1980, Time and temperature in petroleum formation--application of Lopatin's method to petroleum exploration: American Association of Petroleum Geologists Bulletin, v. 64, no. 6, p. 916-926.
- Weed, E. G. A., J. P. Minard, W. J. Perry, Jr., E. C. Rhodehamel, and E. I. Robbins, 1974, Generalized pre-Pleistocene geologic map of the northern United States Atlantic continental margin: U. S. Geological Survey Miscellaneous Geologic Investigations Map I-861, Scale 1:1,000,000.
- Williams, G. L., 1974, Dinoflagellate and spore stratigraphy of the Mesozoic-Cenozoic offshore Eastern Canada, in Offshore Geology of Eastern Canada: Geological Survey of Canada Paper 74-30, v. 2, p. 107-161.
- Williams, G. L., 1977, Dinocysts--their classification, Biostratigraphy, and paleoecology, in A. T. S. Ramsay (ed.), Oceanic Micropaleontology, v. 2, New York, Academic Press, p. 1,231-1,326.

Williams, G. L. and W. W. Brideaux, 1975, Palynologic analyses of Upper Mesozoic and Cenozoic rocks of the Grand Banks, Atlantic Margin: Geological Survey of Canada Bulletin, v. 236, p. 1-163.

Woollam, R. and J. B. Riding, 1983, Dinoflagellate cyst zonation of the English Jurassic: Institute of Geological Sciences Report, v. 83, No. 2, p. 1.

Worsley, T. R., 1971, Calcareous nannofossil zonation of Upper Jurassic and Lower Cretaceous sediments from the Western Atlantic, *in* Proceedings II, International Planktonic Conference, Roma, 1970: Rome, Edizioni Tecnoscienza, p. 1301-1321 .