Chapter 8. References Cited

- Able, K.W., and McAllister, D.E., 1980, Revision of the snailfish genus *Liparis* from Arctic Canada: Ottawa, Department of Fisheries and Oceans, Canadian Bulletin of Fisheries and Aquatic Sciences 208.
- Abookire, A.A., Piatt, J.F., and Norcross, B.L., 2001, Juvenile groundfish habitat in Kachemak Bay, Alaska, during late summer: Alaska Fishery Research Bulletin, v. 8, no. 1, p. 45–56.
- Abookire, A.A., Piatt, J.F., and Robards, M.D., 2000, Nearshore fish distributions in an Alaska estuary in relation to stratification, temperature and salinity: Estuarine, Coastal and Shelf Science, v. 51, no. 1, p. 45–59.
- Acuna, E.I., Goddard, P., and Kotwicki, S., 2003, 2002 bottom trawl survey of the eastern Bering Sea continental shelf: National Marine Fisheries Service, Alaska Fisheries Science Center, Processed Report 2003-01.
- Ahlstrom, E.H., Amaoka, K., Hensley, D.A., Moser, H.G., and Sumida, B.Y., 1984, Pleuronectiformes—Development, Moser, H.G., and others, eds., *in* Ontogeny and systematics of fishes: American Society of Ichthyologists and Herpetologists.
- Akaike, H., 1973, Information theory and an extension of the maximum likelihood principle, *in* Petrov, B.N., and Caski, F., eds., Proceedings of the Second International Symposium on Information Theory: Budapest, Akademiai Kiado, p. 267–281.
- Alaska Fisheries Science Center, 2013, Ichthyoplankton Information System: Alaska Fisheries Science Center, accessed November 29, 2013, at http://access.afsc.noaa.gov/ ichthyo/index.cfm.
- Albers, W.D., and Anderson, P.J., 1985, Diet of Pacific cod, *Gadus macrocephalus*, and predation on the northern pink shrimp, *Pandalus borealis*, in Pavlof Bay, Alaska: Fishery Bulletin, v. 83, no. 4, p. 601–610.
- Albert, O.T., Harbitz, A., and Høines, Å.S., 2003, Greenland halibut observed by video in front of survey trawl— Behaviour, escapement, and spatial pattern: Journal of Sea Research, v. 50, no. 2–3, p. 117–127.
- Alexander, V., 1992, Arctic marine ecosystems, chap. 17 *of*, Peters, R.L., and Lovejoy, T.L., eds., Global warming and biological diversity: New Haven, Conn., Yale University Press, p. 221–232.

- Allen, M.J., and Smith, G.B., 1988, Atlas and zoogeography of common fishes in the Bering Sea and northeast Pacific: U.S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Report NMFS 66, 151 p.
- Allen, M.S., and Hightower, J.E., 2010, Fish population dynamics—Mortality, growth, and recruitment, *in* Hubert, W.A., and Quist, M.C., eds., Inland fisheries management in North America: Bethesda, Md., American Fisheries Society, p. 43–79.
- Alt, K.T., 1969, Taxonomy and ecology of the inconnu, *Stenodus leucichthys nelma*, in Alaska: Anchorage, University of Alaska, Biological Papers of the University of Alaska, no. 12, 63 p.
- Alt, K.T., 1973b, Age and growth of the inconnu (*Stenodus leucichthys*) in Alaska: Journal of the Fisheries Research Board of Canada, v. 30, no. 1, p. 457–459.
- Alt, K.T., 1973a, Contributions to the biology of the Bering cisco (*Coregonus laurettae*) in Alaska: Journal of the Fisheries Research Board of Canada, v. 30, no. 12, p. 1,885–1,888.
- Alt, K.T., 1987, Review of sheefish (*Stenodus leucichthys*) studies in Alaska: Juneau, Alaska Department of Fish and Game, Division of Sport Fish, Fishery Manuscript no. 3, 69 p.
- Alt, K.T., 1988, Biology and management of inconnu (*Stenodus leucichthys*) in Alaska: Finnish Fisheries Research, v. 9, p. 127–132.
- Alt, K.T., 2008, Alaska wildlife notebook series—Sheefish: Alaska Department of Fish and Game.
- Alton, M.S., Bakkala, R.G., Walters, G.E., and Munro, P.T., 1988, Greenland turbot *Reinhardtius hippoglossoides* of the eastern Bering Sea and Aleutian Island region: National Oceanic and Atmospheric Administration, Technical Report NMFS 71, 31 p.
- Alverson, D.L., 1960, A study of annual and seasonal bathymetric catch patterns for commercially important groundfishes of the Pacific Northwest coast of North America: Pacific Marine Fisheries Commission Bulletin 4, 66 p.

Anderson, P.J., and Piatt, J.F., 1999, Community reorganization in the Gulf of Alaska following ocean climate regime shift: Marine Ecology Progress Series, v. 189, p. 117–123.

Anderson, R.O., and Gutreuter, S.J., 1983, Length, weight, and associated structural indices, *in* Nielsen, L.A., and Johnson, D.L., eds., Fisheries Techniques (2d ed.):Bethesda, Md., American Fisheries Society, p. 283–300.

Andriyashev, A.P., 1954, Fishes of the northern seas of the U.S.S.R,—Keys to the fauna of the U.S.S.R.: Academy of Sciences of the U.S.S.R., Zoological Institute, no. 53, 566 p. [In Russian, translation by Israel Program for Scientific Translation, Jerusalem, 1964, 617 p., available from U.S. Department of Commerce, Springfield, Virginia].

Andriyashev, A.P., 1961, Forms of the hook-horn sculpin genus *Artediellus* Jordan (Pisces: Cottidae) in the Bering Sea: Vopr Ikhtiol, v. 1, p. 231–242. [In Russian.]

Andriyashev, A.P., 1970, Cryopelagic fishes of Arctic and Antarctic and their significance in polar ecosystems, *in* Holdgate, M.W., ed., Antarctic ecology: New York, Academic Press, p. 297–304.

Andriashev, A.P., 1986, Agonidae, *in* Whitehead, P.J.P., ed., Fishes of the north-eastern Atlantic and the Mediterranean: Paris, United Nations Educational, Scientific, and Cultural Organization, p. 1,265–1,268.

Andriyashev A.P., and Chernova, N.V., 1994, Annotated list of fishlike vertebrates and fish of the Arctic seas and adjacent waters: Journal of Ichthyology, v. 35, no. 1, p. 81–123.

Anisimov, O.A., Vaughan, D.G., Callaghan, T.V., Furgal, C., Marchant, H., Prowse, T.D., Vilhjálmsson, H., and Walsh, J.E., 2007, Polar regions (Arctic and Antarctic), *in* Parry, M.L., Canziani, O.F., Palutikof, J.P., van der Linden, P.J., and Hanson, C.E., eds., Climate change 2007—Impacts, adaptation and vulnerability, contribution of working group II to the fourth assessment report of the Intergovernmental Panel on Climate Change: Cambridge, U.K., Cambridge University Press, p. 653–685.

Ansaldo, M., Luquet, C.M., Evelson, P.A., Polo, J.M., and Llesuy, S., 2000, Antioxidant levels from different Antarctic fish caught around South Georgia Island and Shag Rocks: Polar Biology, v. 23, no. 3, p. 160–165.

Antonenko, D.V., and Vdovin, A.V., 2001, Seasonal distribution of the common greenling *Hexagrammos stelleri* (Hexagrammidae) in Peter the Great Bay (the Sea of Japan): Journal of Ichthyology, v. 41, p. 524–528.

Archer, S.D., and Johnston, I.A., 1989, Kinematics of labriform and subcarangiform swimming in the Antarctic fish *Notothenia neglecta*: The Journal of Experimental Biology, v. 143, p. 195–210. Armstrong, R.H., Carstensen, R.L., and Wilson, M.F., 2004, Hotspots—Bird survey of Mendenhall Wetlands April 2002 to May 2003: Juneau Audubon Society and Taku Conservation Society, 9 p.

Armstrong, R.H., and Morrow, J.E., 1980, The Dolly Varden charr, *Salvelinus malma*, *in* Balon, E.K., ed., Charrs, salmonid fishes of the genus *Salvelinus*: The Hague, Netherlands, Dr. W. Junk B.V. Publishers, p. 99–140.

Aronovich, T.M., Doroshev, S.I., Spectorova, L.V., and Makhotin, V.M., 1975, Egg incubation and larval rearing of navaga (*Eleginus navaga* Pall.), Arctic Cod (*Boreogadus saida* lepechin) and Arctic flounder (*Liopsetta glacialis* Pall.) in the laboratory: Aquaculture, v. 6, no. 3, p. 233–242.

Arvey, W.D., 1991, Stock status of anadromous Dolly Varden in waters of Alaska's North Slope: Alaska Department of Fish and Game, Fishery Manuscript no. 91-3, 20 p.

Aschan, M., Karamushko, O.V., Byrkjedal, I., Wienerroither, R., Borkin, I.V., and Christiansen, J.S., 2009, Records of the gadoid fish *Arctogadus glacialis* (Peters, 1874) in the European Arctic: Polar Biology, v. 32, no. 7, p. 963–970.

Atkinson, E.G., and Percy, J.A., 1992, Diet comparison among demersal marine fish from the Canadian Arctic: Polar Biology, v. 11, no. 8, p. 567–573.

Axelsson, M., 2005, The circulatory system and its control, *in* Farrell, A.P., and Steffensen, J.F., eds., The physiology of polar fishes: Amsterdam, Elsevier, p. 239–280.

Aydin, K., Gaichas, S., Ortiz, I., Kinzey, D., and Friday, N., 2007, A comparison of the Bering Sea, Gulf of Alaska, and Aleutian Islands large marine ecosystems through food web modeling: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, Alaska Fisheries Science Center, NOAA Technical Memorandum NMFS-AFSC-178, 309 p.

Bailey, K.M., Abookire, A.A., and Duffy-Anderson, J.T., 2008, Ocean transport paths for the early life history stages of offshore-spawning flatfishes—A case study in the Gulf of Alaska: Fish and Fisheries, v. 9, no. 1, p. 44–66.

Bailey, K.M., Brown, E.S., and Duffy-Anderson, J.T., 2003, Aspects of distribution, transport and recruitment of Alaska plaice (*Pleuronectes quadrituberculatus*) in the Gulf of Alaska and eastern Bering Sea: comparison of marginal and central populations: Journal of Sea Research, v. 50, no. 2–3, p. 87–95.

Bailey, K.M., and Picquelle, S.J., 2002, Larval distribution of offshore spawning flatfish in the Gulf of Alaska—Potential transport pathways and enhanced onshore transport during ENSO events: Marine Ecology Progress Series, v. 236, p. 205–217. Bailey, K.M., Powers, D.M., Quattro, J.M., Villa, G., Nishimura, A., Traynor, J.J., and Walters, G., 1999, Population ecology and structural dynamics of Walleye Pollock (*Theragra chalcogramma*), *in* Loughlin, T.R., and Ohtani, K., eds., Dynamics of the Bering Sea: Fairbanks, University of Alaska Sea Grant, AK-SG-99-03, p. 581–613.

Baird, P.A., and Gould, P., 1983, The breeding biology and feeding ecology of marine birds in the Gulf of Alaska: U.S.
Fish and Wildlife Service, Denver Wildlife Research Center, Final Report, Outer Continental Shelf Environmental Assessment Program, Research Unit 341, Final Report 45 (1986), p. 1–120.

Bakkala, R.G., 1970, Synopsis of biological data on the chum salmon, *Oncorhynchus keta* (Walbaum, 1792): U.S. Fish and Wildlife Service, FAO Fisheries Synopsis no. 41, 89 p.

Bakkala, R.G., 1981, Population characteristics and ecology of yellowfin sole, *in* Hood, D.W., and Calder, J.A., eds.: The eastern Bering Sea shelf—Oceanography and resources: National Oceanic and Atmospheric Administration, p. 553–574.

Balanov, A.A., Badaev, O.Z., Napazakov, V.V., and Chuchukalo, V.I., 2006, Distribution and some biological features of *Lycodes raridens* (Zoarcidae) in the western part of the Bering Sea: Journal of Ichthyology, v. 46, no. 2, p. 148–155.

Barber, W.E., Smith, R.L., Vallarino, M., and Meyer, R.M., 1997, Demersal fish assemblages of the northeastern Chukchi Sea, Alaska: Fishery Bulletin, v. 95, no. 2, p. 195–209.

Barber, W.E., Smith, R.L., and Weingartner, T.J., 1994, Fisheries oceanography of the northeast Chukchi Sea: Final Report to U.S. Department of the Interior, Mineral Management Service, Anchorage, Alaska, OCS Study MMS-93-0051 [variously paged].

Beamish, R.J., 1980, Adult biology of the river lamprey (*Lampetra ayresi*) and the Pacific lamprey (*Lampetra tridentata*) from the Pacific Coast of Canada: Canadian Journal of Fisheries and Aquatic Sciences, v. 37, no. 11, p. 1,906–1,923.

Beamish, R.J., McFarlane, G.A., and King, J.R., 2005, Migratory patterns of pelagic fishes and possible linkages between open ocean and coastal ecosystems off the Pacific coast of North America: Deep-Sea Research II, v. 52, p. 739–755.

Bean, T.H., 1887, The fishery resources and fishing-grounds of Alaska, *in* Goode, G.B., ed., The fisheries and fishery industries of the United States, Section III: United States Commission of Fish and Fisheries, p. 81–115. Beauchamp, D.A., Shepard, M.F., and Pauley, G.B., 1983, Species profiles—Life histories and environmental requirements of coastal fishes and invertebrates (Pacific Northwest)—Chinook Salmon: U.S. Fish and Wildlife Service, Biological Services Program FWS/OBS 82 (11.6), 15 p.

Becker, K., Wöhrmann, A.P.A., and Rahmann, H., 1995, Brain gangliosides and cold-adaptation in high-Antarctic fish:
Biochemical Systematics and Ecology, v. 23, nos. 7–8, p. 695–707.

Belyanina, T.N., 1968, Dynamics of smelt populations in sub-arctic waters: Rapports et procès-verbaux des réunions/ Conseil permanent international pour l'exploration de la mer, v. 158, p. 74–79.

Bendock, T.N., 1977, Beaufort Sea estuarine fishery study: Alaska Department of Fish and Game Annual Report, Contract #03-5-022-69, p. 670–729.

Benoit, D., Simard, Y., and Fortier, L., 2008, Hydroacoustic detection of large winter aggregations or Arctic cod (*Boreogadus saida*) at depth in ice-covered Franklin Bay (Beaufort Sea): Journal of Geophysical Research, v. 113, no. CO6S90, 9 p.

Benoit, D., Simard, Y., Gagné, J., Geoffroy, M., and Fortier, L., 2010, From polar night to midnight sun—Photoperiod, sea predation, and the diel vertical migrations of polar cod (*Boreogadus saida*) under landfast ice in the Arctic Ocean: Polar Biology, v. 33, no. 11, p. 1,505–1,520.

Benton, T.G., Plaistow, S.J., and Coulson, T.N., 2006, Complex population dynamics and complex causation— Devils, details, and demography: Proceedings of the Royal Society B, Biological Sciences, v. 273, no. 1591, p. 1,173–1,181.

Benz, G.W., Hocking, R., Kowunna, A., Bullard, S.A., and George, J.C., 2004, A second species of Arctic shark— Pacific sleeper shark (*Somniosus pacificus*) from Point Hope, Alaska: Polar Biology, v. 27, no. 4, p. 250–252.

Benziger, D., and Umminger, B.L., 1973, Role of hepatic glycogenolytic enzymes in the cold-induced hyperglycemia of the killifish, *Fundulus heteroclitus*: Comparative Biochemistry and Physiology, v. 45, no. 3, p. 767–772.

Best, E.A., 1981, Halibut ecology, *in* Hood, D.W., and Calder, J.A., eds., The eastern Bering Sea shelf—Oceanography and resources: Seattle, University of Washington Press, p. 495–508.

Bjørkvoll, E., Grøtan, V., Aanes, S., Sæther, B.E., Engen, S., and Aanes, R., 2012, Stochastic population dynamics and life-history variation in marine fish species: American Naturalist, v. 180, no. 3, p. 372–387.

Blackburn, J.E., and Jackson, P.B., 1982, Seasonal composition and abundance of juvenile and adult marine finfish and crab species in the nearshore zone of Kodiak Island's eastside during April 1978 through March 1979: Outer Continental Shelf Environmental Assessment Program, Alaska Department of Fish and Game, Final Report, Research Unit 552, p. 377–570.

Blackburn, J.E., Jackson, P.B., Warner, I.M., and Dick, M.H., 1981, A survey for spawning forage fish on the east side of the Kodiak Archipelago by air and boat during spring and summer 1979: Outer Continental Shelf Environmental Assessment Program, Final Reports of Principal Investigators, RU 0552, p. 309–376.

Blood, D.M., Duffy-Anderson, J.T., Matarese, A.C., Sohn,
D., Stabeno, P.J., and Cheng, W., 2009, Early life history of Greenland halibut (*Reinhardtius hippoglossoides*) in the eastern Bering Sea based on recent field studies (2007–2009)—Spawning, distribution, and physical processes affecting drift and dispersal: Portland, Oregon, 33rd Annual Larval Fish Conference and American Society of Ichthyologists and Herpetologists.

Bluhm, B., Dunton, K., Grebmeier, J., and Sirenko, B., 2008, Benthos, *in* Hopcroft, R., Bluhm, B., and Gradinger, R., eds., Arctic Ocean synthesis—Analysis of climate change impacts in the Chukchi and Beaufort Seas with strategies for future research: Institute of Marine Sciences, University of Alaska Fairbanks, p. 56–65.

Bluhm, B.A., Gebruk, A.V., Gradinger, R., Hopcroft, R.R., Huettmann, F., Kosobokova, K.N., Sirenko, B.I., and Weslawski, J.M., 2011, Arctic marine biodiversity—An update of species richness and examples of biodiversity change: Oceanography, v. 24, no. 3, p. 232–248.

Bluhm, B.A., and Gradinger, R., 2008, Regional variability in food availability for Arctic marine mammals: Ecological Applications, v. 18, no. 2 Supplement, p. S77-S96.

Bond, W.A., 1982, A study of the fishery resources of Tuktoyaktuk Harbour, southern Beaufort Sea coast, with special reference to life histories of anadromous coregonids: Canadian Technical Report of Fisheries and Aquatic Sciences, no. 1119, 90 p.

Bond, W.A., and Erickson, R.N., 1987, Fishery data from Phillips Bay, Yukon, 1985: Winnipeg, Manitoba, Canadian Data Report of Fisheries and Aquatic Sciences, Central and Artcic Region, Department of Fisheries and Oceans, no. 635, 47 p.

Bond, W.A., and Erickson, R.N., 1989, Summer studies of the nearshore fish community at Phillips Bay, Beaufort Sea coast, Yukon: Winnipeg, Manitoba, Canadian Technical Report of Fisheries and Aquatic Sciences, Central and Arctic Region, Department of Fisheries and Oceans, no. 1676, 102 p. Bond, W.A., and Erickson, R.N., 1993, Fisheries investigations in coastal waters of Liverpool Bay, Northwest Territories: Winnipeg, Manitoba, Canada Department of Fisheries and Oceans, Central and Arctic Region, Canadian Manuscript Report of Fisheries and Aquatic Sciences No. 2204, 59 p.

Bonham, K., 1954, Food of the dogfish *Squalus acanthias*: Washington Department of Fisheries, v. 1, no. 2, p. 25–36.

Booth, S., Zeller, D., and Pauly, D., 2008, Baseline study of marine fisheries catches from Arctic Alaska 1950–2006: Report Fisheries Center, University of British Columbia, Canada, 16 p.

Bradstreet, M.S.W., 1982, Occurrence, habitat use, and behavior of seabirds, marine mammals, and Arctic cod at the Pond Inlet ice edge: Arctic, v. 35, no. 1, p. 28–40.

Bradstreet, M.S.W., Sekerak, A.D., Finley, K.J., Griffiths, W.B., Evans, C.R., Fabijan, M.F., and Stallard, H.E., 1986, Aspects of the biology of Arctic cod (*Boreogadus saida*) and its importance in Arctic marine food chains: Canada Department of Fisheries and Oceans, Canadian Technical Report of Fisheries and Aquatic Sciences, iss. 1491, ISSN 0706-6457, 193 p.

Brand, M.D., Steverding, D., Kadenbach, B., Stevenson, P.M., and Hafner, R.P., 1992, The mechanism of the increase in mitochondrial proton permeability induced by thyroid hormones: European Journal of Biochemistry, v. 206, no. 3, p. 775–781.

Braund, S.R., and Burnham, D.C., 1984, Subsistence economics and marine resource use patterns, *in* Truett, J.C., ed., The Barrow Arch environment and possible consequences of planned offshore oil and gas development: Outer Continental Shelf Environmental Assessment Program, National Oceanic and Atmospheric Administration/Ocean Assessment Division, p. 167–215.

Braund, S.R., and Kruse, J., 2009, Synthesis—Three decades of research on socioeconomic effects related to offshore petroleum development in coastal Alaska: U.S. Department of the Interior, Minerals Management Service, MMS Study Number 2009-006, 405 p., plus 3 appendixes.

Briggs, J.C., 1974, Marine zoogeography: McGraw-Hill, New York, 475 p.

Briggs, J.C., 1995, Global biogeography: Elsevier, Amsterdam, 452 p.

Briggs, J.C., 2003, Marine centres of origin as evolutionary engines: Journal of Biogeography, v. 30, p. 1–18. doi: 10.1046/j.1365-2699.2003.00810.x

Briggs, J.C., 2004, Using marine invertebrate fossils for biogeography: Journal of Biogeography, v. 31, p. 1,711. doi: 10.1111/j.1365-2699.2004.01157.x Briggs, J.C., and Bowan, B.W., 2012, A realignment of marine geographic provinces with particular references to fish distributions: Journal of Biogeography, v. 39, no. 1, p. 12–30.

Brix, O., Thorkildsen, S., and Colosimo, A., 2004,
Temperature acclimation modulates the oxygen binding properties of the Atlantic cod (*Gadus morhua* L.) genotypes-HbI*1/1, HbI*1/2, and HbI*2/2- by changing the concentrations of their major hemoglobin components (results from growth studies at different temperatures): Comparative Biochemistry and Physiology Part A—Molecular and Integrative Physiology, v. 138, no. 2, p. 241–251.

Brodeur, R.D., 1998, In situ observations of the association between juvenile fishes and scyphomedusae in the Bering Sea: Marine Ecology Progress Series, v. 163, p. 11–20.

Brodeur, R.D., and Livingston, P.A., 1988, Food habits and diet overlap of various Eastern Bering Sea fishes: National Oceanic and Atmospheric Administration Technical Memorandum NMFS F/NWC-127, 76 p.

Brodeur, R.D., and Rugen, W.C., 1994, Diel vertical distribution of ichthyoplankton in the northern Gulf of Alaska: Fishery Bulletin, v. 92, no. 2, p. 223–235.

Brown, E.D., 2002, Life history, distribution, and size structure of Pacific capelin in Prince William Sound and the northern Gulf of Alaska: ICES Journal of Marine Science, Journal du Conseil, v. 59, no. 5, p. 983–996.

Brown, J., and Green, J.M., 1976, Territoriality, habitat selection, and prior residency in underyearling *Stichaeus punctatus* (Pisces: Stichaeidae): Canadian Journal of Zoology, v. 54, no. 11, p. 1,904–1,907.

Brown, R.J., 2004, A biological assessment of whitefish species harvested during the spring and fall in the Selawik River Delta Selawik National Wildlife Refuge, Alaska: Alaska Fisheries Technical Report no. 77.

Brown, R.J., Brown, C., Braem, N.M., Carter, W.K., III, Legere, N., and Slayton, L., 2012, Whitefish biology, distribution, and fisheries in the Yukon and Kuskokwim river drainages in Alaska—A synthesis of available information: U.S. Fish and Wildlife Service, Alaska Fisheries Data Series Number 2012-4, 316 p.

Browne, P.J., Laake, L., and DeLong, R.L., 2002, Improving pinnipeds diet analyses through identification of multiple skeletal structures in fecal samples: Fishery Bulletin, v. 100, no. 3, p. 423–433.

Bulatov, O.A., 1983, Distribution of eggs and larvae of Greenland halibut, *Reinhardtius hippoglossoides* (Pleuronectidae), in the eastern Bering Sea: Journal of Ichthyology, v. 23 no. 1, p. 157–159. Bureau of Ocean Energy Management, 2012, Proposed Final Outer Continental Shelf Oil and Gas Leasing Program 2012-2017, U.S. Department of the Interior, 211 p., website accessed November 13, 2013, at http://www.boem.gov/ uploadedFiles/BOEM/Oil_and_Gas_Energy_Program/ Leasing/Five_Year_Program/2012-2017/Program_Area_ Maps/Alaska%20Program%20Areas.pdf.

Bureau of Ocean Energy Management Regulation and Enforcement, 2011, Chukchi Sea Planning Area: U.S. Department of the Interior, Bureau of Ocean Energy Management Regulation and Enforcement, Alaska OCS Region, OCS EIS/EA BOEMHRE 2011-041, Final Supplemental Environmental Impact Statement, v. 1, chaps. I–IV, 590 p.

Burgner, R.L., 1991, Life history of sockeye salmon (*Oncorhynchus nerka*), *in* Groot, C., and Margolis, L., eds., Pacific salmon life histories: University of British Columbia Press, p. 3–117.

Burns, J.J., 1990, Proceedings of a technical workshop on fishes utilized in subsistence fisheries in National Petroleum Reserve-Alaska—Barrow, Alaska, October 26–28, 1988:
Prepared for North Slope Borough, Department of Wildlife Management, Barrow, Alaska, 94 p.

Burton, D.W., 1988, Melanophore comparisons in different forms of ambicoloration in the flatfish *Pseudopleuronectes americanus* and *Reinhardtius hippoglossoides*: Journal of Zoology, v. 214, no. 2, p. 353–360.

Busby, M.S., Matarese, A.C., and Mier, K.L., 2000, Annual, seasonal, and diel composition of larval and juvenile fishes collected by dip-net in Clam Bay, Puget Sound, Washington, from 1985 to 1995: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, NOAA Technical Memorandum NMFS-AFSC-111, 48 p.

Busby, M.S., Mier, K.L., and Brodeur, R.D., 2005, Habitat associations of demersal fishes and crabs in the Pribilof Islands region of the Bering Sea: Fisheries Research, v. 75, no. 1–3, p. 15–28.

Byers, T., and Prach, R.W., 1988, Diet of the kelp snailfish, *Liparis tunicatus*, in Jones Sound, Canadian High Arctic: The Canadian Field-Naturalist, v. 102, no. 2, p. 242–245.

Byrkjedal, I., Brattegard, T., and Møller, P.R., 2009, *Lycodes adolfi* Nielsen and Fosså, 1993 (Teleostei: Zoarcidae) recorded near Jan Mayen and in the eastern part of the Norwegian Sea: Fauna Norvegica, v. 28, p. 1–3.

Byrkjedal, I., Hadler-Jacobsen, S., Rees, D., and Orlov, A., 2014, Molecular and morphological variation in 'saddled' fathead sculpins (Psychrolutidae—*Cottunculus*) of the northeast Atlantic, with a synonymy of three species: Journal of Fish Biology, v. 85, no. 3, p. 821–837.

Byrkjedal, I., Langhelle, G., Wenneck, T.d.L., and
Wienerroither, R., 2011, *Lycodes adolfi* Nielsen and Fosså, 1993 (Teleostei: Zoarcidae) found in the Arctic Ocean:
Polar Biology, v. 34, no. 3, p. 465–467.

Campbell, H.A., Fraser, K.P.P., Bishop, C.M., Peck, L.S., and Egginton, S., 2008, Hibernation in an Antarctic fish—On ice for winter: PLoS ONE: v. 3, iss. 3, e1743, 9 p., doi:10.1371/ journal.pone.0001743.

Carey, A.G., Jr, 1987, Particle flux beneath fast ice in the shallow southwestern Beaufort Sea, Arctic Ocean: Marine Ecology Progress Series, v. 40, p. 247–257.

Carey, A.G., Jr., and Ruff, R.E., 1977, Ecological studies of the benthos in the western Beaufort Sea with special reference to bivalve molluscs, *in* Dunbar, M.J., ed., Polar oceans: Calgary, Alberta, Arctic Institute North America, p. 505–530.

Carey, M.P., Levin, P.S., Townsend, H., Minello, T.J., Sutton, G.R., Francis, T., Harvey, C.J., Toft, J.E., Arkema, K.K., Burke, J.L., Kim, C-K., Guerry, A., Plummer, M., Spiridonov, G., and Ruckelshaus, M., 2013, Characterizing coastal food webs with qualitative links to bridge the gaps between theory and the practice of ecosystem-based management: ICES Journal of Marine Science, v. 71, no. 3, p. 713–724, doi:10.1093/icesjms/fst012.

Carey, M.P., Sanderson, B.L., Barnis, K.A., and Olden, J.L., 2012, Native invaders—Challenges for science, management, policy, and society: Frontiers in Ecology and the Environment, v. 10, no. 7, p. 373–381.

Carlson, H.R., 1995, Consistent yearly appearance of age-0 Walleye Pollock, *Theragra chalcogramma*, at a coastal site in southeastern Alaska, 1973–1994: Fishery Bulletin, v. 93, no. 2, p. 386–390.

Carmack, E., and Kulikov,Y., 1998, Wind-forced upwelling and internal Kelvin wave generation in Mackenzie Canyon, Beaufort Sea: Journal of Geophysical Research, v. 103, no. 18, p. 447–458.

Carmack, E., and Wassmann, P., 2006, Food webs and physical-biological coupling on pan Arctic shelves— Unifying concepts and comprehensive perspectives: Progress in Oceanography, v. 71, nos. 2–4, p. 446–477.

Carmack, Eddy, Winsor, Peter, and Williams, William, 2015, The contiguous panarctic Riverine Coastal Domain—A unifying concept: Progress in Oceanography, v. 139, p. 13–23, accessed December 5, 2015, at http://dx.doi. org/10.1016/j.pocean.2015.07.014.

Carothers, C., Cotton, S., and Moerlein, K., 2013, Subsistence use and knowledge of salmon in Barrow and Nuiqsut, Alaska: U.S. Department of the Interior, Bureau of Ocean Energy Management, OCS Study BOEM 2013-0015, 52 p. Carscadden, E., and Vilhjálmsson, H., 2002, Capelin—What are they good for? Introduction: ICES Journal of Marine Science, v. 59, p. 863–869.

Chambers, R.C., 1997, Environmental influences on egg and propagule sizes in marine fishes, *in* Chambers, R.C., and Trippel, E.A., eds., Early life history and recruitment in fish populations: London, Chapman and Hall, p. 63–102.

Chang, Y.J., Sun, C.L., Zhang, Y., and Yeh, S.Z., 2010, Incorporation climate changes into population dynamic modeling—An individual-based modeling approach for lobster: Canadian Journal of Fisheries Aquatic Science, v. 68, no. 1, p. 122–136.

Chen, A.-L., and Sakurai, Y., 1993, Age and growth of saffron cod (*Eleginus gracilis*): Scientific Reports of Hokkaido Fisheries Experimental Station, v. 42, p. 251–264.

Cheng, C.-H.C., Cziko, P.A., and Evans, C.W., 2006, Nonhepatic origin of notothenioid antifreeze reveals pancreatic synthesis as common mechanism in polar fish freezing avoidance: Proceedings of the National Academy of Sciences of the USA, v. 103, no. 27, p. 10,491–10,496.

Chereshnev, I., Nazarkin, M.V., Skopets, M.B., Pitruk, D., Shestakov, A.V., Yabe, M., and others, 2001, Annotated list of fish-like vertebrates and fish in Tauisk Bay (northern part of the Sea of Okhotsk), *in* Andreev, A.V., and Bergmann, H.H., eds., Biodiversity and ecological status along the northern coast of the Sea of Okhotsk—A collection of study reports: Dalnauka Vladivostok, Russia, Institute of Biological Problems of the North, p. 64–86.

Chereshnev, I., Shestakov, A.V., and Skopets, M.B., 1999, On the distribution of silver smelts of the genus *Hypomesus* (Osmeridae) in the northern part of the Sea of Okhotsk: Journal of Ichthyology, v. 39, no. 7, p. 498–503.

Chernova, N.V., 1991, Aquarium observations of the snail fishes *Careproctus reinhardti and Liparis liparis*: Journal of Ichthyology, v. 31, no. 9, p. 65–70.

Chernova, N.V., 2008, Systematics and phylogeny of fish of the genus *Liparis* (Liparidae, Scorpaeniformes): Journal of Ichthyology, v. 48, no. 10, p. 831–852.

Chernova, N.V., 2011, Distribution patterns and chronological analysis of fish fauna of the Arctic region: Journal of Ichthyology, v. 51, no. 10, p. 825–924.

Cheung, W.W.L., Lam, V.W.Y., Sarmiento, J.L., Kearney, K., Watson, R., and Pauly, D., 2009, Projecting global marine biodiversity impacts under climate change scenarios: Fish and Fisheries, v. 10, p. 235–251.

Christiansen, J.S., ed., 2003, TUNU-1 Expedition—The fish fauna of the NE Greenland fjord systems—Technical report: Tromsø, Norway, University of Tromsø, Norwegian College of Fishery Science, Institute of Aquatic Resources, 33 p. Christiansen, J.S., Chernitsky, A.G., and Karamushko, O.V., 1995, An Arctic teleost fish with a noticeably high body fluid osmolality: a note on the navaga, *Eleginus navaga* (Pallas 1811), from the White Sea: Polar Biology, v. 15, no. 4, p. 303–306.

Christiansen, J.S., Fevolden, S.E., Karamushko, O.V., and Karamushko, L.J., 1998, Maternal output in polar fish reproduction, *in* di Prisco, G., Pisano, E., and Clarke, eds., Fishes of Antractica—A biological overview: Milan, Italy, Springer, p. 41–52.

Christiansen, J.S., Karamushko, O.V., Fevolden, S-E., and Præbel, K., 2008, Marine fishes of North East Greenland— Diversity and adaptation: International Council for the Exploration of the Sea, ICES2008/B, 19 p.

Christiansen, J.S., Mecklenburg, C.W., and Karamushko, O.V., 2014, Arctic marine fishes and their fisheries in light of global climate change: Global Change Biology, v. 20, p. 352–359.

Christiansen, J.S., Reist, J.D., Brown, R.J., Brykov, V.A., Christensen, G., Christoffersen, K., Cott, P., Crane, P., Dempson, J.B., Docker, M., and others, 2013, Fishes, *in* Meltofte, H., ed. Arctic biodiversity assessment—Status and trends in Arctic biodiversity: Akureyri, Iceland, Conservation of Arctic Flora and Fauna, p. 193–245, plus apps., http://www.arcticbiodiversity.is/the-report/chapters/ fishes.

Christensen, V., and Pauly, D., 1992, ECOPATH II—A software for balancing steady-state ecosystem models and calculating network characteristics: Ecological Modeling, v. 61, nos. 3–4, p. 169–185.

Chu, P.B., Wang, Q., and Bourke, R.H., 1999, A geometric model for the Beaufort/Chukchi Sea thermohaline structure: Journal of Atmospheric and Oceanic Technology, v. 16, p. 613–632.

Chuchukalo, V.I., Efimkin, A.Y., and Lapko, V.V., 1995, Feeding of some planktivorous fishes in the Sea of Okhotsk during the summer: Russian Journal of Marine Biology, v. 21, p. 112–115.

Clement, J.P., Bengtson, J.L., and Kelly, B.P., 2013, Managing for the future in a rapidly changing Arctic—A report to the President: Washington, D.C., Interagency Working Group on Coordination of Domestic Energy Development and Permitting in Alaska (Hayes, D.J., Chair): 59 p.

Close, D.A., Fitzpatrick, M.S., and Li, H.W., 2002, The ecological and cultural importance of a species at risk of extinction—Pacific lamprey: Fisheries, v. 27, no. 7, p. 19–25.

Clow, G.D., DeGange, A.R., Derksen, D.V., and Zimmerman, C.E., 2011, Climate change considerations, chap. 4 *of* Holland-Bartels, Leslie, and Pierce, Brenda, eds., An evaluation of the science needs to inform decisions on Outer Continental Shelf energy development in the Chukchi and Beaufort Seas, Alaska: U.S. Geological Survey Circular 1370, p. 81–109.

Coachman, L.R., Aagaard, K., and Tripp, R.B., 1975, Bering Strait-the regional physical oceanography: Seattle, University of Washington Press, 172 p.

Coad, B.W., and Reist, J.D., 2004, Annotated list of the Arctic marine fishes of Canada: Canadian Manuscript Report of Fisheries and Aquatic Sciences, Fisheries and Oceans Canada, no. 2674, 112 p.

Coad, B.W., Waszczuk, H., and Labignan, I., 1995, Encyclopedia of Canadian fishes: Canadian Museum of Nature and Canadian Sportfishing Productions, Inc., 928 p.

Coggan, R., 1996, Growth—Ration relationships in the Antarctic fish *Notothenia coriiceps* R. maintained under different conditions of temperature and photoperiod: Journal of Experimental Marine Biology and Ecology, v. 210, p. 23–35.

Cohen, D.M., Inada, T., Iwamoto, T., and Scialabba, N., 1990, FAO species catalogue, volume 10—Gadiform fishes of the world (Order Gadiformes)—An annotated and illustrated catalogue of cods, hakes, grenadiers and other gadiform fishes known to date: Rome, Food and Agriculture Organization of the United Nations, FAO Fisheries Synopsis, no. 125, v. 10, 442 p.

Colonell, J.M., and Gallaway, B.J., 1997, Wind-driven transport and dispersion of age-0 Arctic ciscoes along the Alaska Beaufort coast: American Fisheries Society Symposium, no. 19, p. 90–103.

Committee on the Status of Endangered Wildlife in Canada, 2004, COSEWIC assessment and update status report on the Bering cisco *Coregonus laurettae* in Canada: Ottawa, Canada, Committee on the Status of Endangered Wildlife in Canada, 19 p.

Cook, A.M., and Bundy, A., 2012, Use of fishes as sampling tools for understanding biodiversity and ecosystem functioning in the ocean: Marine Ecology Progress Series, v. 454, p. 1–18.

Collette, B.B., and Klein-MacPhee, G., 2002, Bigelow and Schroeder's fishes of the Gulf of Maine (3rd ed.): Washington, D.C., Smithsonian Institution Press, 882 p.

Collette, B.B., and Vecchione, M., 1995, Interactions between fisheries and systematics: Fisheries, v. 20, no. 1, p. 20–25.

Coombs, S., and Montgomery, J., 1994, Function and evolution of superficial neuromasts in an Antarctic notothenioid fish: Brain Behavior and Evolution, v. 44, p. 287–298.

Cooper, D.W., Maslenikov, K.P., and Gunderson, D.R., 2007, Natural mortality rate, annual fecundity, and maturity at length for Greenland halibut (*Reinhardtius hippoglossoides*) from the northeastern Pacific Ocean: Fishery Bulletin, v. 105, no. 2, p. 296–304.

Cossins, A.R., and Prosser, C.L., 1978, Evolutionary adaptation of membranes to temperature: Proceedings of the National Academy of Sciences of the USA, v. 75, no. 4, p. 2,040–2,043.

Courcelles, D., 2011, Re-evaluation of the length-weight relationship of Pacific halibut (*Hippoglossus stenolepis*), *in* IPHC Report of Assessment and Research Activities 2011: International Pacific Halibut Commission (IPHC), p. 459–470 [Also available at http://www.iphc.int/library/ raras/254-rara-2011.html.]

Courtney, M., 2014, Dispersal patterns and summer distribution of adult Dolly Varden from the Wulik River, Alaska, evaluated with satellite telemetry [abs.]: Anchorage, Alaska Marine Science Symposium, January 20–24, 2014, p. 15.

Coyle, K.O., Gillispie, J.A., Smith, R.L., and Barber, W.E. 1997, Food habits of four demersal Chukchi Sea fishes, *in* Reynolds, J.B., ed., Symposium 19, Proceedings of the Fish Ecology in Arctic North America Symposium: American Fisheries Society, Fairbanks, Alaska, May 19–21, 1992, p. 310–318.

Craig, P.C., 1984, Fish use of coastal waters of the Alaskan Beaufort Sea—A review: Transactions of the American Fisheries Society, v. 113, no. 3, p. 265–282.

Craig, P.C., 1987, Subsistence fisheries at coastal villages in the Alaskan Arctic, 1970–1986: Department of the Interior, Minerals Management Service, OCS Study MMS 87-6044, Technical Report No. 129, 63 p.

Craig, P.C., 1989a, An introduction to anadromous fishes in the Alaskan Arctic: Biological Papers of the University of Alaska, v. 24, p. 27–54.

Craig, P.C., 1989b, Subsistence fisheries at coastal villages in the Alaskan Arctic, 1970–1986: Biological Papers University of Alaska, no. 24, p. 131–152.

Craig, P.C., Griffiths, W.B., Haldorson, L., and McElderry, H., 1982, Ecological studies of Arctic cod (*Boreogadus saida*) in Beaufort Sea coastal waters, Alaska: Canadian Journal of Fisheries and Aquatic Sciences, v. 39, no. 3, p. 395–406. Craig, P.C., Griffiths, W.B., Haldorson, L., and McElderry, H., 1984a, Distributional patterns in an Alaska Arctic lagoon: Polar Biology, v. 4, no. 1, p. 9–18.

Craig, P.C., Griffiths, W.B., Johnson, S.R., and Schell, D.M., 1984b, Trophic dynamics in an Arctic lagoon, *in* Barnes, P.W., and Reimnitz, E., eds., The Alaskan Beaufort Sea—Ecosystems and environments: New York, Academic Press, p. 347–380.

Craig, P.C., and Haldorson, L.J., 1981, Beaufort Sea Barrier Island Lagoon ecological process studies—Final report, Simpson Lagoon—Fish: U.S. Department of Commerce and U.S. Department of the Interior, Biological Studies, p. 384–649.

Craig, P.C., and Haldorson, L.J., 1986, Pacific salmon in the North American Arctic: Arctic, v. 39, no. 1, p. 2–7.

Craig, P.C., and McCart, P.J., 1974, Fall spawning and overwintering areas of fish populations along routes of proposed pipeline between Prudhoe Bay and the Mackenzie Delta 1972–1973, chap. 3 *of* McCart, P.J., ed., Life histories of anadromous and freshwater fish in the Western Arctic: Canadian Arctic Gas Study Ltd./Alaskan Arctic Gas Study Company, Biological Report Series, v. 20, p. 229.

Craig, P.C., and McCart, P.J., 1975, Fish utilization of nearshore coastal waters between the Colville and Mackenzie Rivers with an emphasis on anadromous species, *in* Craig, P.C., ed., Fisheries investigations in a coastal region of the Beaufort Sea: Calgary, Alberta, Canadian Arctic Gas Study Ltd. and Alaskan Arctic Gas Study Co., Biological Report Series 34: p. 172–219.

Craig, P.C., and Schmidt, D.R., 1985, Fish resources at Point Lay, Alaska: Barrow, Alaska, LGL Alaska Research Associates, Inc., North Slope Borough, Materials Source Division: 105 p.

Crane, P., Viavant, T., and Wenburg, J., 2005, Overwintering patterns of Dolly Varden *Salvelinus malma* in the Sagavanirktok River in the Alaska North Slope inferred using mixed-stock analysis: U.S. Fish and Wildlife Service, Alaska Fisheries Technical Report, no. 84, 25 p.

Crawford, R.E., and Jorgenson, J.K., 1996, Quantitative studies of Arctic cod (*Boreogadus saida*) schools— Important energy stores in the Arctic food web: Arctic, v. 49, no. 2, p. 181–193.

Crawford, R.E., Vagle, S., and Carmack, E.C., 2012, Water mass and bathymetric characteristics of polar cod habitat along the continental shelf and slope of the Beaufort and Chukchi Seas: Polar Biology, v. 35, no. 2, p. 179–190. Criscione, C.D., Watral, V., Whipps, C.M., Blouin, M.S., Jones, S.R.M., and Kent, M.L., 2002, Ribosomal DNA sequences indicate isolated populations of *Ichthyophonus hoferi* in geographic sympatry in the north-eastern Pacific Ocean: Journal of Fish Disease, v. 25, no. 10, p. 575–582.

Cross, J.N., Fresh, K., Miller, B.S., Simenstad, C.A., Steinfort, S.N., and Fegley, J.C., 1978, Nearshore fish and invertebrates assemblages along the Strait of Juan de Fuca including food habits of the common inshore fishes: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Environmental Research Laboratories, NOAA Technical Memorandum ERL MESA-32.

Cui, X., Grebmeier, J.M., Cooper, L.W., Lovvorn, J.R., North, C.A., and Kolts, J.M., 2009, Spatial distributions of groundfish in the northern Bering Sea in relation to environmental variation: Marine Ecology Progress Series, v. 19, no. 6, p. 1,596–1,613.

Cziko, P.A., Evans, C.W., Cheng, C.-H.C., and DeVries, A.L., 2006, Freezing resistance of antifreeze-deficient larval Antarctic fish: The Journal of Experimental Biology, v. 209, p. 407–420.

Datsky, A.V., 2015, Fish fauna of the Chukchi Sea and perspectives of its commercial use: Journal of Ichthyology, v. 55, p. 185–209.

Datsky A.V., and Andronov, P. Yu., 2007, Upper shelf ichtyocenose in the northwestern part of the Bering Sea, Magadan: NASC FEB RAS, 261 p. [In Russian.]

Davenport, J., 1992, Animal life at low temperature: London, Chapman and Hall, 246 p.

Davies, P.L., Baardsnes, J., Kuiper, M.J., and Walker, V.K., 2002, Structure and function of antifreeze proteins:
Physiological Transactions of the Royal Society London B, Biological Science, v. 357, no. 1423, p. 927–935.

Davies, P.L., Hew, C.L., and Fletcher, G.L., 1988, Fish antifreeze proteins—Physiology and evolutionary biology: Canadian Journal of Zoology, v. 66, no. 12, p. 2,611–2,617.

Davison, W., 2005, Antarctic fish skeletal muscle and locomotion, *in* Farrell, A.P., and Steffensen, J.F., eds., The physiology of polar fishes: Amsterdam, Elsevier, p. 317–349.

Davison, W., Axelsson, M., Nilsson, S., and Forster, M.E., 1997, Cardiovascular function in Antarctic notothenioid fishes: Comparative and Biochemical Physiology, v. 118A, p. 1,001–1,008.

DeCicco, A.L., 1992, Long-distance movements of anadromous Dolly Varden between Alaska and the U.S.S.R.: Arctic, v. 45, no. 2, p. 120–123. DeCicco, A.L., 1996, Abundance of Dolly Varden overwintering in the Wulik River, Northwest Alaska, during 1994/1995: Alaska Department of Fish and Game, Fishery Data Series, No. 96-3, 24 p.

DeCicco, A.L., 1997, Movements of postsmolt anadromous Dolly Varden in northwestern Alaska, *in* Reynolds, J.B., ed., Fish ecology in Arctic North America: American Fisheries Society Symposium no. 19, Fairbanks, Alaska, May 19–21, 1992 [Proceedings], p. 175–183.

DeGange, A.R., and Thorsteinson, L.K., 2011, Ecological and subsistence context, chap. 3 *of* Holland-Bartels, L., and Pierce, B., eds., An evaluation of the science needs to inform decisions on Outer Continental Shelf energy development in the Chukchi and Beaufort Seas, Alaska: U.S. Geological Survey Circular 1370, p. 41–80.

DeGraff, D., 1974, The life history of the pond smelt *Hypomesus olidus* Pallas (Osmeridae) in a small unnamed lake in the northern Yukon Territory: Arctic Gas Biological Report Series, v. 18, no. 2, p. 89.

DeGraaf, D.A., 1986, Aspects of the life history of the pond smelt (*Hypomesus olidus*) in the Yukon and Northwest Territories: Arctic, v. 39, no. 3, p. 260–263.

DeGraaf, D.A., and Machniak, K., 1977, Fisheries investigations along the cross delta pipeline route in the MacKenzie Delta, *in* McCart, P.J., ed., Studies to determine the impact of gas pipeline development on aquatic ecosystems: Calgary, Alberta, Canadian Arctic Gas Study, Ltd. and Alaskan Arctic Gas Study Co., Biological Report Series 39, p. 1–169.

Deibel, D., and Daly, L.L., 2007, Zooplankton processes in Arctic and Antarctic polynyas, *in* Smith, W.O., Jr., and Barber, D., eds., Polynyas—Windows into Polar oceans: Elsevier Oceanography Series 74, p. 271–322.

DeMartini, E.E., 1970, Reproductive colorations, paternal behavior and egg masses of kelp greenling, Hexagrammos decagrammus, and whitespotted greenling, H. stelleri: Northwest Science, v. 60, no. 1, p. 32–35.

Denstad, J.-P., Aunaas, T., Börseth, J.B., Vollan Aarset, A., and Zachariassen, K.E., 1987, Thermal hysteresis antifreeze agents in fishes from Spitsbergen waters: Polar Research, v. 5, no. 2, p. 171–174.

Deriso, R.B., 1980, Harvesting strategies and parameter estimation for an age-structured model: Canadian Journal of Fisheries and Aquatic Sciences, v. 37, no. 2, p. 268–282.

Desjardins, M., Le François, N.R., Fletcher, G.L., and Blier, P.U., 2006, Seasonal modulation of plasma antifreeze protein levels in Atlantic (*Anarhichas lupus*) and spotted wolfish (*A. minor*): Journal of Experimental Marine Biology and Ecology, v. 335, no. 1, p. 142–150.

Desjardins, M., Le François, N.R., Fletcher, G.L., and Blier, P.U., 2007, High antifreeze protein levels in wolfish (*Anarchichas lupus*) make them an ideal candidate for culture in cold, potentially ice laden waters: Aquaculture, v. 272, no. 1–4, p. 667–674.

DeVries, A.L., 1980, Biological antifreezes and survival in freezing environments, *in* Gilles, B., ed., Animals and environmental fitness, vol. 1: Oxford, England, Pergamon Press, p. 583–607,

DeVries, A.L., and Cheng, C.-H. C., 2005, Antifreeze proteins and organismal freezing avoidance in polar fishes, *in* Farrell, A.P., and Steffensen, J.F., eds., The physiology of Polar fishes: Amsterdam, Elsevier, p. 155–201.

DeVries, A.L., and Lin, Y., 1977, The role of glycoprotein antifreezes in the survival of Antarctic fishes, *in* Llano, G.A., ed., Adaptations within Antarctic ecosystems: Houston, Texas, Gulf Publishing, p. 439–458.

DeVries, A.L., and Steffensen, J.F., 2005, The Arctic and Antarctic polar marine environments, *in* Farrell, A.P., and Steffensen, J.F., eds., The physiology of polar fishes: Amsterdam, Elsevier, p. 1–24.

Dick, M.H., and Warner, I.M., 1982, Pacific sand lance, *Ammodytes hexapterus Pallas*, in the Kodiak island group, Alaska: Syesis, v. 15, p. 43–50.

Dobbs, G.H., and DeVries, A.L., 1975, Renal function in Antarctic teleosts fish—Serum and urine composition: Marine Biology, v. 29, p. 59–70.

Dodson, J.J., Tremblay, S., Colombani, F., Carscadden, J.E., and Lecomte, F., 2007, Trans-Arctic dispersals and the evolution of a circumpolar marine fish species complex, the capelin (*Mallotus villosus*): Molecular Ecology, v. 16, p. 5,030–5,043.

Doney, S.C., Fabry, V.J., Feely, R.A., and Klepas, J.A., 2009, Ocean acidification—The other CO₂ problem: Annual Review of Marine Science, v. 1, p. 169–192.

Doney, S.C., and others, 2012, Climate change impacts on marine ecosystems: Annual Review of Marine Science, v. 4, p. 11–37.

Dorn, M., Aydin, K., Barbeaux, S., Guttormsen, M., Megrey, B., Spalinger, K., and others, 2007, Gulf of Alaska walleye pollock, *in* The Plan Team for the Groundfish Fisheries of the Gulf of Alaska, compilers, Stock assessment and fishery evaluation report for the groundfish resources of the Gulf of Alaska: Anchorage, Alaska, North Pacific Fishery Management Council, p. 51–168.

Doyle, M.J., Mier, K.L., Busby, M.S., and Brodeur, R.D., 2002, Regional variation in springtime ichthyoplankton assemblages in the northeast Pacific Ocean: Progress in Oceanography, v. 53, no. 2, p. 247–281.

Driedzic, W.R., Clow, K.A., Short, C.E., and Ewart, K.V., 2006, Glycerol production in rainbow smelt (*Osmerus mordax*) may be triggered by low temperature alone and is associated with the activation of glycerol-3-phosphate dehydrogenase and glycerol-3-phosphatase: Journal of Experimental Biology, v. 209, no. 6, p. 1,016–1,023.

Driedzic, W.R., and Short, C.E., 2007, Relationship between food availability, glycerol and glycogen levels in lowtemperature challenged rainbow smelt *Osmerus mordax*: Journal of Experimental Biology, v. 210, no. 16, p. 2,866–2,872.

Dudnik, Y.I., and Dolganov, V.N., 1992, Distribution and abundance of fish on the continental slopes of the Sea of Okhotsk and of the Kuril Islands during the summer of 1989: Journal of Ichthyology, v. 32, no. 9, p. 58–76.

Dudnik, Y.I., and Usol'tsev, E.A., 1964, The herrings of the eastern part of the Bering Sea, *in* Moiseev, P.A., ed., Soviet fisheries investigations in the Northeastern Pacific, part 1: Jerusalem, Israel Program for Scientific Translations [1968], p. 236–240.

Duffy-Anderson, J.T., Doyle, M.J., Mier, K.L., Stabeno, P.J., and Wilderbuer, T.K., 2010, Early life ecology of Alaska plaice (*Pleuronectes quadrituberculatus*) in the eastern Bering Sea— Seasonality, distribution, and dispersal: Journal of Sea Research, v. 64, no. 1–2, p. 3–14.

Dunbar, M.J., and Hildebrand, H.H., 1952, Contributions to the study of the fishes of Ungava Bay: Journal of the Fisheries Research Board of Canada, v. 9, no. 2, p. 83–126.

Duncker, B.P., Koops, M.D., Walker, V.K., and Davies, P.L., 1995, Low temperature persistence of type I antifreeze protein is mediated by cold-specific mRNA stability: FEBS Letters v. 377, no. 2, p. 185–188.

Dunn, J.R., and Matarese, A.C., 1987, A review of the early life history of northeast Pacific gadoid fishes: Fisheries Research, v. 5, nos. 2–3, p. 163–184.

Dunn, J.R., and Sample, T.M., 1976, Greenland turbot (family Pleuronectidae) *in* Pereya, W.T., Reeves, J.E., and Bakkala, R.G., eds., Demersal fish and shellfish resources of the eastern Bering Sea in the baseline year 1975: Seattle, Washington, National Marine Fisheries Service, Northwest and Alaska Fisheries Science Center, Processed Report, p. 475–487.

Dunn, J.R., and Vinter, B.M., 1984, Development of larvae of the saffron cod, *Eleginus gracilis*, with comments on the identification of gadid larvae in Pacific and Arctic waters contiguous to Canada and Alaska: Canadian Journal of Fisheries and Aquatic Sciences, v. 41, no. 2, p. 304–318. Dunton, K.H., Goodall, J.L., Schonberg, S.V., Grebmeier, J.M., and Maidment, D.R., 2005. Multi-decadal synthesis of benthic–pelagic coupling in the western Arctic: role of cross-shelf advective processes, Deep-Sea Research II, v. 52, p. 3,462–3,477.

Dunton, K.H., Reimnitz, E., and Schonberg, S., 1982, An Arctic kelp community in the Alaskan Beaufort Sea: Arctic, v. 35, no. 4, p. 465–484.

Dunton, K.H., Schonberg, S.V., and Cooper, L.W., 2012, Food web structure of the Alaskan nearshore shelf and estuarine lagoons of the Beaufort Sea: Estuaries and Coasts, v. 35, no. 2, p. 416–435.

Dunton, K.H., Weingartner, T., and Carmack, E.C., 2006, The nearshore western Beaufort Sea ecosystem: Progress in Oceanography, v. 71, no. 2–4, p. 362–378.

D'yakov, Y.P., 1982, The fecundity of the Greenland halibut, *Reinhardtius hippoglossoides* (Pleuronectidae), from the Bering Sea: Journal of Ichthyology, v. 22, no. 5, p. 59–64.

Dykstra, C.L., Olsen, W., Williams, G.H., and Gregg, J.L., 2012, *Ichthyophonus* prevalence in Pacific halibut, *in*IPHC Report of Assessment and Research Activities 2012: International Pacific Halibut Commission (IPHC), p. 437–446, [Also available at http://www.iphc.int/publications/rara/2012/rara2012437 ichthyophonus.pdf.]

Eastman, J.T., 1997, Comparison of the Antarctic and Arctic fish faunas: Cybium, v. 21, no. 4, p. 335–352.

Ege, R., and Krogh, A., 1914, On the relation between the temperature and the respiratory exchange in fishes: Internationale Revue der gesamten Hydrobiologie und Hydrographie, v. 7, no. 1, p. 48–55.

Egginton, S., 1996, Blood rheology of Antarctic fishes— Viscosity adaptations at very low temperatures: Journal of Fish Biology, v. 48, p. 513–521.

Ehrlich, P., Dobkin, D.S., and Wheye, D., 1988, The birder's handbook: New York, Simon and Schuster, Inc., 816 p.

Elith, J., and Leathwick, J.R., 2009, Species distribution models—Ecological explanation and prediction across space and time: Annual Review of Ecology, Evolution, and Systematics, v. 40, p. 677–697.

Ellis, D.V., 1962, Observations on the distribution and ecology of some Arctic fish: Arctic, v. 15, no. 3, p. 179–189.

Emmett, R.L., Stone, S.L., Hinton, S.A., and Monaco, M.E., 1991, Distribution and abundance of fishes and inverteprates in west coast estuaries, Volume II—Species life history summaries: National Oceanic and Atmospheric Administration/NOS Strategic Environmental Assessments Division, ELMR Report no. 8, 327 p. Enevoldsen, L.T., Heiner, I., DeVries, A.L., and Steffensen, J.F., 2003, Does fish from the Disko Bay area of Greenland possess antifreeze proteins during the summer?: Polar Biology, v. 26, no. 6, p. 365–370.

Ennis, G.P., 1968, Occurrences of the staghorn sculpin (*Gymnocanthus tricuspis*) in Newfoundland waters: Journal of the Fisheries Research Board of Canada, v. 25, no. 12, p. 2,729–2,731.

Ennis, G.P., 1969, The biology of the shorthorn sculpin *Myoxocephalus scorpius* (L.) in Newfoundland waters: Saint John's, Newfoundland, Memorial University, Master's thesis.

Ennis, G.P., 1970a, Age, growth, and sexual maturity of the shorthorn sculpin, *Myoxocephalus scorpius*, in Newfoundland waters: Journal of the Fisheries Research Board of Canada v. 27, no. 12, p. 2,155–2,158.

Ennis, G.P., 1970b, Reproduction and associated behaviour in the shorthorn sculpin, *Myoxocephalus scorpius* in Newfoundland waters: Journal of the Fisheries Research Board of Canada, v. 27, no. 11, p. 2,037–2,045.

Eschmeyer, W.N., Frick, R., Fong, J.D., and Polack, D.A., 2010, Marine fish diversity—History of knowledge and discovery (Pisces): Zootaxa 2525, p. 19–50.

Eschmeyer, W.N., Frick, R., and van der Laan, R., eds, 2015, Catalog of fishes—Genera, species, references: California Academy of Sciences, Web site accessed June 17, 2016, at http://researcharchive.calacademy.org/research/ichthyology/ catalog/fishcatmain.asp.

Essington, T.E., Kitchell, J.F., and Walters, C.J., 2001, The von Bertalanffy growth function, bioenergetics, and consumption rates of fish: Canadian Journal of Fisheries and Aquatic Sciences, v. 58, no. 11, p. 2,129–2,138.

Evans, C.L., Reist, J.D., and Minns, C.K., 2002, Life history characteristics of freshwater fishes occurring in the Northwest Territories and Nunavut, with major emphasis on riverine habitat requirements: Canadian Manuscript Report of Fisheries and Aquatic Sciences, no. 2614, 169 p.

Evans, C.W., Gubala, V., Nooney, R., Williams, D.E., Brimble, M.A., and DeVries, A.L., 2011, How do Antarctic notothenioid fishes cope with internal ice? A novel function for antifreeze glycoproteins: Antarctic Science, v. 23, no. 1, p. 57–64.

Fabry, V.J., McClintock, J.B., Mathis, J.T., and Grebmeier, J.M., 2009, Ocean acidification at high latitudes—The bellweather: Oceanography, v. 22, no. 4, p. 160.

Fadeev, N.S., 1965, Comparative outline of the biology of flatfishes in the southeastern part of the Bering Sea and condition of their resources, *in* Moiseev, P.A., ed., Soviet fisheries investigations in the northeastern Pacific, Part IV: Jerusalem, Israel Program for Scientific Translations, p. 112–146.

Fadeev, N.S., 1970, The fishery and biological characteristics of yellowfin soles in the eastern part of the Bering Sea *in* Moiseev, P.A., ed., Soviet fisheries investigations in the northeastern Pacific, Part V: Jerusalem, Israel Program for Scientific Translations, p. 332–396.

Falk-Petersen, I.-B., Frivoll, V., Gulliksen, B., and Haug, T., 1986, Occurrence and size/age relations of polar cod, *Boreogadus saida* (Lepechin), in Spitsbergen coastal waters: Sarsia, v. 71, no. 3–4, p. 235–245.

Falk-Petersen, I.-B., Frivoll, V., Gulliksen, B., Haug, T., and Vader, W., 1988, Age/size relations and food of two snailfishes, *Liparis gibbus* and *Careproctus reinhardtii* (Teleostei, Liparididae) from Spitsbergen coastal waters: Polar Biology, v. 8, no. 5, p. 353–358.

Farwell, M.K., Green, J.M., and Pepper, V.A., 1976, Distribution and known life history of *Stichaeus punctatus* in the northwest Atlantic: Copeia, v. 1976, no. 3, p. 598–602.

Fechhelm, R.G., Baker, J.S., Griffiths, W.B., and Schmidt, D.R., 1989, Localized movement patterns of least cisco (*Coregonus sardinella*) and Arctic cisco (*C. autumnalis*) in the vicinity of a solid-fill causeway: Biological Papers of the University of Alaska, v. 24, p. 75–106.

Fechhelm, R.G., Craig, P.C., Baker, J.S., and Gallaway, B.J., 1984, Fish distribution and use of nearshore waters in the northeastern Chukchi Sea: Final Report, LGL Ecological Research Associates Inc., National Oceanic and Atmospheric Administration, OMPA/OCSEAP, 190 p.

Fechhelm, R.G., and Fissel, D.B., 1988, Wind-aided recruitment of Canadian Arctic cisco (*Coregonus autumnalis*) into Alaskan waters: Canadian Journal of Fisheries and Aquatic Sciences, v. 45, no. 5, p. 906–910.

Fechhelm, R.G., and Griffiths, W.B., 1990, The effect of wind on the recruitment of Canadian Arctic cisco (*Coregonus autumnalis*) into the central Alaskan Beaufort Sea: Canadian Journal of Fisheries and Aquatic Sciences, v. 47, p. 2,164–2,171.

Fechhelm, R.G., Griffiths, W.B., Bryan, J.D., Gallaway,
B.J., and Wilson, W.J., 1995a, Application of an *in situ* growth model—Inferred instance of interspecific trophic competition between anadromous fishes of Prudhoe Bay,
Alaska: Transactions of the American Fisheries Society, v. 124, no. 1, p. 55–69.

Fechhelm, R.G., Griffiths, W.B., Wilson, W.J., Gallaway, B.J., and Bryan, J.D., 1995b, Intra- and interseasonal changes in the relative condition and proximate body composition of Broad Whitefish from the Prudhoe Bay region of Alaska: Transactions of the American Fisheries Society, v. 124, no. 4, p. 508–519.

Fechhelm, R.G., Griffiths, W.B., Wilson, W.J., Trimm, B.A., and Colonell, J.M., 1996, The 1995 fish and oceanography study in Mikkelsen Bay, Alaska: Anchorage, Alaska, Prepared by LGL Alaska Research Associates and Woodward-Clyde Consultant for BP Exploration (Alaska) Inc., 102 p. plus apps.

Fechhelm, R.G., Streever, B., and Gallaway, B.J., 2007, The Arctic cisco (*Coregonus autumnalis*) subsistence and commercial fisheries, Colville River, Alaska—A conceptual model: Arctic, v. 60, no. 4, p. 421–429.

Fechhelm, R.G., Wade, G.D., and Link, M.R., 2006, Nearshore Beaufort Sea fish monitoring in the Prudhoe Bay region, 2005: Anchorage, Alaska, Report to British Petroleum (Alaska) Inc. by LGL Alaska Research Associates, Inc., 72 p. plus apps.

Fedorov, V.V., 1986, Cottidae, *in* Whitehead, P.J.P., Bauchot, M.-L., Hureau, J.-C., Nielsen, J., and Tortonese, E., eds., Fishes of the North-eastern Atlantic and the Mediterranean: UNESCO, Paris, v. 3, p. 1,243–1,260.

Fedorov, V.V., 2000, Species composition, distribution and habitation depths of the northern Kuril Islands fish and fish like species, *in* Koteneva, B.N., ed., Commercial and biological studies of fishes in the Pacific waters of the Kuril Islands and adjacent areas of the Okhotsk and Bering Seas in 1992–1998: Moscow, VNIRO Publishing, collected papers, p. 7–41. [In Russian.]

Field, L.J., 1988, Pacific sand lance, *Ammodytes hexapterus*, with notes on related *Ammodytes species*, *in* Wilimovsky, N.J., Incze, L.S., and Westrheim, S.J., eds., Species synopses, life histories of selected fish and shellfish of the northeast Pacific and Bering Sea: Seattle, University of Washington, Washington Sea Grant Program and Fisheries Research Institute.

Finley, K.J., and Gibb, E.J., 1982, Summer diet of the narwhal (*Monodon monoceros*) in Pond Inlet, northern Baffin Island: Canadian Journal of Zoology, v. 60, no. 12, p. 3,353–3,363.

Fletcher, G.L., 1975, The effects of capture, "stress," and storage of whole blood on the red blood cells, plasma proteins, glucose, and electrolytes of the winter flounder (*Pseudopleuronectes americanus*): Canadian Journal of Zoology, v. 53, no. 2, p. 197–206. Fletcher, G.L., 1977, Circannual cycles of blood plasma freezing point and Na+ and CL– concentrations in Newfoundland winter flounder (*Pseudopleuronectes americanus*)—Correlation with water temperature and photoperiod: Canadian Journal of Zoology, v. 55, no. 5, p. 789–795.

Fletcher, G.L., 1981, Effects of temperature and photoperiod on the plasma freezing point depression, Cl⁻ concentration, and protein "antifreeze" in winter flounder: Canadian Journal of Zoology, v. 59, no. 2, p. 193–201.

Fletcher, G.L., Addison, R.F., Slaughter, D., and Hew, C.L., 1982, Antifreeze proteins in the Arctic shorthorn sculpin (*Myoxocephalus scorpius*): Arctic, v. 35, no. 2, p. 302–306.

Fletcher, G.L., Haya, K., King, M.J., and Reisman, H.M., 1985a, Annual antifreeze cycles in Newfoundland, New Brunswick and Long island winter flounder, *Pseudopleuronectes americanus*: Marine Ecology Progress Series, v. 21, p. 205–212.

Fletcher, G.L., Hew, C.L., and Davis, P.L., 2001, Antifreeze proteins of teleost fishes: Annual Review of Physiology, v. 63, p. 359–390.

Fletcher, G.L., Hew, C.L., Li, X.M., Haya, K., and Kao, M.H., 1985b, Year-round presence of high levels of plasma antifreeze peptides in a temperate fish, ocean pout (*Macrozoarces americanus*): Canadian Journal of Zoology, v. 63, no. 3, p. 488–493.

Fletcher, G.L., Idler, D.R., Vaisius, A., and Hew, C.L., 1989, Hormonal regulation of AFP gene expression in winter flounder: Fish Physiology and Biochemistry, v. 7, no. 1–6, p. 387–393.

Fletcher, G.L., Kao, M.H., and Haya, K., 1984, Seasonal and phenotypic variations in plasma protein antifreeze levels in a population of marine fish, sea raven (*Hemitripterus americanus*): Canadian Journal of Fisheries and Aquatic Sciences, v. 41, no. 5, p. 819–824.

Fletcher, G.L., King, M.J., and Kao, M.H., 1987, Low temperature regulation of antifreeze glycopeptides levels in Atlantic cod (*Gadus morhua*): Canadian Journal of Zoology, v. 65, no. 2, p. 227–233.

Fluharty, D., 2012, Recent developments at the Federal level in ocean policymaking in the United States: Coastal Management, v. 40, no. 2, p. 209–221.

Fordham, D.A., and others, 2013, Population dynamics can be more important than physiological limits for determining range shifts under climate change, Global Change Biology, v. 19, p. 3,224–3,237. Forrest, A., Sampei, M., Hatton, H., Makabe, R., Sasaki, H., Fukuchi, M., Wassman, P., and Fortier, L., 2007, Particulate organic carbon fluxes on the slope of the MacKenzie Shelf (Beaufort Sea)—Physical and biological forcing of shelfbasin exchanges: Journal of Marine Systems: v. 68, no. 1–2, p. 39–54.

Forrester, C.R., 1964, Laboratory observations on embryonic development and larvae of the Pacific cod (*Gadus macrocephalus* Tilesius): Journal of the Fisheries Research Board of Canada, v. 21, no. 1, p. 9–16.

Forsberg, J.E., 2011, Age distribution of Pacific halibut in the 2011 IPHC stock assessment setline survey, *in* International Pacific Halibut Commission (IPHC) Report of Assessment and Research Activities 2011: International Pacific Halibut Commission, p. 553–560. [Also available at http://www.iphc.int/library/raras/254-rara-2011.html.]

Fortier, L., Sirois, P., Michaud, J., and Barber, D., 2006, Survival of Arctic cod larvae (*Boreogadus saida*) in relation to sea ice and temperature in the Northeast Water Polynya (*Greenland Sea*): Canadian Journal of Fisheries and Aquatic Sciences, v. 63, no. 7, p. 1,608–1,616.

Fourney, R.M., Fletcher, G.L., and Hew, C.L., 1984, The effects of long day lengths on liver antifreeze mRNA in winter flounder: Canadian Journal of Zoology, v. 62, no. 8, p. 1,456–1,460.

Foy, R.J., Crapo, C.A., and Kramer, D.E., 2006, Investigating the roles of temperature and exercise in the development of chalkiness in Pacific halibut: International Pacific Halibut Commission, Technical Report No. 50.

Fresh, K.L., 1979, Distribution and abundance of fishes occurring in the nearshore surface waters of northern Puget Sound, Washington: Seattle, University of Washington, Master's thesis.

Frey, K.E., Arrigo, K.R., and Gradinger, R.R., 2011, Arctic report card, update for 2011—Arctic Ocean primary productivity: National Oceanic and Atmospheric Administration Arctic Theme Page Web site, accessed October 25, 2012, at http://www.arctic.noaa.gov/report11/ primary_productivity.html.

Froese, R., and Pauly, D., eds., 2012, FishBase—Global information system on fishes: FishBase database, accessed July 8, 2012, at http://www.fishbase.org.

Frost, K.J., and Lowry, L.F., 1980, Feeding of ribbon seals (*Phoca fasciata*) in the Bering Sea in spring: Canadian Journal of Zoology, no. 58, v. 9, p. 1,601–1,607.

Frost, K.J., and Lowry, L.F., 1981, Trophic importance of some marine gadids in Northern Alaska and their bodyotolith size relationships: Fishery Bulletin, v. 79, no. 1, p. 187–192.

Frost, K.J., and Lowry, L.F., 1983, Demersal fishes and invertebrates trawled in the northeastern Chukchi and western Beaufort Seas 1976–1977: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Technical Report NMFS-SSRF-764, 22 p.

Frost, K.J., and Lowry, L.F., 1984, Trophic relationships of vertebrate consumers in the Alaskan Beaufort Sea, *in* Barnes, P.W., Schell, D.M., and Reimnitz, E., eds., The Alaska Beaufort Sea—Ecosystems and environments: San Diego, Academic Press, p. 382–401.

Fruge, D.J., and Wiswar, D.W., 1991, First records of the Bering wolffish, *Anarhichas orientalis*, for the Alaskan Beaufort Sea: Canadian Field-Naturalist, v. 105, no. 1, p. 107–109.

Fruge, D.J., Wiswar, D.W., Dugan, L.J., and Palmer, D.E., 1989, Fish population characteristics of Arctic National Wildlife Refuge coastal waters, summer 1988: Fairbanks, Alaska, U.S. Fish and Wildlife Service, Fishery Assistance office, Progress Report, 73 p.

Frydenberg, O., Møller, D., Névdal, G., and Sick, K., 1965, Hemoglobin polymorphism in Norwegian cod populations: Hereditas, v. 53, no. 1, p. 257–271.

Furniss, R.A., 1975, Inventory and cataloging of Arctic area waters: Alaska Department of Fish and Game, Federal Aid in Fish Restoration, Annual Report of Performance, Project F-9-7, Study G-1-1, v. 16, 47 p.

Gallaway, B.J., and Fechhelm, R.G., 1997, Anadromous and amphidromous fishes, *in* Truett, J.C., and Johnson, S.R., eds., The natural history of an Arctic oil field: San Diego, Academic Press, p. 349–369.

Gallaway, B.J., Griffiths, W.B., Craig, P.C., Gazey, W.J., and Helmericks, J.W., 1983, An assessment of the Colville River Delta stock of Arctic cisco—Migrants from Canada?: Biological Papers of the University of Alaska, no. 21, p. 4–23.

Gallaway, B.J., and Norcross, B.L., eds., 2011, A synthesis of diversity, distribution, abundance, age, size and diet of fishes in the lease sale 193 Area of the northeastern Chukchi Sea: Anchorage, Alaska, Final report prepared for CononcoPhillips Alaska, Inc., Shell Exploration and Production Co., and StatOil USA E&P, Inc., 218 p.

Gamito, S., 1998, Growth models and their use in ecological modeling—An application to a fish population: Ecological Modeling, v. 113, nos. 1–3, p. 83–94.

Garrett, D.L., Pietsch, T.W., Utter, F.M., and Hauser, L., 2007, The hybrid sole, *Inopsetta ischyra* (Teleostei: Pleuronectiformes: Pleuronectidae)—Hybrid or biological species?: Transactions of the American Fisheries Society, v. 136, no. 2, p. 460–468. Garrison, K.J., and Miller, B.S., 1982, Review of the early life history of Puget Sound fishes: Seattle, University of Washington, Fisheries Research Institute, 729 p.

Gaston, A.J., Cairns, D.K., Elliot, R.D., and Noble, D.G., 1985, A natural history of Digges Sound: Report Series Canadian Wildlife Service no. 46, Canadian Government Publishing Centre, Canada Communication Group, 63 p.

George, C., Moulton, L.L., and Johnson, M., 2007, A field guide to the common fishes of the North Slope of Alaska: Alaska Department of Wildlife Management, North Slope Borough, 93 p.

Georgette, S., and Shiedt, A., 2005, Whitefish—Traditional ecological knowledge and subsistence fishing in the Kotzebue Sound, Alaska: Alaska Department of Fish and Game, Technical Report No. 290, 148 p.

Gieseg, S., Cuddihy, S., Hill, J.V., and Davison, W., 2000, A comparison of plasma vitamin C and E levels in two Antarctic and two temperate water fish species: Comparative Biochemistry and Physiology, Part B— Biochemistry and Molecular Biology, v. 125, no. 3, p. 371–378.

Gillispie, J.G., Smith, R.L., Barbour, E., and Barber, W.E., 1997, Distribution, abundance, and growth of Arctic cod in the Northeastern Chukchi Sea, *in* Reynolds, J.B., ed., Symposium 19, Proceedings of the Fish Ecology in Arctic North America Symposium, Fairbanks, Alaska, May 19–21, 1992: American Fisheries Society, p. 81–89.

Gillman, D.V., and Kristofferson, A.H., 1984, Biological data on Pacific herring (*Clupea harengus pallasi*) from Tuktoyaktuk Harbour and the Liverpool Bay area, Northwest Territories, 1981 to 1983: Winnipeg, Manitoba, Canada, Canadian Data Report of Fisheries and Aquatic Sciences, Department of Fisheries and Oceans, Western Region, no. 485, 22 p.

Gjøsaeter, J., 1981, Growth, production and reproduction of the Myctophid fish *Benthosema glaciale* from western Norway and adjacent seas: Fiskeridirektoratets skrifter, Serie Havundersøkelse, v. 17, no. 3, p. 79–108.

Goddard, S.V., and Fletcher, G.L., 2002, Physiological ecology of antifreeze proteins—A northern perspective, *in* Ewart, K.V., and Hew, C.L., eds., Molecular aspects of fish and marine biology, volume 1—Fish antifreeze proteins: Singapore, World Scientific, p. 17–60.

Goddard, S.V., Kao, M.H., and Fletcher, G.L., 1999, Population differences in antifreeze production cycles of juvenile Atlantic cod reflect adaptations to overwintering environment: Canadian Journal of Fisheries and Aquatic Sciences, v. 56, no. 11, p. 1,991–1,999. Goetz, F.W., McCauley, L., Goetz, G.W., and Norberg, B., 2005, Using global genome approaches to address problems in cod mariculture: ICES Journal of Marine Science, v. 63, no. 2, p. 393–399.

Gomelyuk, V.E., 2000, Comparative analysis of everyday behavior and mode of the life of three species of *Hexagrammos* (Hexagrammidae, Scorpaeniformes) in summer: Journal of Ichthyology, v. 40, no. 1, p. 74–85.

Gong, H., Croft, K., Driedzic, W.R., and Ewart, K.V., 2011, Chemical chaperoning action of glycerol on the antifreeze protein of rainbow smelt: Journal of Thermal Biology, v. 36, no. 1, p. 78–83.

Gong, Z., Fletcher, G.L., and Chew, C.L., 1992, Tissue distribution of fish antifreeze protein mRNAs: Canadian Journal of Zoology, v. 70, no. 4, p. 810–814.

Grant, W.S., Bakkala, R.G., Utter, F.M., Teel, D.J., and Kobayashi, T., 1983, Biochemical genetic population structure of yellowfin sole, *Limanda aspera*, of the North Pacific Ocean and Bering Sea: Fishery Bulletin, v. 81, no. 4, p. 667–677.

Grebmeier, J.M., Cooper, L.W., Ashjian, C.A., Bluhm, B.A., Campbell, R.B., Dunton, K.E., Moore, J., Okkonen, S., Sheffield, G., Trefry, J., and Pasternak, S.Y., 2015, Pacific Marine Arctic Regional Synthesis (PacMARS) Final Report: North Pacific Research Board, 259 p.

Grebmeier, J.M., Cooper, L.W., Feder, H.M., and Sirenko, B.I., 2006a, Ecosystem dynamics of the Pacific-influenced North Bering and Chukchi Seas in the Amerasian Arctic: Progress in Oceanography, v. 71, issues 2–4, p. 331–361.

Grebmeier, J.M. and Maslowski, W., eds., 2014, The Pacific Arctic Region: Ecosystem status and trends in a rapidly changing environment, Springer Science+Business Media Dordrecht, 450 p.

Grebmeier, J.M., Overland, J.E., Moore, S.E., Farley, E.V., Carmack, E.C., Cooper, L., Frey, K.E., Helle, J.J., McLaughlin, F.A., and McNutt, S.L., 2006b, A major ecosystem shift in the northern Bering Sea: Science, v. 311, no. 5766, p. 1,461–1,464.

Green, J.M., 1974, A localized mass winter kill of cunners in Newfoundland: Canadian Field-Naturalist, v. 88, p. 96–97.

Green, J.M., and Farwell, M., 1971, Winter habits of the cunner, *Tautogolabrus adspersus* (Walbaum 1792), in Newfoundland: Canadian Journal of Zoology, v. 49, no. 12, p. 1,497–1,499.

Green, J.M., and Mitchell, L.R., 1997, Biology of the fish doctor, an eelpout, from Cornwallis Island, Northwest Territories, Canada: American Fisheries Society Symposium 19, p. 140–174. Green, J.M., and Steele, D.H., 1975, Observations on marine life beneath sea ice, Resolute Bay, N.W.T., *in* Proceedings of the Circumpolar Conference on Northern Ecology, September 15–18, 1975: Ottawa, Ontario, National Research Council of Canada, p. II-77–II-86.

Greenwald, N., Callimanis, S., Garty, A., and Peters, E., 2006, The Bering Sea—A biodiversity assessment of vertebrate species: Portland, Oreg., Center for Biological Diversity, Pacific Environment, 67 p. plus ACCESS database.

Gregg, J.L., Anderi, D.M., and Kimura, D.K., 2006, Improving the precision of otolith-based age estimates for Greenland halibut (*Reinhardtius hippoglossoides*) with preparation methods adapted for fragile sagittae: Fishery Bulletin, v. 104, no. 4, p. 643–648.

Greiner, B., 2009, Estimation of Dolly Varden and documentation of water and weather parameters on the Hulahula River, Alaska 2008: Meadville, Pennsylvania, Allegheny College, Department of Environmental Science, Senior thesis, 36 p.

Griffiths, W., Craig, P., Walder, G., and Mann, G., 1975,
Fisheries investigations in a coastal region of the Beaufort
Sea (Nunaluk Lagoon, Yukon), *in* Craig, P.C., ed., Fisheries
investigations in a coastal region of the Beaufort Sea:
Calgary, Alberta, Canadian Arctic Gas Study, Ltd. and
Alaskan Arctic Gas Study Co., Arctic Gas Biological Report
Series 34 (2), 219 p.

Griffiths, W.B., Fechhelm, R.G., Gallaway, B.J., and Fissel, D.B., 1988, Yukon fish and effects of causeway: Prepared for Indian and Northern Affairs Canada, LGL Ltd. and Arctic Science Ltd., 124 p.

Griffiths, W.B., Fechhelm, R.G., Gallaway, B.J., Martin, L.R., and Watson, W.J., 1998, Abundance of selected fish species in relation to temperature and salinity patterns in the Sagavanirktok Delta, Alaska, following construction of the Endicott Causeway: Arctic, v. 51, no. 2, p. 94–104.

Griffiths, W.B., Gallaway, B.J., Gazey, W.J., and Dillinger, R.E., Jr., 1992, Growth and condition of Arctic Cisco and Broad Whitefish as indicators of causeway-induced effects in the Prudhoe Bay Region, Alaska: Transactions of the American Fisheries Society, v. 121, no. 5, p. 557–577.

Grigorev, S.S., 1995, Characteristics of flatfish eggs and larva samples near eastern Kamchatka and in the western Bering Sea during June-July 1991, *in* Alaska Sea Grant Program, eds., Proceedings of the International Symposium on North Pacific Flatfish: University of Alaska, Fairbanks, Alaska Sea Grant Program, AK-SG-95-04, p. 101–116.

Grigorev, S.S., and Fadeev, N.S., 1995, Distribution of Alaska plaice, flathead sole, and Bering flounder eggs in the eastern Bering Sea during April–July, 1988–1990, *in* Alaska Sea Grant Program, eds., Proceeding of the International Symposium on North Pacific Flatfish: Fairbanks, University of Alaska, Alaska Sea Grant Program, AK-SG-95-04.

Gritsenko, O.F., Churikov, A.A., and Rodionova, S.S., 1984, The reproductive ecology of the Arctic smelt, *Osmerus mordax dentex* (Osmeridae), in the rivers of Sakhalin Island: Journal of Ichthyology, v. 24, no. 3, p. 23–33.

Gross, M.R., 1987, Evolution of diadromy in fines: American Fisheries Society Symposium, no. 1, p. 14–25.

Grossberg, R., and Cunningham, C.W., 2001, Genetic structure in the sea from populations to communities, chap. 3 *of* Bertness M.D., Gaines, S.D., and Hay, M.E., eds., Marine community ecology: Sunderland, Massachusetts, Sinauer Associates, Inc., p. 61–84.

Gunderson, D.R., 1980, Using r-K selection theory to predict natural mortality: Canadian Journal of Fisheries and Aquatic Sciences, v. 37, no. 12, p. 2,266–2,271.

Gustafson, R.G., Lenarz, W.H., McCain, B.B., Schmitt, C.C., Grant, W.S., Builder, T.L., and Methot, R.D., 2000, Status review of Pacific hake, Pacific cod, and walleye pollock from Puget Sound, Washington: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, NOAA Technical Memorandum NMFS-NWFSC-44.

Haertel, L., and Osterberg, C., 1967, Ecology of zooplankton, benthos and fishes in the Columbia River Estuary: Ecology, v. 48, no. 3, p. 459–472.

Haldorson, L.J., and Craig, P., 1984, Life history and ecology of a Pacific-Arctic population of rainbow smelt in coastal waters of the Beaufort Sea: Transactions of the American Fisheries Society, v. 113, no. 1, p. 33–38.

Haldorson, L., Pritchett, M., Sterritt, D., and Watts, J., 1993,
Abundance patterns of marine fish larvae during spring in a southeastern Alaskan bay: Fishery Bulletin, v. 91, no. 1,
p. 36–44.

Hamada, T., 1966, Studies on fluctuations in the abundance of larval sand-lance in the Harima-nada and Osaka Bay
I—Relation between the progeny abundance and the age composition of parent fish: Bulletin of the Japanese Society of Fisheries Oceanography, v. 32, no. 5, p. 393–398.

Hamazaki, T., Fair, L., Watson, L., and Brennan, E., 2005, Analyses of Bering Sea bottom-trawl surveys in Norton Sound—Absence of regime shift effects on epifauna and demersal fish: ICES Journal of Marine Science, v. 62, no. 8, p. 1,597–1,602. Hardie, D.C., 2003, The evolutionary ecology of Arctic fishes: InfoNorth v. 56, no. 4, p. 430–433.

Harris, C.K., and Hartt, A.C., 1977, Assessment of pelagic and nearshore fish in three bays on the east and south coasts of Kodiak Island, Alaska: Seattle, University of Washington, Fisheries Research Institute, FRI-UW-7719.

Harry, G.Y., and Hartley, J.R., 1981, Northern fur seals in the Bering Sea, *in* Hood, D.W., and Calder, J.A., eds., The Eastern Bering Sea shelf: Oceanography and Resources: Seattle, University of Washington Press, p. 847–867.

Hart, J.L., 1973, Pacific fishes of Canada: Ottawa, Fisheries Research Board of Canada, 740 p.

Hart, P.J.B., and Reynolds, J.D., eds., 2002, Handbook of fish biology and fisheries—2 volume set: Blackwell Science, Ltd., 856 p.

Harwood, L.A., Sandstrom, S., and Linn, E., 2009, Status of anadromous Dolly Varden (*Salvelinus malma*) of the Rat River, Northwest Territories, as assessed through sampling of the subsistence fishery (1995–2007): Canadian Manuscript Report of Fisheries and Aquatic Sciences 2891, 52 p.

Harley, C.D.G., Hughs, A.R., Hutgren, K.M., Miner, B.G., Sorte, C.J.B., Thomber, C.S., Rodriguez, L.F., Tomanek, L., and Williams, S.L., 2006, The impacts of climate change in coastal systems: Ecological Letters, v. 9, p. 228–241.

Hatch, S.A., and Sanger, G.A., 1992, Puffins as samplers of juvenile pollock and other forage fish in the Gulf of Alaska: Marine Ecology Progress Series, v. 80, p. 1–14.

Hauser, L., Spies, I., and Loher, T., 2006, Microsatellite screening in Pacific halibut (*Hippoglossus stenolepis*) and a preliminary examination of population structure based on observed DNA variation: Seattle, Washington, International Pacific Halibut Commission Scientific Report, no. 81.

Hayes, D.L., and Kuletz, K.J., 1997, Decline of pigeon guillemot populations in Prince William Sound, Alaska, and apparent changes in distribution and abundance of their prey, *in* Forage fishes in marine ecosystems—Proceedings of the International Symposium on the Role of Forage Fishes in Marine Ecosystems: University of Alaska, Fairbanks, Alaska Sea Grant College Program, p. 699–702.

Healey, M.C., 1991, Life history of Chinook salmon (*Oncorhynchus tshawytscha*), *in* Groot, C., and Margolis, L., eds., Pacific salmon life histories, Vancouver, British Columbia, University of British Columbia Press, p. 313–393.

Heard, W.R., 1991, Life history of pink salmon (*Oncorhynchus gorbuscha*), *in* Groot, C., and Margolis, L., eds., Pacific salmon life histories: Vancouver, British Columbia, UBC Press, p. 121–230. Hebert P.D.N., and Wearing-Wilde J., eds., 2002, Canada's Polar Life: Web page, CyberNatural Software, University of Guelph, accessed October 15, 2013, at http://www.polarlife. ca.

Heikkinen, R.K., Luoto, M., Araújo, M.B., Virkkala, R.V., Thuiller, W., and Sykes, M.T., 2006, Methods and uncertainties in bioclimatic envelope modelling under climate change: Progress in Physical Geography, v. 30, no. 6, p. 1–17.

Hershberger, P.K., Pacheco, C.A., Gregg, J.L., Purcell, M.K., and LaPatra, S.E., 2008, Differential survival of Ichthyophonus isolates indicates parasite adaptation to its host environment: Journal of Parasitology, v. 94, no. 5, p. 1,055–1,059.

Hershberger, P., Rhodes, L., Kurath, K., and Winton, J., 2013, Infectious diseases of fishes in the Salish Sea: Fisheries, v. 38, no. 9, p. 402–409.

Hew, C.L., Fletcher, G.L., and Ananthanarayanan, V.S.,
1980, Antifreeze proteins from the shorthorn sculpin, *Myoxocephalus scorpius*: isolation and characterization:
Canadian Journal of Biochemistry, v. 58, no. 5, p. 377–383.

Hew, C.L., Wang, N.-C., Joshi, S., Fletcher, G.L., Scott, G.K., Hayes, P.H., Buettner, B., and Davies, P.L., 1988, Multiple genes provide the basis for antifreeze protein diversity and dosage in ocean pout, *Macrozoarces americanus*: Journal of Biological Chemistry v. 263, p. 12,049–12,055.

Hildebrandt, N., Bergmann, M., and Knust, R., 2011, Longevity and growth efficiency of two deep-dwelling Arctic zoarcids and comparison with eight other zoarcid species from different climatic regions: Polar Biology, v. 34, no. 10, p. 1,523–1,533.

Hill, J.V., and Egginton, S., 2010, Control of branchial artery tone in fish—Effects of environmental temperature and phylogeny: Physiological Biochemistry and Zoology, v. 83, no. 1, p. 33–42.

Hjorth, J.P., 1974, Molecular and genetic structure of multiple hemoglobins in the eelpout, *Zoarces viviparous* L: Biochemical Genetics, v. 13, no. 5–6, p. 379–391.

Hobson, E.S., 1986, Predation on the Pacific sand lance, *Ammodytes hexapterus* (Pisces: Ammodytidae), during the transition between day and night in southeastern Alaska: Copeia, v. 1986, no. 1, p. 223–226.

Hoff, G.R., 2006, Biodiversity as an index of regime shift in the eastern Bering Sea: Fishery Bulletin, v. 104, no. 2, p. 226–237.

Hoff, G.R., and Britt, L.L., 2003, The 2002 Eastern Bering Sea upper continental slope survey of groundfish and invertebrate resources: U.S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Memorandum, NMFS-AFSC-141, Department of Commerce, 286 p.

Hoff, G.R., Stevenson, D.E., and Orr, J.W., 2015, Guide to the gadiform fishes of the eastern North Pacific: U.S. Department of Commerce, National Oceanic and Atmospheric Aadministration Technical Memorandum NMFS-AFSC-309, 68 p. doi:10.7289/V5/TM-AFSC-309, available at http://www.afsc.noaa.gov/Publications/ AFSC-TM/NOAA-TM-AFSC-309.pdf.

Hoff, G.R., and Britt, L.L., 2005, Results of the 2004 Eastern Bering Sea upper continental slope survey of groundfish and invertebrate resources: National Oceanic and Atmospheric Administration Technical Memorandum, 300 p.

Hognestad, P.T., 1968, Observations on polar cod in the Barents Sea: Rapports et procès-verbaux des reunions: Conseil permanent international pour l'exploration de la mer., v. 158, p. 126–130.

Holeton, G.F., 1974, Metabolic cold adaptation of polar fish— Fact or artefact?: Physiological Zoology, v. 47, no. 3, p. 137–152.

Holladay, B.A., and Norcross, B.L., 1995, Diet diversity as a mechanism for partitioning nursery grounds of pleuronectids: Proceedings of the International Symposium on North Pacific Flatfish, Alaska Sea Grant AK-SG-95-04, p. 177–203.

Holland-Bartels, L., and Pierce, B., eds., 2011, An evaluation of the science needs to inform decisions on Outer Continental Shelf energy development in the Chukchi and Beaufort Seas, Alaska: U.S. Geological Survey Circular 1370, 278 p.

Hollowed, A.B., Planque, B., and Loeng, H., 2013, Potential movement of fish and shellfish stocks from the sub-Arctic to the Arctic Ocean: Fisheries Oceanography, v. 22, no. 5, p. 355–370.

Holmes, W.N., and Donaldson, E.M., 1969, The body compartments and distribution of electrolytes: *in* Hoar, W.S. and Randall, D.J., eds., Fish physiology, volume 1: New York, Academic Press, p. 1–89,

Hop, H., and Graham, M., 1995, Respiration of juvenile Arctic cod (*Boreogadus saida*)—Effects of acclimation, temperature, and food intake: Polar Biology, v. 15, no. 5, p. 359–367.

Hop, H., Welch, H.E., and Crawford, R.E., 1997, Population structure and feeding ecology of Arctic cod schools in the Canadian High Arctic: American Fisheries Society Symposium 19, p. 13–39.

Hopky, G.E., and Ratynski, R.A., 1983, Relative abundance, spatial and temporal distribution, age and growth of fishes in Tuktoyaktuk Harbour, N.W.T., 28 June to 5 September, 1981: Canadian Manuscript Report of Fisheries and Aquatic Sciences 1983, 77 p.

Hopcroft, R., Bluhm, B., and Gradinger, R., eds., 2008, Arctic Ocean synthesis—Analysis of climate change impacts in the Chukchi and Beaufort Seas with strategies for future research: Fairbanks, University of Alaska, 184 p.

Houston, J., and McAllister, D.E., 1990a, Status of the Bering wolfish, *Anarhichas orientalis*, in Canada: The Canadian Field-Naturalist, v. 104, no. 1, p. 20–23.

Houston, J., and McAllister, D.E., 1990b, Status of the blackline prickleback, *Acantholumpenus mackayi*, in Canada: The Canadian Field-Naturalist, v. 104, no. 1, p. 24–28.

Howard, E.I., Blakeley, M.P., Haertlein, M., Petit-Haertlein,
I., Mitschler, A., Fisher, S.J., Cousido-Siah, A., Salvay,
A.G., Popov, A., Muller-Dieckmann, C., Petrova, T., and
Podjarny, A., 2010, Neutron structure of type-III antifreeze
protein allows the reconstruction of AFP-ice interface:
Journal of Molecular Recognition, v. 24, no. 4, p. 724–732.

Howland, K.L., Gendron, M., Tonn, W.M., and Tallman, R.F., 2004, Age determination of a long-lived coregonid from the Canadian North—Comparison of otoliths, fin rays and scales in inconnu (*Stenodus leucichthys*): Annales Zoologici Fennici, Finnish Zoological and Botanical Publishing Board, p. 205–214.

Hufford, G.L., 1974, WEBSEC 71-72—An ecological survey in the Beaufort Sea, August–September 1971–1972: U.S. Coast Guard Oceanographic Unit report No. CG 373-64, 268 p.

Hunt, B.M., Hoefling, K., and Cheng, C.-H. C., 2003, Annual warming episodes in seawater temperatures in McMurdo Sound in relationship to endogenous ice in notothenioid fish: Antarctic Science, v. 15, no. 3, p. 333–338.

Hunt G.L. Jr., and McKinnell, S., 2006, Interplay between top-down, bottom-up, and wasp-waist control in marine ecosystems: Progress in Oceanography, v. 68, no. 2–4, p. 115–124 Hunt, J.L., Jr., Blanchard, A.L., Boveng, Peter, Dalpadado, Padmini, Drinkwater, K.F., Eisner, Lisa, Hopcroft, R.R., Kovacs, K.M., Norcross, B.L., Renaud, Paul, Reigstad, Marit, Renner, Martin, Skjoldal, H.R., Whitehouse, Andy, and Woodgate, R.A, 2013, The Barents and Chukchi Seas— Comparison of two Arctic shelf ecosystems: Journal of Marine Systems, v. 109–110, p. 43–68.

Huntington, H.P., 2000, Traditional knowledge of the ecology of belugas, *Delphinapterus leucas*, in Cook Inlet, Alaska: Marine Fisheries Review, v. 62, p. 134–140.

Huse, G., and Ellingsen, I., 2008, Capelin migrations and climate change—A modeling analysis: Climatic Change, v. 87, no. 1–2, p. 177–197.

Ianelli, J.N., Wilderbuer, T.K., and Nichol, Dan, 2007, Assessment of Greenland turbot in the eastern Bering Sea and Aleutian Islands: North Pacific Fisheries Management Council, Bering Sea and Aleutian Islands SAFE Document, p. 513–562, accessed April 6, 2014, at http://www.afsc. noaa.gov/refm/docs/2007/BSAIturbot.pdf.

Imsland, A.K., Foss, A., Naevdal, G., Johansen, T., Folkvord, A., Stefansson, S.O., and Jonassen, T.M., 2004, Variations in growth in haemoglobin genotypes of Atlantic cod: Fish Physiology and Biochemistry, v. 30, no. 1, p. 47–55.

International Pacific Halibut Commission, 1998, The Pacific halibut—Biology, fishery, and management: Seattle, Washington, International Pacific Halibut Commission Technical Report, no. 40.

Irvine, J.R., Macdonald, R.W., Brown, R.J., Godbout, L., Reist, J.D., and Carmack, E.C., 2009, Salmon in the Arctic and how they avoid lethal low temperatures: North Pacific Anadromous Fish Commission, Bulletin no. 5, p. 39–50.

Irvine, G., and Meyer, R., 1989, Overview of the fishes occurring in the Alaska Arctic, *in* Meyer, R.M., and Johnson, T.M., eds., Fishery oceanography—A comprehensive formulation of technical objectives for offshore application in the Arctic: Anchorage, Alaska, U.S. Department of the Interior, Minerals Management Service, OCS Study MMS 88-0042. p. 6–25.

Isakson, J.S., Houghton, J.P., Rogers, D.E., and Parker, S.S., 1986, Fish use of inshore habitats north of the Alaska Peninsula, June–September 1984 and June–July 1985— Final report: U.S. Department of the Interior, Minerals Management Service, Alaska OCS Region, OCS EIS/EA, 380 p.

Jakobsson, M., Polyak, L., Edwards, M., Kleman, J., and Coakley, B., 2008, Glacial geomorphology of the Central Arctic Ocean—The Chukchi Borderland and the Lomonosov Ridge: Earth Surface Processes and Landforms, v. 33, p. 526–545. Jarvela, L.E., and Thorsteinson, L.K., 1999, The epipelagic fish community of Beaufort Sea coastal waters, Alaska: Arctic, v. 52, no. 1, p. 80–94.

Jay, C.V., Marcot, B.G., and Douglas, D.C., 2011, Projected status of the Pacific walrus (*Odobenus rosmarus divergenes*) in the twenty-first century: Polar Biology, v. 34, no. 7, p. 1,065–1,084.

Jensen, A.S., 1948, A contribution to the ichthyofauna of Greenland: Spoila Zoologica Musei Hauniensis, v. 9, p. 27–57.

Jewett, S.C., and Feder, H.M., 1980, Autumn food of adult starry flounders, *Platichthys stellatus*, from the northeastern Bering Sea and the southeastern Chukchi Sea: Journal de Conseil International pour l'Exploration de la Mer., v. 39, no. 1, p. 7–14.

Johnson, L., 1997, Living with uncertainty, *in* Reynolds, J., ed., Fish ecology in Arctic North America: American Fisheries Society Symposium 19, p. 340–345.

Johnson, M.L., Fiscus, C.H., Ostenson, B.T., and Barbour, M.L., 1966, Marine mammals, *in* Wilimovsky, N.J., and Wolfe, J.N., eds., Environment of the Cape Thompson Region, Alaska: Oak Ridge, Tennessee, United States Atomic Energy Commission, Division of Technical Information, p. 877–924.

Johnson, S.W, Neff, A.D., Lindeberg, J.F., and Maselko, J.M., 2012, Atlas of nearshore fishes of Alaska—A synthesis of marine surveys from 1998 to 2011: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Technical Memorandum NMFS-AFSC-239, 261 p.

Johnson, S.W., Neff, A.D., and Thedinga, J.F., 2005, An atlas on the distribution and habitat of common fishes in shallow nearshore waters of southeastern Alaska: Alaska Fisheries Science Center, Technical Memorandum NMFS-AFSC-157, 98 p.

Johnson, S.W., Thedinga, J.F., and Lindeberg, M.R., 2012, Nearshore fish atlas of Alaska: National Oceanic and Atmospheric Administration Fisheries, accessed February 2012 at http://www.fakr.noaa.gov/habitat/fishatlas/.

Johnson, S.W., Thedinga, J.F., Neff, A.D., and Hoffman, C.A., 2010, Fish fauna in nearshore waters of a Barrier Island in the western Beaufort Sea, Alaska: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Technical Memorandum, NMFSAFSC-210, 28 p.

Johnson, T.P., and Johnston, I.A., 1991, Temperature adaptation and the contractile properties of live muscle fibres from teleosts fish: Journal of Comparative Physiology B, v. 161, no. 1, p. 27–36. Johnston, I.A., 1985, Sustained force development— Specializations and variation among the vertebrates: Journal of Experimental Biology, v. 115, p. 239–251.

Johnston, I.A., and Altringham, J.D., 1985, Evolutionary adaptation of muscle power output to environmental temperature—Force-velocity characteristics of skinned fibres isolated from Antarctic, temperate and tropical marine fish: Pflügers Archiv, v. 405, no. 2, p. 136–140.

Johnston, I.A., and Battram, J., 1993, Feeding energetics and metabolism in demersal fish species from Antarctic, temperate and tropical environments: Marine Biology, v. 115, no. 1, p. 7–14.

Johnston, I.A., Calvo, J., Guderley, H., Fernandez, D., and Palmer, L., 1998, Latitudinal variation in the abundance and oxidative capacities of muscle mitochondria in perciform fishes: The Journal of Experimental Biology, v. 201, no. 1, p. 1–12.

Jones, M.L., and Den Beste, J., 1977, Tuft Point and adjacent coastal areas fisheries projects: Calgary, Alberta, Aquatic Environments Ltd., 152 p.

Jordan, D.S., 1891, Relations of temperature to vertebrae among fishes: Proceedings of The United States National Museum: v. 14, no. 845, p. 107–120.

Jordan, D.S., and Evermann, B.W., 1896, The fishes of North and Middle America—A descriptive catalogue of the species of fish-like vertebrates found in the waters of North America, North of the Isthmus of Panama, part 1: Bulletin of the United States National Museum, no. 47, 1,240 p.

Jørgensen, O.A., 1997, Pelagic occurrence of Greenland halibut, *Reinhardtius hippoglossoides* (Walbaum), in west Greenland waters: Journal of Northwest Atlantic Fisheries Science, v. 21, p. 39–50.

Junquera, S., Román, E., Morgan, J., Sainza, M., and Ramilo, G., 2003, Time scale of ovarian maturation in Greenland halibut (*Reinhardtius hippoglossoides*, Walbaum): ICES Journal of Marine Science, v. 60, no. 4, p. 767–773.

Kajimura, H., and Loughlin, T.R., 1988, Marine mammals in the oceanic food web of the eastern subarctic Pacific: University of Tokyo, Bulletin of the Ocean Research Institute, v. 26, no. 2, p. 187–233.

Karamushko, L.I., Shatunovskii, M.I., and Christiansen, I. Sh., 2004, Metabolic rate and metabolic adaptations in fishes of different latitudes: Journal of Ichthyology, v. 44, p. 647–654.

Karpenko, V.I., and Vasilets, P.M., 1996, Biology of smelt (Osmeridae) in the Korf-Karagin coastal area of the southwestern Bering Sea, *in* Mathisen, O.A., and Coyle, K.O., eds., Ecology of the Bering Sea—A review of Russian literature: Fairbanks, University of Alaska, Alaska Sea Grant Program, AK-SG-96-01, p. 217–235.

Karpov, A.K., and Novikov, G.G., 1981, Hemoglobin alloforms in cod, *Gadus morhua* (Gadiformes, Gadidae), their functional characteristics and occurrence in populations: Journal of Ichthyology, v. 6, p. 45–49.

Kartavtsev, Y.P., Sharina, S.N., Goto, T., Chichvarkhin, A.Y., Balanov, A.A., Vinnikoc, K.A., Invankov, V.N., and Hanzawa, N., 2008, Cytochrome oxidase 1 gene sequence analysis in six flatfish species (Teleostei Pleuronectidae) of Far East Russia with inferences in phylogeny and taxonomy: DNA Sequence v. 19, no. 6, p. 479–489.

Katsanevakis, S., 2006, Modelling fish growth—Model selection, multi-modal inference and model selection uncertainty: Fisheries Research v. 81, p. 229–235.

Kawasaki, T., 1980, Fundamental relations among the selections of life history in marine teleosts: Bulletin of the Japanese Society of Scientific Fisheries, v. 46, no. 3, p. 289–293.

Kendel, R.E., Johnston, R.A.C., Kozak, M.D., and Lobsiger, U., 1974, Movements, distribution, populations and food habits of fish in the western coastal Beaufort Sea: Department of the Environment, Canada, Interim Report of Beaufort Sea Project, Study B1, 64 p.

Kendel, R.E., Johnston, R.A.C., Lobsiger, U., and Kozak, M.D., 1975, Fishes of the Yukon coast: Victoria, British Columbia, Department of the Environment (Canada), Beaufort Sea Project, Technical Report 6, 114 p.

Kennett, J.P., 1982, Marine geology: Englewood Cliffs, New Jersey, Prentice-Hall, 813 p.

Ketchen, K.S., 1975, Age and growth of dogfish *Squalus acanthias* in British Columbia waters: Journal of the Fisheries Research Board of Canada, v. 32, no. 1, p. 43–59.

Kim, S.T., 2001, Winter migrations of shelf fish to the continental slope: Journal of Ichthyology, v. 41, no. 8, p. 564–574.

Kim, S.T., and Biryukov, I.A., 1998, Distribution and some features of the biology of *Reinhardtius hippoglossoides matsuurae* (Pleuronectidae) and *Sebastolobus macrochir* (Scorpaenidae) from the eastern coast of Sakhalin: Journal of Ichthyology, v. 38, no. 1, p. 143–146. Kim, S.T., Biryukov, I.A., and Pometeev, E.V., 1999, Species composition, distribution, and structural changes in the flatfish community of the Tatarskii Strait during the fishing period 1994–1997: Journal of Ichthyology, v. 39, no. 6, p. 433–440.

Kim, S.T., and Shepeleva, O.N., 2001, The structure of shelf ichthyocenoses of northeastern Sakhalin and Terpeniya Bay: Journal of Ichthyology, v. 41, no. 9, p. 711–722.

Kinder, T.H., and Schumacher, J.D., 1981, Circulation over the continental shelf of the southeastern Bering Sea, *in* Hood, D.W., and Calder, J.A., eds., The Eastern Bering Sea shelf—Oceanography and resources, vol. 1: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Office of Marine Pollution Assessment, p. 53–75.

King, J.R., and McFarlane, G.A., 2003, Marine fish life history strategies—Applications to fishery management: Fisheries Management and Ecology, v. 10, no. 4, p. 249–264.

Knipowitsch, N., 1906, Ichthyologische Untersuchungen im Eismeer, I. Lycodes und Lycenchelys: Mémoires de l'Academie Imperiale des Sciences de Saint Petersbourg, v. 19, no. 1, p. 1–130, 1 pl.

Kobelev, E.A., 1989, Some biological characteristics of the Arctic flounder, *Liopsetta glacialis*, in the southeastern area of the Barents Sea: Journal of Ichthyology, v. 29, no. 6, p. 16–21.

Kodolov, L.S., and Matveychuk, S.P., 1995, Stock condition of Greenland turbot (*Reinhardtius hippoglossoides matsuurae* Jordan et Snyder) in the northwestern Bering Sea: Anchorage, Alaska Sea Grant Program AK-SG-95-04.

Kolpakov, N.V., 2005a, Diversity and seasonal dynamics of ichthyocenosis of the Circumlittoral of Russkaya Bight (Northern Primor's): Journal of Ichthyology, v. 45, no. 9, p. 744–753.

Kolpakov, N.V., 2005b, On the biology of *Platichthys stellatus* (Pleuronectidae) in coastal waters of northern Primorye: Journal of Ichthyology, v. 45, no. 8, p. 594–606.

Kostow, K., 2002, Oregon lampreys—Natural history, status, and analysis of management issues—Information reports: Corvallis, Oregon Department of Fish and Wildlife, Oregon State University, Klamath Waters Digital Library, accessed at http://digitallib.oit.edu/cdm/ref/collection/kwl/id/3483.

Knight, C.A., Cheng, C.C., and DeVries, A.L., 1991, Adsorption of alpha-helical antifreeze peptides on specific ice crystal surface planes: Biophysical Journal, v. 59, no. 2, p. 409–418.

Krogh, A., 1916, The respiratory exchange of animals and man: London, Longmans, 173 p. Krylov, A.A., Andreeva, I.A., Vogt, C., Backman, J., Krupskaya, V.V., Grikurov, G.E., and others, 2008, A shift in heavy and clay mineral provenance indicates a middle Miocene onset of a perennial sea ice cover in the Arctic Ocean: Paleoceanography 23, PA1S06.

Kucheryavyi, A.V., Savvaitova, K.A., Gruzdeva, M.A., and Pavlov, D.S., 2007, Sexual dimorphism and some special traits of spawning behavior of the Arctic lamprey *Lethenteron camtschaticum*: Journal of Ichthyology, v. 47, no. 7, p. 481–485.

Kucheryavyi, A.V., Savvaitova, K.A., Pavlov, D.S., Gruzdeva, M.A., Kuzishchin, K.V., and Stanford, J.A., 2007, Variations of life history strategy of the Arctic lamprey *Lethenteron camtschaticum* from the Utkholok River (Western Kamchatka): Journal of Ichthyology, v. 47, no. 1, p. 37–52.

Kulka, D., Hood, C., and Huntington, J., 2007, Recovery strategy for northern wolffish (*Anarhichas denticulatus*) and spotted wolffish (*Anarhichas minor*), and management plan for Atlantic Wolffish (*Anarhichas lupus*) in Canada: St. John's, Fisheries and Oceans Canada, Newfoundland and Labrador Region, 103 p.

Kupriyanov, S.V., 1996, Distribution and biological indices of yellowfin sole (*Pleuronectes asper*) in the southwestern Bering Sea, *in* Mathisen, O.A., and Coyle, K.O., eds., Ecology of the Bering Sea—A review of Russian literature: Fairbanks, University of Alaska, Alaska Sea Grant Report no. 96-01, 306 p.

Lacho, G., 1991, Stomach content analyses of fishes from Tuktoyaktuk Harbour, N.W.T., 1981: Winnipeg, Manitoba, Canadian Data Report of Fisheries and Aquatic Sciences, Central and Arctic Region, Department of Fisheries and Oceans, no. 853, 15 p.

Laevastu, T., Marasco, R., Bax, N., Honkalehto, T., Fredin, R., Fukuhara, F., Gallagher, A., Ingraham, J., Livingston, P., Miyahara, R., and Pola, N., 1985, Evaluation of the effects of oil development on commercial fisheries in the eastern Bering Sea: Anchorage, Alaska, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Outer Continental Shelf Environmental Assessment Program, Final Report for Research Unit 643, 47 p.

Lang, G.M., Livingston, P.A., and Dodd, K., 2005, Groundfish food habits and predation on commercially important prey species in the eastern Bering Sea from 1997 through 2001: U.S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Memorandum NMFS-AFSC-158, 249 p. Lang, G.M., Livingston, P.A., Pacunski, R.E., Parkhurst, J., and Yang, M.-S., 1991, Groundfish food habits and predation on commercially important prey species in the eastern Bering Sea from 1984 to 1986: U.S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Memorandum NMFS F/NWC-207, 240 p.

Lannig, G., Eckerle, L.G., Serendero, I., Sartoris, F.-J., Fischer, T., Knust, R., Johansen, T., and Pörtner, H.-O., 2003, Temperature adaptation in eurythermal cod (*Gadus morhua*)—A comparison of mitochondrial enzyme capacities in boreal and Arctic populations: Marine Biology, v. 142, no. 3, p. 589–599.

Lawrence, M.J., Lacho, G., and Davies, S., 1984, A survey of the coastal fishes of the southeastern Beaufort Sea: Canadian Technical Report of Fisheries and Aquatic Sciences, no. 1220, 178 p.

Lecklin, T., Nash, G., and Egginton, S., 1995, Do fish acclimated to low temperature improve microcirculatory perfusion by adapting red cell rheology: Journal of Experimental Biology, v. 198, no. 8, p. 1,801–1,808.

Leong, H., Brander, K., Carmack, E., Denisenko, S., Drinkwater, K., Hansen, B., Kovacs, K., Livingston, P., McLaughlin, F., and Sakshaug, E., 2005, Marine systems, *in* Arctic climate impact assessment: Cambridge, United Kingdom, Cambridge University Press, p. 453–538.

Levin, P.S., Fogarty, M.J., Murawski, S.A., and Fluharty, D., 2009, Integrated ecosystem assessments—Developing the scientific basis for ecosystem-based management of the ocean: PLoS ONE, v. 7, no. 1, p. 23–28.

LGL Limited, 2014, LGL and Greeneridge publications on fish: LGL Limited Web page, accessed May 15, 2014, at http://lgl.com/en/fish.

Lindeman, R.L., 1942, The trophic-dynamic aspect of ecology: Ecology, v. 23, p. 399–418.

Litzow, M.A., 2007, Alaska Fisheries Science Center quarterly research report, January–March, 2007: National Oceanic and Atmospheric Administration Fisheries, Alaska Fisheries Science Center Web page, accessed April 2, 2014, at http:// www.afsc.noaa.gov/Quarterly/jfm2007/divrptsRACE4.htm.

Litzow, M.A., Piatt, J.F., Prichard, A.K., and Roby, D.D., 2002, Response of pigeon gullemots to variable abundance of high-lipid and low-lipid prey: Oecologia, v. 132, no. 2, p. 286–295.

Livingston, P.A., Ward, A., Lang, G.M., and Yang, M.-S., 1993, Groundfish food habits and predation on commercially important prey species in the eastern Bering Sea from 1987 to 1989: National Oceanic and Atmospheric Administration Technical Memorandum NMFS-AFSC-11, 202 p.

Logerwell, E., Busby, M., Carothers, C., Cotton, S., Duffy-Anderson, J., Farley, E., Goddard, P., Heintz, R., Holladay, B., Horne, J., Johnson, S., Lauth, B., Moulton, L., Neff, D., Norcross, B., Parker-Stetter, S., Seigle, J., and Sformo, T., 2015, Fish communities across a spectrum of habitats in the western Beaufort Sea and Chukchi Sea: Progress in Oceanography, no. 136, p. 115–132.

Logerwell, E., Rand, K., Parker-Setter, S., Horne, J.,
Weingartner, T., and Bluhm, B., 2009, Beaufort Sea marine fish monitoring 2008—Pilot survey and test of hypotheses: Anchorage, Alaska, Final Report from U.S. Department of Commerce, National Oceanic and Atmospheric Administration/National Marine Fisheries Service to U.S. Department of the Interior, Minerals Management Service, Alaska OCS Region, OCE Study MMS 2008-062, 258 p.

Logerwell, E., Rand, K., and Weingartner, T.J., 2011, Oceanographic characteristics of the habitat of benthic fish and invertebrates in the Beaufort Sea: Polar Biology, v. 34, p. 1,783–1,796.

Loher, T., 2012, Potential mechanisms, research avenues, and management action associated with size at age and growth of Pacific halibut, *in* IPHC Report of Assessment and Research Activities 2012: International Pacific Halibut Commission (IPHC) online report, p. 457–486, accessed February 16, 2014, at http://www.iphc.int/publications/ rara/2012/rara2012457_sizeatage.pdf.

Loher, T., and Seitz, A.C., 2006, Seasonal migration and environmental conditions of Pacific halibut *Hippoglossus stenolepis*, elucidated from pop-up archival transmitting (PAT) tags: Marine Ecology Progress Series, v. 317, p. 259–271.

Longshan, L., Yongjun, C., Yungchih, L., Jing, Z., Punqing, S., Xingguang, Y., Risheng, W., and Kwang-tsao, S., 2014, Composition of fish species in the Bering and Chukchi Seas and their responses to changes in the ecological environment, Acta Oceanologica Sinica, v. 33, no. 6, p. 63-73.

Lourie, S.A., and Vincent, A.C.J., 2004, Using biogeography to help set priorities in marine conservation: Conservation Biology, v. 18, no. 4, p. 1,004–1,020.

Love, M.S., 2011, Certainly more than you wanted to know about the fishes of the Pacific coast: Santa Barbara, Calif., Really Big Press, 649 p. Love, M.S., Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2005, Resource inventory of marine and estuarine fishes of the West Coast and Alaska—A checklist of North Pacific and Arctic Ocean species from Baja California to the Alaska-Yukon border: U.S. Geological Survey, OCS Study MMS 2005-030 and USGS/NBII 2005-001, 276 p.

Low, W.-K., Lin, Q., Ewart, K.V., Fletcher, G. L., and Hew, C.L., 2002, The skin-type antifreeze polypeptides—A new class of Type I AFPs, *in* Ewart, K.V., and Hew, C.L., eds., Molecular aspects of fish and marine biology, volume 1— Fish antifreeze proteins: New Jersey, World Scientific, p. 161–186.

Lowry, L.F., and Frost, K.J., 1981, Distribution, growth, and foods of Arctic cod (*Boreogadus saida*) in the Bering, Chukchi, and Beaufort Seas: Canadian Field-Naturalist, v. 95, no. 2, p. 186–191.

Lowry, L.F., Frost, K.J., and Burns, J.J., 1980, Variability in the diet of ringed seals, *Phoca hispida*, in Alaska: Canadian Journal of Fisheries and Aquatic Sciences, v. 37, no. 12, p. 2,254–2,261.

Luksenburg, J.A., Pedersen, T., and Falk-Petersen, I.-B., 2004, Reproduction of the shorthorn sculpin *Myoxocephalus scorpius* in northern Norway: Journal of Sea Research, v. 51, no. 2, p. 157–166.

Luzier, C.W., Schaller, H.A., Brostrom, J.K., Cook-Tabor, C., Goodman, D.H., Nelle, R.D., and others, 2011, Pacific Lamprey (*Entosphenus tridentatus*)—Assessment and template for conservation measures: U.S. Fish and Wildlife Service, 282 p.

Lynghammar, A., Christiansen, J.S., Mecklenburg, C.W., Karamushko, O.V., Møller, P.R., Gallucci, V.F., and Moller, P.R., 2012, Species richness and distribution of chondrichthyan fishes in the Arctic Ocean and adjacent seas: Biodiversity, v. 14, no. 1, p. 57–66.

Macdonald, J.A., 1981, Temperature compensation in the peripheral nervous systems—Antarctic vs temperate poikilotherms: Journal of Comparative Physiology, v. 142, p. 411–418.

Macdonald, J.A., and Montgomery, J.C., 1982, Thermal limits of neuromuscular function in an Antarctic fish: Journal of Comparative Physiology, v. 147, p. 237–250.

Macdonald, J.A., and Montgomery, J., 2005, The nervous system, *in* Farrell, A.P., and Steffensen, J.F., eds., The physiology of polar fishes: Amsterdam, Elsevier, p. 351–383.

Magdanz, J.S., Smith, H., Braem, N., Fox, P., and Koster, D.S., 2011, Patterns and trends in subsistence fish harvests, Northwest Alaska, 1994–2004: Alaska Department of Fish and Game, Technical Paper no. 366, 104 p.

Majewski, A.R., Sareault, J.E., and Reist, J., 2006, Fish catch data from offshore sites in the Mackenzie River estuary and Beaufort Sea during the open water season, August 2004 aboard the CCGS Nahidik: Winnipeg, Manitoba, Fisheries and Oceans Canada, no. 2771, 42 p.

Maksimenkov, V.V., and Tokranov, A.M., 1993, The diet of the pond smelt, *Hypomesus olidus*, in the Bol'shaya River Estuary (Western Kamchatka): Journal of Ichthyology, v. 33, no. 9, p. 11–21.

Mann, G., 1974, Life history types of the least cisco (*Coregonus sardinella*, Valenciennes) in the Yukon Territory, North Slope and eastern Mackenzie River Delta drainages, *in* McCart, P.J., ed., Life histories of three species of freshwater fishes in Beaufort Sea drainages, Yukon Territory,: Arctic Gas Biological Report Series, v. 18, no. 3, 132 p.

Markon, C.J., Trainor, S.F., and Chapin, F.S., III, eds., 2012, The United States National Climate Assessment—Alaska Technical Regional Report: U.S. Geological Survey Circular 1379, 148 p.

Marliave, J.B., 1975, The behavioral transformation from the planktonic larval stage of some marine fishes reared in the laboratory: Vancouver, University of British Columbia, Ph.D. dissertation.

Marshall, N.B. 1953, Egg size in Arctic, Antarctic and deepsea fishes: Evolution, v. 7, p. 328–341.

Martin, D.J., Glass, D.R., Whitmus, C.J., Simenstad, C.A., Milward, D.A., Volk, E.C., Stevenson, M.L., Nunes, P., Savoie, M., and Grotefendt, R.A., 1986, Distribution, seasonal abundance, and feeding dependencies of juvenile salmon and non-salmonid fishes in the Yukon River Delta: Outer Continental Shelf Environmental Assessment Program, Reports of Principal Investigators, U.S. Department of Commerce and U.S. Department of the Interior, 388 p.

Martin, M.H., and Clausen, D.M., 1995, Data report—1993 Gulf of Alaska Bottom Trawl Survey: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, NOAA Technical Memorandum, NMFS-AFSC-59, 217 p.

Marz, S., 2010, Arctic sea ice ecosystem—A summary of species that depend on and associate with sea ice and projected impacts from sea ice changes: Report to the Arctic Council Working Group on the Conservation of Arctic Flora and Fauna, 64 p. Maslenikov, K.P., Orr, J.W., and Stevenson, D.E., 2013, Range extensions and significant distributional records for eight-two species of fishes in Alaska marine waters: Northwestern Naturalist, v. 94, no. 1, p. 1–21.

Matarese, A.C., Blood, D.M., Picquelle, S.J., and Benson, J.L., 2003, Atlas of abundance and distribution patterns of ichthyoplankton from the northeast Pacific Ocean and Bering Sea ecosystems based on research conducted by the Alaska Fisheries Science Center (1972–1996): National Oceanic and Atmospheric Administration Professional Paper NMFS 1, 281 p.

Matarese, A.C., Kendall, J., Arthur W., Blood, D.M., and Vinter, B.M., 1989, Laboratory guide to early life history stages of northeast Pacific fishes: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service Technical Report NMFS 80, 652 p.

Mathis, J.T., and Questel, J.M., 2013, Assessing seasonal changes in carbonate parameters across small spatial gradients in the Northeastern Chukchi Sea: Continental Shelf Research, v. 67, p. 42–51, accessed February 3, 2014, at http://www.sciencedirect.com/science/article/pii/ S0278434313001416.

Maule, A.G., and Thorsteinson, L.K., 2012, Feasibility plan to assess the effects of dispersed oil on Arctic Cod: Final report prepared by the U.S. Department of the Interior, U.S. Geological Survey for the Bureau Ocean Energy Management, OCS Study BOEM 2011-062, 52 p.

McAllister, D.E., 1959, Records of marine fishes from fresh water in British Columbia: The Canadian Field-Naturalist, v. 73, no. 1, p. 13–14.

McAllister, D.E., 1975, Ecology of the marine fishes of Arctic Canada, in Proceedings of the Circumpolar Conference on Northern Ecology, September 15–18, 1975: Ottawa, National Research Council of Canada, p. II-49–II-65.

McAllister, D.E., Anderson, M.E., and Hunter, J.G., 1981, Deep-water eelpouts, Zoarcidae, from Arctic Canada and Alaska: Canadian Journal of Fisheries and Aquatic Sciences, v. 38, no. 7, p. 821–839.

McArdle, H.J., and Johnston, I.A., 1980, Evolutionary temperature adaptation in fish muscle sarcoplasmic reticulum: Journal of Comparative Physiology, v. 135B, p. 157–164.

McCart, P.J., Craig, P.C., and Bain, H., 1972, Report of fisheries investigations in the Sagavanirtok River and neighboring drainages: Bellevue, Wash., Aquatic Environments Ltd., Report to Alyeska Pipeline Service Company, 165 p.

McConnaughey, T., and McRoy, C.P., 1979, ¹³C label identifies eelgrass (*Zostera marina*) carbon in an Alaskan estuarine food web: Marine Biology, v. 53, no. 3, p. 263–269.

McDowall, R.M., 1987, The occurrence and distribution of diadromy among fishes: American Fisheries Society Symposium, no. 1, p. 1–13.

McDowall, R.M., 2008, Jordan's and other ecogeographical rules, and the vertebral number in fishes: Journal of Biogeography, v. 35, p. 501–508.

McGregor, S.B., 1990, Seasonal and ontogenetic changes in meiofauna in the diets of postmetamorphic flatfishes: University of Alaska, Fairbanks, Master's thesis.

McKechnie, R.J., and Fenner, R.B., 1971, Food habits of white sturgeon, *Acipenser transmontanus* in San Pablo and Suisun Bays, California: California Fish and Game, v. 57, p. 209–212.

McKinney, M.L., 1997, Extinction vulnerability and selectivity—Combining ecological and paleontological views: Annual Review of Ecology and Systematics, v. 28, p. 495–516.

McLaren, I.A., 1958, The biology of the ringed seal (*Phoca hispida* Schreber) in the Eastern Canadian Arctic: Bulletin Fisheries Research Board of Canada 118, 97 p.

McPhail, J.D., and Lindsey, C.C., 1970, Freshwater fishes of northwestern Canada and Alaska: Bulletin of the Fisheries Research Board of Canada Bulletin 173, 381 p.

Mecklenburg, C.W., 2003a, Family Anarhichadidae Bonaparte 1846—Wolffishes: California Academy of Sciences Annotated Checklists of Fishes, no. 10, 6 p.

Mecklenburg, C.W., 2003b, Family Hemitripteridae Gill 1872—Sea ravens or sailfin sculpins: California Academy of Sciences Annotated Checklists of Fishes no. 5, 6 p.

Mecklenburg, C.W., 2003c, Family Zaproridae Jordan and Evermann 1898—Prowfishes: California Academy of Sciences Annotated Checklists of Fishes No. 13.

Mecklenburg, C.W., Byrkjedal, I., Christiansen, J.S., Karamushko, O.V., Lynghammar, A., and Møller, P.R., 2013, List of marine fishes of the Arctic Region annotated with common names and zoogeographic characterizations: Akureyri, Iceland, Conservation of Arctic Flora and Fauna, 35 p.

Mecklenburg, C.W., Byrkjedal, I., Karamushko, O.V., and Møller, P.R., 2014, Atlantic fishes in the Chukchi Borderland: Marine Biodiversity, v. 44, no. 1, p. 127–150, doi 10./1007/s12526-013-0192-1. Mecklenburg, C.W., and Mecklenburg, T.A., 2009, Arctic marine fish museum specimens, 2nd ed., Metadata report and database submitted to ArcOD, Institute of Marine Science: University of Alaska, Fairbanks, by Point Stephens Research, metadata report accessed August 8, 2012, at http://www.arcodiv.org/Database/Fish_datasets.html.

Mecklenburg, C., and Mecklenburg, T.A., 2011a, Longhead Dab—*Limanda proboscidea* Gilbert, 1896: Arctic Ocean Diversity—Census of Marine Life, accessed April 2011 at http://www.arcodiv.org/Fish/Limanda proboscidea.html.

Mecklenburg, C.W., and Mecklenburg, T.A., 2011b, Polar Eelpout—*Lycodes polaris* (Sabine, 1824): Arctic Ocean Diversity Web site, accessed 2011 at http://www.arcodiv. org/Fish/Lycodes_polaris.html.

Mecklenburg, C.W., Mecklenburg, T.A., Sheiko, B.A., and Chernova, N.V., 2006, Arctic marine fish museum specimens—Metadata report and database submitted to ArcOD, Institute of Marine Science: University of Alaska Fairbanks by Point Stephens Research, metadata report, http://dw.sfos.uaf.edu/rest/metadata/ArcOD/2007F1.

Mecklenburg, C.W., Mecklenburg, T.A., Sheiko, B.A., and Steinke, D., 2016, Pacific Arctic marine fishes: Akureyri, Iceland, Conservation of Arctic Flora and Fauna, Monitoring Series Report No. 23, 406 p., accessed May 10, 2016, at http://caff.is/monitoring-series/370-pacific-arcticmarine-fishes.

Mecklenburg, C.W., Mecklenburg, T.A., and Thorsteinson, L.K., 2002, Fishes of Alaska: Bethesda, Maryland, American Fisheries Society, 1,116 p.

Mecklenburg, C.W., Møller, P.R., and Steinke, D., 2011, Biodiversity of Arctic marine fishes—Taxonomy and zoogeography: Marine Biodiversity, v. 41, no. 1, p. 109– 140, Online Resources 1–5.

Mecklenburg, C.W., Norcross, B.L., Holladay, B.A., and Mecklenburg, T.A., 2008, Fishes, *in* Hopcroft, R., Bluhm, B., and Gradinger, R., eds., Arctic Ocean Synthesis— Analysis of climate change impacts in the Chukchi and Beaufort Seas with strategies for future research: Fairbanks, Alaska, Institute of Marine Science, University of Alaska, p. 66–79.

Mecklenburg, C.W., and Sheiko, B.A., 2003, Family Cyclopteridae Bonaparte 1831—Lumpsuckers: California Academy of Sciences Annotated Checklists of Fishes no. 6, 17 p.

Mecklenburg, C.W., Stein, D.L., Sheiko, B.A., Chernova, N.V., Mecklenburg, T.A., and Holladay, B.A., 2007, Russian–American long-term census of the Arctic—Benthic fishes trawled in the Chukchi Sea and Bering Strait, August 2004: Northwestern Naturalist, v. 88, no. 3, p. 168–187. Mecklenburg, C.W., and Steinke, D., 2015, Ichthyofaunal baselines in the Pacific Arctic region and RUSALCA study area, Oceanography, v. 28, no. 3, p. 158–189.

Mellilo, J.M., Richmond, T.C., and Yohe, G.W., eds., 2014, Climate change impacts in the United States—The third annual national climate assessment: U.S. Global Climate Research Program, 841 p.

Meyer Ottesen, C.A., Hop, H., Christiansen, J.S., and Falk-Petersen, S., 2011, Early life history of the daubed shanny (Teleostei: *Leptoclinus maculatus*) in Svalbard waters: Marine Biodiversity, v. 41, no. 3, p. 383–394.

Mikhail, M.Y., and Welch, H.E., 1989, Biology of Greenland cod, *Gadus ogac*, at Saqvaqjuac, northwest coast of Hudson Bay: Environmental Biology of Fishes, v. 26, p. 49–62.

Miller, B.S., 1967, Stomach contents of adult starry flounder and sand sole in East Sound, Orcas Island, Washington: Journal of the Fisheries Research Board of Canada, v. 24, no. 12, p. 2,515–2,521.

Miller, B.S., and Kendall, A.W., 2009, Early life history of marine fishes: Berkley, University of California Press, 437 p.

Miller, B.S., Simenstad, C.A., Moulton, L.L., Fresh, K.L., Funk, F.C., Karp, W.A., and others, 1977, Puget Sound baseline program—Nearshore fish survey: University of Washington, Fishery Research Institute, FRI-UW-7710, 219 p.

Miller, D.J., and Lea, R.N., 1976, Guide to the coastal marine fishes of California: California Fisheries Bulletin no. 157, 255 p.

Miller, D.J., and Schmidtke, J., 1956, Report on the distribution and abundance of Pacific herring (*Clupea pallasi*) along the coast of central and southern California: California Fish and Game, v. 42, no. 3, p. 163–187.

Miller, J.M., Burke, J.S., and Fitzhugh, G.R., 1991, Early life history patterns of Atlantic North Atlantic American flatfish—Likely (and unlikely) factors controlling recruitment: Netherlands Journal of Sea Research, v. 27, p. 261–275.

Miller, T.J., 2007, Contribution of individual-based coupled physical-biological models to understanding recruitment in marine fish populations: Marine Ecology Progress Series, v. 347, p. 127–138.

Milliman, J.D., and Meade, R.H., 1983, World-wide delivery of river sediment to the oceans: Journal of Geology, v. 91, no. 1, p. 1–21.

Miner, P., and Rivasplata, A., 1994, Thresholds of significance; criteria for defining significance: Sacramento, Calif., CEQA Technical Advice Series, Governor's Office of Planning and Research, 12 p.

Minerals Management Service, 2008, Beaufort Sea and Chukchi Sea planning areas—Oil and gas lease sales 209, 212, 217, and 221: U.S. Department of the Interior, Minerals Management Service Alaska OCS Region, OCS EIS/EA: MMS 2008-0055, 538 p.

Mineva, T.A., 1964, On the biology of some flatfishes in the eastern Bering Sea, *in* Moiseev, P.A. ed., Soviet fisheries investigations in the northeastern Pacific, Part II: Jerusalem, Israel Program for Scientific Translations, p. 227–235.

Mito, K.-I., Nishimura, A., and Yanagimoto, T., 1999, Ecology of groundfishes in the eastern Bering Sea, with emphasis on food habits, *in* Loughlin, T.R., and Ohtani, K., eds., Dynamics of the Bering Sea: Fairbanks, University of Alaska Sea Grant, p. 537–580.

Moiseev, P.A., 1953, Cod and flounders of far-eastern seas: Izvestiya TINRO, Proceedings Pacific Science Research Institute of Marine Fisheries and Oceanography, v. 40, p. 1–287.

Moles, A., and Norcross, B.L., 1995, Sediment preference in juvenile Pacific flatfishes: Netherlands Journal of Sea Research, v. 34, no. 1–3, p. 177–182.

Møller, P.R., 2000, Restoration of the taxon *Lycodes marisalbi*, with notes on its disjunct Arctic distribution: Journal of Fish Biology, v. 57, no. 6, p. 1,404–1,415.

Møller, P.R., 2001, Redescription of the *Lycodes pallidus* species complex (Pisces, Zoarcidae), with a new species from the Arctic/North Atlantic Ocean: Copea, v. 2001, no. 4, p. 972–996.

Møller, P.R., and Jørgensen, O.A., 2000, Distribution and abundance of eelpouts (Pisces, Zoarcidae) off West Greenland: Sarsia, v. 85, no. 1, p. 23–48.

Møller, P.R., Nielsen, J.G., Knudsen, S.W., Poulsen, J.Y., Sünksen, K., and Jørgensen, O.A., 2010, A checklist of the fish fauna of Greenland waters: Zootaxa, v. 2378, p. 1–84.

Monterio, R.S.O., 2002, Fish growth modeling: growth of the European anchovy (*Engraulis encrasicolus*) in the Tagus Estuary, Portugal: European Diploma Eruditionum Altarum Modelling of the Marine Environment, 53 p.

Montgomery, J.C., and Macdonald, J.A., 1984, Performance of motor systems in Antarctic fishes: Journal of Comparative Physiology, v. 154A, p. 241–248.

Montgomery, J.C., and Sutherland, K.B.W., 1997, Sensory development of the Antarctic silverfish *Pleurogramma antarcticum*—A test for the ontogenetic shift hypothesis: Polar Biology, v. 18, p. 112–115.

Moran, C., and Melan, R., 2001, Temperature-dependent conduction properties in Arctic fish peripheral nerves: Polar Biology, v. 24, p. 9–15.

Mork, J., Giskeödegård, R., and Sundnes, G., 1983, Haemoglobin polymorphism in *Gadus morhua*—Genotypic differences in maturing age and within-season gonad maturation: Helgoländer Meeresunters, v. 36, p. 313–322.

Morris, R.W. and Schneider, M.J., 1969, Brain fatty acids of an Antarctic fish, *Trematomus bernacchii*: Comparatuve Biochemistry and Physiology, v. 28, p. 1,461–1,465.

Morrow, J.E., 1980, The freshwater fishes of Alaska: Anchorage, Alaska Northwest Publishing Company, 248 p.

Moser, H.G., 1996, The early stages of fishes in the California current region: Atlas, California Cooperative Oceanic Fisheries Investigations, no. 33, 1,505 p.

Mouillot, D., Bellwood, D.R., Baraloto, C., Chave, J., and Galzin, R., 2013, Rare species support vulnerable functions in high-diversity ecosystems: PLoS Biology, v. 11, no. 5, e1001569, doi:10.1371/journal.pbio.1001569.

Moulton, L.L., and Penttila, D.E., 2000, Forage fish spawning distribution in San Juan County and protocols for sampling intertidal and nearshore regions: Mount Vernon, Washington, San Juan County Forage Fish Assessment Project, Northwest Straits Commission, 53 p.

Moulton, L.L., Seavey, B., and Pausanna, J., 2010, History of an under-ice subsistence fishery for Arctic Cisco and Least Cisco in the Colville River, Alaska: Arctic, v. 63, no. 4, p. 381–390.

Moulton, L.L., and Tarbox, K.E., 1987, Analysis of Arctic cod movements in the Beaufort Sea nearshore region, 1978–79: Arctic, v. 40, no. 1, p. 43–49.

Moyle, P.B., Kiernan, J.D., Crain, P.K., and Quiñones, R.M., 2013, Climate change vulnerability of native and alien freshwater fishes of California—A systematic assessment approach: PLoS ONE, v. 8, no. 5, e63883, doi:10.1371/ journal.pone.0063883.

Murray, H.M., Hew, C.L., and Fletcher, G.L., 2002, Skin-type antifreeze protein expression in integumental cells of larval winter flounder: Journal of Fish Biology, v. 60, no. 6, p. 1,391–1,406. Murray, H.M., Hew, C.L., and Fletcher, G.L., 2003, Spatial expression patterns of skin-type antifreeze protein in winter flounder (*Pseudopleuronectes americanus*) epidermis following metamorphosis: Journal of Morphology, v. 257, no. 1, p. 78–86.

Musick, J.A., 1999, Criteria to define extinction risk in marine fishes: Fisheries, v. 24, no. 12, p. 6–14.

Mueter, F.J., Bond, N.A., Ianelli, J.N., and Hollowed, A.B., 2011, Expected declines in recruitment of walleye pollock (*Theragra chalcogramma*) in the eastern Bering Sea under future climate change: ICES Journal of Marine Science, v. 68, no. 6, p. 1,284–1,296.

Mueter, F.J., Broms, C., Drinkwater, K.F., Friedland, K.D., Hare, J.A., Hunt, G.L., Jr., Melle, W., and Taylor, M., 2009, Ecosystem response to recent oceanographic variability in high-latitude Northern Hemisphere ecosystems: Progress in Oceanography, v. 81, no. 1–4, p. 93–110.

Mueter, F.J., Lauth, R., Andrews, A., de Robertis, A., Britt, L., Farley, E., Norcross, B., Sigler, M., and Wilson, C., 2013, The marine fish communities of the Bering and Chukchi Sea: Paper presented at the 2013 Sea Grant Lowell Wakefield Symposium, Responses of Arctic Marine Ecosystems to Climate Change, March 26–29, 2013, Anchorage, Alaska.

Mueter, F.J., and Litzow, M.A., 2008, Sea ice retreat alters the biogeography of the Bering Sea continental shelf: Ecological Applications, v. 18, no. 2, p. 309–320.

Mueter, F.J., and Norcross, B.L., 2002, Spatial and temporal patterns in the demersal fish community on the shelf and upper slope regions of the Gulf of Alaska: Fishery Bulletin 100, no. 3, p. 559–581.

Murphy, S.M., Mueter, F.J., and Braund, S.R., 2007, Variation in the abundance of Arctic Cisco in the Colville River— Analysis of existing data and local knowledge, volume 1—Final report: U.S. Department of the Interior, Minerals Management Service, OCS Study MMS 2007-042, 240 p.

Musienko, L.N., 1970, Reproduction and development of Bering Sea fishes, *in* Moiseev, P.A., ed., Soviet fisheries investigations in the northeastern Pacific—part V: Jerusalem, Israel Program for Scientific Translations, p. 161–224.

Nahrgang, J., Camus, L., Broms, F., Christiansen, J.S., and Hop, H., 2010, Seasonal baseline levels of physiological and biochemical parameters in polar cod (*Boreogadus saida*)—Implications for environmental monitoring: Marine Pollution Bulletin, v. 60, no. 8, p. 1,336–1,345. Napazakov, V.V., and Chuchukalo, V.I., 2003, Diet and trophic status of mass species of sculpins (Cottidae) in the western part of the Bering Sea in autumn: Journal of Ichthyology, v. 43, no. 3, p. 236–244.

National Aeronautics and Space Administration, 2012, Understanding earth—The icy Arctic: Greenbelt, Maryland, Goddard Space Flight Center, 12 p.

National Oceanic and Atmospheric Administration, 2008, West coast of North America living marine resources data atlas: Rockville, Maryland, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, Strategic Assessment Branch, Ocean Assessment Division, Office of Oceanography and Marine Assessment, National Ocean Service.

National Oceanic and Atmospheric Administration, 2009, World ocean database 2009: National Oceanographic and Atmospheric Administration, National Centers for Environmental Information database, accessed October 20, 2010, at http://www.nodc.noaa.gov/OC5/WOD09/ pr_wod09.html.

National Oceanic and Atmospheric Administration, 2010, Age and growth program: National Oceanic and Atmospheric Administration, accessed April 2011 at http://www.afsc. noaa.gov/refm/age/Stats/max_age.htm.

National Oceanic and Atmospheric Administration, 2012, Groundfish Assessment Program: National Oceanic and Atmospheric Administration database, accessed November 10, 2013, at http://www.afsc.noaa.gov/RACE/groundfish/ default.php.

National Oceanic and Atmospheric Administration, 2013a, Effects of oil and gas activities in the Arctic Ocean— Supplemental draft environmental impact statement: Silver Spring, Md., U.S. Department of Commerce, National Marine Fisheries Service, Office of Protected Resources, v. 1, chaps. 1–3, 445 p.

National Oceanic and Atmospheric Administration, 2013b, Effects of oil and gas development in the Arctic Ocean, Supplemental Draft Environmental Impact Statement: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, National Marine Fisheries Service, 445 p., accessed April 2, 2013 at http://www.nmfs. noaa.gov/pr/permits/eis/arctic sdeis vol1.pdf.

National Oceanic and Atmospheric Administration., 2014, Summer bottom surface temperatures—Eastern Bering Sea: Alaska Fisheries Science Center, accessed November 24, 2014, at http://access.afsc.noaa.gov/reem/EcoWeb/content/ pdf/EBSSummerBottomTemps.pdf. National Oceanic and Atmospheric Administration, 2014, Alaska Fisheries Science Center: National Oceanic and Atmospheric Administration Fisheries, Web site accessed November 24, 2014, at http://www.afsc.noaa.gov/.

National Research Council, 1996, The Bering Sea ecosystem: Washington D.C., National Academies Press, 307 p.

National Snow and Ice Data Center, 2011a, Monthly sea ice extent for 1980 and 2007: National Snow and Ice Data Center database, accessed April 15, 2011, at http://nsidc. org/.

National Snow and Ice Data Center, 2011b, NSIDC BIST, Compare data, spreadsheet—Sea Ice Index—Extent and concentrations trends: accessed April 15, 2011, at http:// nsidc.org/cgi-bin/bist/bist.pl?config=seaice_extent_ trends&submit=Go%21.

National Snow and Ice Data Center, 2011c, State of the cryosphere—Is the cryosphere sending signals about climate change?: Boulder, Colorado, University of Colorado, accessed April 15, 2011, at http://nsidc.org/sotc/ sea_ice.html.

Naumenko, E.A., 1996, Distribution, biological condition, and abundance of capelin (*Mallotus villosus socialis*) in the Bering Sea, *in* Mathisen, O.A., and Coyle, K.O., eds., Ecology of the Bering Sea: Fairbanks, University of Alaska, Alaska Sea Grant College Program, p. 237–256.

Nazarov, D.Y., Kucheryavyi, A.V., Savvaitova, K.A., Gruzdeva, M.A., Kuzishchin, K.V., and Pavlov, D.S., 2011, Population structure of Arctic lamprey *Lenthenteron camtschaticum* from the Kol River (western Kamchatka): Journal of Ichthyology, v. 51, no. 4, p. 277–290.

Negulescu, P.A., Rubinsky, B., Fletcher, G.L., and Machen, T.E., 1992, Fish antifreeze proteins block Ca entry into rabbit parietal cells: American Journal of Physiology, Cell Physiology, v. 263, p. 1,310–1,313.

Nelson, J.S., 2006, Fishes of the World: Hoboken, New Jersey, John Wiley and Sons, Inc., 416 p.

Nelson, J.S., Crossman, E.J., Espinosa- Pérez, H., Findley, L.T., Gilbert, C.R., Lea, R.N., and Williams, J.D., 2004, Common and scientific names of fishes from the United States, Canada, and Mexico (6th ed.): American Fisheries Society Special Publication 29, 386 p.

Nelson, R., and Bouchard, C., 2013, Arctic Cod (*Boreogadus saida*) population structure and connectivity as examined with molecular genetics: Final Report to the North Pacific Research Board, NRPB Project 1125, 39 p.

Nichol, D.G., 1995, Spawning and maturation of female yellowfin sole in the eastern Bering Sea, *in* Alaska Sea Grant Program, Proceedings of the International Symposium on North Pacific Flatfish: Alaska Sea Grant Program, AK-SG-95-04, p. 35–50.

Nichol, D.G., 1997, Effects of geography and bathymetry on growth and maturity of yellowfin sole, *Pleuronectes asper*, in the eastern Bering Sea: Fishery Bulletin, v. 95, no. 3, p. 494–503.

Nichol, D.G., 1998, Annual and between-sex variability of yellowfin sole, *Pleuronectes asper*, spring-summer distributions in the eastern Bering Sea: Fishery Bulletin, v. 96, no. 3, p. 547–561.

Nichol, D.G., and Acuna, E.I., 2001, Annual and batch fecundities of yellowfin sole, *Limanda aspera*, in the eastern Bering Sea: Fishery Bulletin, v. 99, no. 1, p. 108–122.

Nielsen, J.G., and Fosså, S.A., 1993, *Lycodes adolfi*, a new species of eelpout (Zoarcidae) from Greenland: Cybium, v. 17, p. 39–44.

Nielsen, J.L., Ruggerone, G.T., and Zimmerman, C.E., 2012, Adaptive strategies and life history characteristics in a warming climate—Salmon in the Arctic?: Environmental Biology of Fishes, v. 96, no. 10–11, p. 1,187–1,226.

Nikolsky, G.V., 1963, The ecology of fishes: London, Academic Press, 352 p.

Norcross, B.L., Blanchard, A., and Holladay, B.A., 1999, Comparison of models for defining nearshore flatfish nursery areas in Alaska waters: Fisheries Oceanography, v. 8, p. 50–67.

Norcross, B.L., Holladay, B.A., Busby, M.S., and Mier, K.L., 2010, Demersal and larval fish assemblages in the Chukchi Sea: Deep-Sea Research II, v. 57, no. 1–2, p. 57–70.

Norcoss, B.L., Holladay, B.A., and Mecklenburg, C.W., 2013a, Recent and historical distribution and ecology of demersal fishes in the Chukchi Sea Planning Area: Fairbanks, Alaska, OCS Study BOEM 2012-073, 210 p.

Norcross, B.L., Müter, F.-J., and Holladay, B.A., 1997, Habitat models for juvenile pleuronectids around Kodiak Island, Alaska: Fishery Bulletin, v. 95, no. 3, p. 504–520.

Norcross, B.L., Raborn, S.W., Holladay, B.A., Gallaway, B.J., Crawford, S.T., Priest, J.T., Edenfield, L.E. and Meyer, R., 2013b, Northeastern Chukchi Sea demersal fishes and associated environmental characteristics, 2009–2010, Continental Shelf Research v. 67, p. 77–95.

North Pacific Fishery Management Council, 2009, Fishery management plan for fish resources of the Arctic management area: Anchorage, Alaska, North Pacific Fishery Management Council, 158 p. North Pacific Fishery Management Council, 2010, Fishery management plan for groundfish of the Bering Sea and Aleutian Islands management area—Appendices—Fishery management plan for groundfish of the BSAI Management Area: North Pacific Fisheries Management Council, 233 p.

Novikov, N.P., 1964, Basic elements of the biology of the Pacific halibut (*Hippoglossus hippoglossus stenolepis* Schmidt) in the Bering Sea, *in* Moiseev, P.A., ed., Soviet fisheries investigations in the northeastern Pacific, Part II: Jerusalem, Israel Program for Scientific Translations, p. 175–219.

Nozères, C., Archambault, D., Chouinard, P.-M., Gauthier, J., Miller, R., Parent, E., Schwab, P., Savard, L., and Dutil, J.-D., 2010, Identification guide for marine fishes of the estuary and northern Gulf of St. Lawrence and sampling protocols used during trawl surveys between 2004 and 2008: Canadian Technical Report of Fisheries and Aquatic Sciences no. 2866.

Ochman, S., and Dodson, J.J., 1982, Composition and structure of the larval and juvenile fish community of the Eastman River and estuary, James Bay: Naturaliste Canada, v. 109, p. 803–813.

Odum, E.P., 1985, Ecologia: Rio de Janeiro, Brazil, Interamericana, 434 p.

Odum, W.E., and Heald, E.J., 1975, The detritus-based food web of an estuarine mangrove community, *in* Cronin, L.E., ed., Estuarine research: New York, Academic Press, Inc., p. 265–286.

Ofelt, C.H., 1975, Food habits of nesting bald eagles in southeast Alaska: The Condor, v. 77, p. 337–338.

O'Connell, M., Dillon, M.C., Wright, J.M., Bentzen, P., Merkouris, S.E., and Seeb, J.E., 1998, Genetic structuring among Alaskan Pacific herring populations identified using microsatellite variation: Journal of Fish Biology, v. 53, no. 1, p. 150–163.

O'Dea, N.R., and Haedrich, R.L., 2001, COSEWIC status report on the northern wolffish *Anarhichas denticulatus* in Canada, *in* Committee on the Status of Endangered Wildlife in Canada, Status report on the northern wolffish Anarhichas denticulatus in Canada: Ottawa, Committee on the Status of Endangered Wildlife in Canada, p. 1–21.

O'Grady, S.M., and DeVries, A.L., 1982, Osmotic and ionic regulation in polar fishes: Journal of Experimental Marine Biology and Ecology, v. 57, no. 2–3, p. 219–228.

Olden, J.D., Lawler, J.J., and Poff, N.L., 2008, Machine learning methods without tears—A primer for ecologists: The Quarterly Review of Biology, v. 83, no. 2, p. 171–193. Orcutt, H.G., 1950, The life history of the starry flounder *Platichthys stellatus* (Pallas): Fish Bulletin, California Fish and Game, v. 78, 64 p.

Orlov, A.M., 1999, Some aspects of trophic relations among Pacific predatory fishes off the northern Kuril Islands and southeastern Kamchatka, *in* MacKinlay, D., and Houlihan, D., eds., Gutshop '98: American Fisheries Society, p. 41–52.

Orlov, A.M., 2003, Impact of eddies on spatial distributions of groundfishes along waters off the northern Kuril Islands, and southeastern Kamchatka (north Pacific Ocean): Indian Journal of Marine Sciences, v. 32, no. 2, p. 95–111.

Orlov, A.M., 2005, Bottom trawl-caught fishes and some features of their vertical distribution in the Pacific waters off the north Kuril Islands and south-east Kamchatka, 1993– 1999: Aqua, Journal of Ichthyology and Aquatic Biology of Fishes, v. 9, no. 4, p. 139–160.

Orr, A.J., Banks, A.S., Mellman, S., Huber, H.R., and DeLong, R.L., 2004, Examination of the foraging habits of Pacific harbor sea (*Phoca vitulina richardsoni*) to describe their use of the Umpqua River, Oregon, and their predation on salmonids: Fishery Bulletin, v. 102, no. 1, p. 108–117.

Orr, J.W., and Matarese, A.C., 2000, Revision of the genus *Lepidopsetta* Gill, 1862 (Teleostei:Pleuronectidae) based on larval and adult morphology, with a description of a new species from the North Pacific Ocean and Bering Sea: Fishery Bulletin, v. 98, p. 539–582.

Orr, J.W., Wildes, S., Kai, Y., Raring, N., Nakabo, T., Katugin, O., and Guyon, J., 2015, Systematics of North Pacific sand lances of the genus Ammodytes based on molecular and morphological evidence, with the description of a new species from Japan: Fishery Bulletin no. 113, p. 129–156.

Orsi, J.A., and Landingham, J.H., 1985, Numbers, species, and maturity stages of fish captured with beach seines during spring 1981 and 1982 in some nearshore marine waters of southeastern Alaska: National Oceanic and Atmospheric Administration, Technical Memorandum NMFS F/NWC-86, 68 p.

Overland, J.E., and Wang, M., 2013, When will the summer Arctic be nearly sea ice free?: Geophysical Research Letters, v. 40, p. 2,097–2,101.

Pacunski, R.E., Livingston, P.A., and Miller, B.S., 1998, Food of flathead sole *Hippoglossoides elassodon* in the eastern Bering Sea: National Oceanic and Atmospheric Administration, Technical Memorandum NMFS-AFSC-90, 36 p.

Paemelaere, E., and Dobson, S.F., 2011, Fast and slow life histories of carnivores: Canadian Journal of Zoology, v. 89, no. 8, p. 692–704. Page, L.M., Espinosa-Pérez, H., Findley, L.T., Gilbert, C.R., Lea, R.N., Mandrak, N.E., Mayden, R.L., and Nelson, J.S., 2013, Common and scientific names of fishes from the United States, Canada, and Mexico: Bethesda, Md., American Fisheries Society Special Publication 34, 384 p.

Pahlke, K.A., 1985, Preliminary studies of capelin (*Mallotus villosus*) in Alaskan waters: Alaska Department of Fish and Game, Informational Leaflet No. 250, 64 p.

Paine, R.T., 1966, Food web complexity and species diversity: The American Naturalist, v. 100, no. 910, p. 65–75.

Palmer, D.E., and Dugan, L.J., 1990, Fish population characteristics of Arctic National Wildlife Refuge coastal waters, summer 1989: Fairbanks, Alaska, U.S. Fish and Wildlife Service, Progress Report, 83 p.

Palsson, W.A., 1990, Pacific cod (*Gadus macrocephalus*) in Puget Sound and adjacent waters—Biology and stock assessment: Olympia, State of Washington, Department of Fisheries, Technical Report no. 112, 137 p.

Panchenko, V.V., 2001, Reproduction peculiarities of plain sculpin *Myoxocephalus jaok* in Peter the Great Bay, Sea of Japan: Russian Journal of Marine Biology, v. 27, p. 111–112.

Panchenko, V.V., 2002, Age and growth of sculpins of the genus *Myoxocephalus* (Cottidae) in Peter the Great Bay (the Sea of Japan): Journal of Ichthyology, v. 42, no. 7, p. 516–522.

Panchenko, V.V., 2005, Seasonal distribution of antlered sculpin, *Enophrys diceraus* (Cottidae) in Peter the Great Bay, Sea of Japan: Russian Journal of Marine Biology, v. 31, p. 273–278.

Parker-Stetter, S.L., Horne, J.K., and Weingartner, T.J., 2011, Distribution of polar cod and age-0 fish in the U.S. Beaufort Sea: Polar Biology v. 34, p. 1,543–1,557.

Parin, N.V., 2004, Ichtyofauna of the seas of Russia—
Biological diversity and fishery potential: Izvestiya TINRO, v. 137, p. 226–231.

Parin, N.V., Evseenko, S.A., and Vasil'eva, E.D., 2014, Ryby morei Rossii—Annotirovannyi catalog (Fishes of Russian seas—Annotated catalogue): Moscow: KMK Scientific Press, 733 p. [In Russian.]

Parrish, J.D., Braun, D.P., and Unnach, R.S., 2003, Are we conserving what we say we are?—Measuring ecological integrity within protected areas: BioScience, v. 53, no. 9, p. 851–860.

Patton, J.C., Gallaway, B.J., Fechhelm, R.G., and Cronin, M.A., 1997, Genetic variation of microsatellite and mitochondrial DNA markers in Broad Whitefish (*Coregonus nasus*) in the Colville and Sagavanirktok Rivers in northern Alaska: Canadian Journal of Fisheries and Aquatic Sciences, v. 54 no. 7, p. 1,548–1,556.

Pauly, D., and Swartz, W., 2007, Marine fish catches in North Siberia (Russia, FAO Area 18), *in* Zeller, D., and Pauly, D., eds., Reconstruction of marine fisheries catches for key countries and regions (1950–2005): Fisheries Centre, University of British Columbia, Canada, Fisheries Centre Research Reports, v. 15, no. 2, p. 17–33.

Pauly, D., and Watson, R., 2005, Background and interpretation of the 'Marine Trophic Index' as a measure of biodiversity: Philosophical Translations, The Royal Society B, v. 360, no. 1454, p. 415–423.

Pauley, G.B., Bowers, K.L., and Thomas, G.L., 1988, Species profiles—Life histories and environmental requirements of coast fishes and invertebrates (Pacific Northwest)—Chum salmon: U.S. Fish and Wildlife Service and U.S. Army Corps of Engineers, Biological Report no. 82 (11.81), TR EL-82-4, 17 p.

Percy, R., 1975, Fishes of the outer Mackenzie Delta: Victoria, British Columbia, Beaufort Sea Project, Beaufort Sea Technical Report, no. 8, 114 p.

Pertseva-Ostroumova, T.A., 1961, The reproduction and development of far-eastern flounders: Moscow, Academy Nauk U.S.S.R. Institute Okeanol, 486 p. [In Russian.]

Petersen, M.F., and Steffensen, J.F., 2003, Preferred temperature of juvenile Atlantic cod *Gadus morhua* with different haemoglobin genotypes at normoxia and moderate hypoxia: Journal of Experimental Biology, v. 206, p. 359–364.

Philo, L.M., George, J.C., and Moulton, L.L., 1993b, The occurrence and description of anadromous fish in the Dease Inlet/Admiralty Bay, Alaska area, 1988–1990: Barrow, Alaska, Department of Wildlife Management, North Slope Borough and MJM Research, 150 p.

Philo, L.M., George, J.C., and Moulton, L.L., 1993a, The occurrence and description of anadromous and freshwater fish in Teshekpuk Lake, Alaska, 1990–1992: Barrow, Alaska, Department of Wildlife Management, North Slope Borough and MJM Research, 98 p.

Piatt, J.F., and Springer, A.M., 2003, Advection, pelagic food webs and the biogeography of seabirds in Beringia: Marine Ornithology, v. 31, p. 141–154. Piatt, J.F., Wells, J.L., MacCharles, A., and Fadely, B.S., 1989, The distribution of seabirds and fish in relation to ocean currents in the southeastern Chukchi Sea, *in* Montevecchi, W.A., and Gaston, A.J., eds., Studies of high-latitude seabirds 1—Behavioral, energetic, and oceanographic aspects of seabird feeding ecology: Canadian Wildlife Service, Occasional Papers, no. 68, p. 21–31.

Pietsch, T.W., 1993, Systematics and distribution of cottid fishes of the genus *Triglops* Reinhardt (Teleostei: Scorpaeniformes): Zoological Journal of the Linnean Society, v. 109, p. 335–393.

Pirtle, J.L., and Mueter, F.J., 2011, Beaufort Sea fish and their trophic linkages—Literature search and synthesis: Anchorage, Alaska, U.S. Department of the Interior, Bureau of Ocean Management, Regulation, and Enforcement, BOEMRE 2011-021, 41 p.

Pockett, S., and Macdonald, J.A., 1986, Temperature dependence of neurotransmitter release in the Antarctic fish *Pagothenia borchgrevinki*: Experientia, v. 42, no. 4, p. 414–415.

Policansky, D., and Sieswerda, P., 1979, Early life history of the starry flounder, *Platichthys stellatus*, reared through metamorphosis in the laboratory: Transactions of the American Fisheries Society, v. 108, no. 3, p. 326–327.

Polyak, L., Alley, R.B., Andrews, J.T., Brigham-Grette, J., Cronin, T.M., Darby, D.A. and others, 2010, History of sea ice in the Arctic: Quaternary Science Reviews v. 29, p. 1,757–1,778.

Ponomarenko, V.P., 1968, Some data on the distribution and migrations of polar cod in the seas of the Soviet Arctic: Rapports et Procès-Verbeaux des Réunions du Conseil International pour l'Exploration de la Mer, v. 158, p. 131–135.

Pörtner, H.O., 2002, Physiological basis of temperaturedependent biogeography—Trade-offs in muscle design and performance in polar ectotherms: Journal of Experimental Biology, v. 205, p. 2,217–2,230.

Pörtner, H.O., Berdal, B., Blust, R., Brix, O., Colosimo, A., De Wachter, B., Giuliani, A., Johansen, T., Fischer, T., Knust, R., Lannig, G., Naevdal, G., Nedenes, A., Nyhammer, G., Sartoris, F.J., Serendero, I., Sirabella, P., Thorkildsen, S., and Zakhartsev, M., 2001, Climate induced temperature effects on growth performance, fecundity of recruitment in marine fish—Developing a hypothesis for cause and effect relationships in Atlantic cod (*Gadus morhua*) and common eelpout (*Zoarces viviparous*): Continental Shelf Research, v. 21, no. 18–19, p. 1,975–1,997. Pörtner, H.O., Bock, C., Rainer, K., Lanning, G., Lucassen, M., Mark, F.C., and Satoris, F.J., 2008, Cod and climate in a latitudinal cline: physiological analyses of climate effects in marine fishes: Climate Research, v. 37, p. 253–270.

Pörtner, H.O., Lucassen, M., and Storch, D., 2005, Metabolic biochemistry—Its role in thermal tolerance and in the capacities of physiological and ecological function, *in* Farrell, A.P., and Steffensen, J.F., eds., The physiology of polar fishes: Amsterdam, Elsevier, p. 79–154.

Power, G., 1997, A review of fish ecology in Arctic North America: American Fisheries Society Symposium, no. 19, p. 13–39.

Praebel, K., Hunt, B., Hunt, L., DeVries, A.L., 2009, The presence of quantification of splenic ice in the McMurdo Sound notothenioid fish, *Pagothenia borchgrevinki* (Boulenger, 1902)—Comparative biochemistry and physiology part A: Molecular and Integrative Physiology, v. 154, no. 4, p. 564–569.

Praebel, K., and Ramlov, H., 2005, Antifreeze activity in the gastrointestinal fluids of *Arctogadus glacialis* (Peters, 1874) is dependent on food type: Journal of Experimental Biology, v. 208, p. 2,609–2,613.

Protection of the Arctic Marine Environment, 2013, Large Marine Ecosystems (LMEs) of the Arctic area—Revision of the LME map 15 May 2013, 2nd ed.: Arctic Council, 24 p.

Pruter, A.T., and Alverson, D.L., 1962, Abundance, distribution, and growth of flounders in the south-eastern Chukchi Sea: Journal de Conseil International pour l'Exploration de la Mer., v. 27, p. 79–99.

Quast, J.C., 1974, Density distribution of juvenile Arctic cod, *Boreogadus saida*, in the eastern Chukchi Sea in the fall of 1970: Fishery Bulletin, v. 72, no. 4, p. 1,094–1,105.

R Core Team, 2013, R—A language and environment for statistical computing: Vienna, Austria, R Foundation for Statistical Computing, accessed November 5, 2013, at http://www.R-project.org/.

Rand, K.M., and Logerwell, E.A., 2011, The first demersal trawl survey of benthic fish and invertebrates in the Beaufort Sea since the late 1970s: Polar Biology, v. 34, no. 4, p. 475–488.

Rasmussen, C., Purcell, M.K., Gregg, J.L., LaPatra, S.E., Winton, J.R., and Hershberger, P.K., 2010, Sequence analysis of the internal transcribed spacer (ITS) region reveals a novel clade of *Ichthyophonus sp.* from rainbow trout: Diseases of Aquatic Organisms, v. 89, p. 179–183.

Rass, T.S., 1941, Analogous or parallel variations in structure and development of fishes in the northern and Arctic seas: Jubilee Publication of the Moscow Society of Naturalists 1805–1940, p. 1–60. Rass, T.S., 1989, Regularity in the increase of egg mass size among marine fishes and invertebrates polewards from the Equator to high latitudes: Rapports et Procès-Verbeaux des Réunions du Conseil International pour l'Exploration de la Mer, v. 191, p. 443–444.

Ratynski, R.A., 1983, Mid-summer ichthyoplankton populations of Tuktoyaktuk Harbor, N.W.T.: Canadian Technical Report of Fisheries and Aquatic Sciences, no. 1218, 21 p.

Raymond, J.A., 1992, Glycerol is a colligative antifreeze in some northern fishes: Journal of Experimental Zoology, v. 262, no. 3, p. 347–352.

Raymond, J.A., 1994, Seasonal variations of trimethylamine oxide and urea in the blood of a cold-adapted marine teleost, the rainbow smelt: Fish Physiology and Biochemistry, v. 13, no. 1, p. 13–22.

Raymond, J.A., 1997, Responses of marine fishes to freezing temperatures—A new look at colligative mechanisms, *in* Willis, J.S., ed., Advances in molecular and cell biology: Greenwich, Conn., JAI Press, p. 33–142.

Raymond, J.A., and DeVries, A.L., 1977, Adsorption inhibition as a mechanism of freezing resistance in polar fishes: Proceedings of the National Academy of Sciences of the USA, v. 74, no. 6, p. 2,589–2,593.

Raymond, J.A., and DeVries, A.L., 1998, Elevated concentrations and synthetic pathways of trimethylamine oxide and urea in some teleosts fishes of McMurdo Sound, Antarctica: Fish Physiology and Biochemistry, v. 18, no. 4, p. 387–398.

Raymond, J.A., Lin, Y., and DeVries, A.L., 1975, Glycoprotein and protein antifreezes in two Alaskan fishes: Journal of Experimental Zoology, v. 193, no. 1, p. 125–130.

Reist, J.D., and Chang-Kue, K.T.J., 1997, The life history and habitat usage of broad whitefish in the lower Mackenzie River Basin, *in* Tallman, R.F., and Reist, J.D., eds., The proceedings of the broad whitefish workshop—The biology, traditional knowledge and scientific management of broad whitefish (*Coregonus nasus* (Pallas)) in the Lower Mackenzie River: Canadian Technical Report of Fisheries and Aquatic Sciences, no. 2193, p. 63–84.

Reist, J.D., Wrona, F.J., Prowse, T.D., Power, M., Dempson, J.B., Beamish, R.J., King, J.R., Carmichael, T.J., and Sawatzky, C.D., 2006, General effects of climate change on Arctic fishes and fish populations, AMBIO: A Journal of the Human Environment v. 35, no. 7, p. 370–380.

Renaud, P.E., Carroll, M.L., and Ambrose, W.G., Jr., 2008,
Effects of global warming on Arctic sea-floor communities and its consequences for higher trophic levels, *in* Duarte,
C.M., ed., Impacts of global warming on polar ecosystems:
Bilbao, Spain, Fundación BBVA, p. 141–177.

Reynolds, W.W., 1977, Temperature as a proximate factor in orientation behavior: Journal of the Fisheries Research Board of Canada, v. 34, no. 5, p. 734–739.

Richardson, E.S., Reist, J.D., and Minns, C.K., 2001, Life history characteristics of freshwater fishes occurring in the Northwest Territories and Nunavut, with major emphasis on lake habitat requirements: Canadian Manuscript Report of Fisheries and Aquatic Sciences no. 2569, 156 p.

Ricker, W.E., 1975, Computation and interpretation of biological statistics of fish populations: Ottawa, Canada, Bulletin of Fisheries Research Board of Canada, no. 191, 382 p.

Robards, M.D., and Piatt, J.F., 1999, Biology of the genus *Ammodytes*, the sand lances, *in* Robards, M.D., and others, eds., Sand land—A review of biology and predator relations and annotated bibliography: U.S. Forest Service, Pacific Northwest Research Station, Research Paper PNW-RP-521.

Robards, M.D., Piatt, J.F., and Rose, G.A., 1999, Maturation, fecundity, and intertidal spawning of Pacific sand lance in the northern Gulf of Alaska: Journal of Fish Biology v. 54, no. 5, p. 1,050–1,068.

Robards, M.D., Rose, G.A., and Piatt, J.F., 2002, Growth and abundance of Pacific sand lance, *Ammodytes hexapterus*, under differing oceanographic regimes: Environmental Biology of Fishes, v. 64, no. 4, p. 429–441.

Robinson, L.M., Elith, J., Hobday, A.J., Pearson, R.G., Kendall, B.E., Possingham, H.P., and Richardson, A.J., 2011, Pushing the limits in marine species distribution modeling—Lessons from the land present challenges and opportunities: Global Ecology and Biogeography, v. 20, p. 789–802.

Robles, V., Cabrita, E., Fletcher, G.L., Shears, M.A., King, M.J., and Herráez, M.P., 2005, Vitrification assays with embryos from a cold tolerant sub-arctic fish species: Theriogenology v. 64, no. 7, p. 1,633–1,646.

Roessig, J.M., Woodley, C.M., Cech, J.J., Jr., and Hansen, L.J., 2004, Effects of global climate change on marine and estuarine fishes and fisheries: Reviews in Fish Biology and Fisheries, v. 14, no. 2, p. 251–275. Rogers, D.E., Rabin, D.J., Rogers, B.J., Garrison, K.J., and Wangerin, M.E., 1979, Seasonal composition and food web relationships of marine organisms in the nearshore zone of Kodiak Island—including ichthyoplankton, meroplankton (shellfish), zooplankton, and fish: Seattle, University of Washington College of Fisheries, Fisheries Research Institute, Technical Report FRI-UW-7906, 123 p.

Romanuk, T.N., Hayward, April, and Hutchings, J.A., 2011, Trophic level scales positively with body size in fishes: Global Ecology and Biogeography, v. 20, p. 231–240, doi: 10.1111/j.1466-8238.2010.00579.x.

Ronholt, L.L., Wakabayashi, K., Wilderbuer, T.K., Yamaguchi, H., and Okada, K., 1986, Ground fish resource of the Aleutian Island waters based on the U.S.–Japan trawl survey, June–November 1980: Bulletin International North Pacific Fisheries Commission 48, p. 251.

Rooper, C.N., 2002, English sole transport during pelagic stages on the Pacific Northwest coast, and habitat use by juvenile flatfish in Oregon and Washington estuaries: Seattle, University of Washington, Ph.D. dissertation.

Rose, G.A., 2005a, Capelin (*Mallotus villosus*) distribution and climate: a sea "canary" for marine ecosystem change: ICES Journal of Marine Science, v. 62, no. 7, p. 1,524–1,530.

Rose, G.A., 2005b, On distributional responses of North Atlantic fish to climate change: ICES Journal of Marine Science, v. 62, no. 7, p. 1,360–1,374.

Rosenberg, A., Bigford, T.E., Leathery, S., Hill, R., and Bickers, K., 2000, Ecosystem approaches to fishery management through essential fish habitat: Bulletin of Marine Science, v. 66, no. 3, p. 535–542.

Rosenthal, R.J., 1980, Shallow water fish assemblages in the northeastern Gulf of Alaska—Habitat evaluation, species composition, abundance, spatial distribution and trophic interaction, *in* Bureau of Land Management, Environmental assessment of the Alaskan Continental Shelf, final reports of principal investigators: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, p. 451–540.

Rosenthal, R.J., Haldorson, L.J., Field, L.J., Moran-O'Connell, V., LaRiviere, M.G., Underwood, J., and others, 1982, Inshore and shallow offshore bottomfish resources in the southeastern Gulf of Alaska (1981–82): Alaska Coastal Research and University of Alaska, Juneau, 166 p.

Roy, K., Jablonski, D., Vaelntine, J.W., and Rosenberg,
G., 1998, Marine latitudinal gradients—Tests of causal hypotheses: Proceedings of the National Academy of Science, v. 95, no. 7, p. 3,699–3,702.

Rubinsky, B., Arav, A., Mattioli, M., and DeVries, A.L., 1990, The effect of antifreeze glycopeptides on membrane potential changes at hypothermic temperatures: Biochemical and Biophysical Research Communications, v. 173, no. 3, p. 1,369–1,374.

Russian-American Long-Term Census of the Arctic, 2009, Expedition to the Bearing Strait and Pacific side of the Arctic Ocean: National Oceanic and Atmospheric Administration Web site, accessed April 15, 2013, at http:// www.arctic.noaa.gov/rusalca/2009.

Sadorus, L.L., Lauth, R., and Ranta, A.M., 2012, Size and age composition of Pacific halibut in NMFS Bering Sea shelf trawl surveys, *in* IPHC Report of Assessment and Research Activities 2012: International Pacific Halibut Commission online report, p. 593–600, accessed February 16, 2014, at http://www.iphc.int/publications/rara/2012/rara2012593_ beringtrawlcomp.pdf.

Safronov, S.N., 1986, Peculiarities of reproduction and principles of change in the fecundity of Pacific navaga, *Eleginus gracilis* (Gadidae): Journal of Ichthyology, v. 26, no. 5, p. 59–68.

Salveson, S.J., 1976, Alaska plaice (family Pleuronectidae), *in* Pereya, W.T., Reeves, J.E., and Bakkala, R.G., eds., Demersal fish and shellfish resources of the Eastern Bering Sea in the baseline year 1975: National Oceanic and Atmospheric Administration, p. 489–495.

Salveson, S.J., and Alton, M.S., 1976, Yellowfin sole (family Pleuronectidae) *in* Pereya, W.T., Reeves, J.E., and Bakkala, R.G., eds.: Demersal fish and shellfish resources of the Eastern Bering Sea in the baseline year 1975: Seattle, Washington, National Marine Fisheries Service, Northwest and Alaska Fisheries Center, Processed Report, p. 439–459.

Sameoto, D.D., 1988, Feeding of lantern fish Benthosema glaciale off the Nova Scotia shelf: Marine Ecology Progress Series, v. 44, p. 113–129.

Sandercock, F.K., 1991, Life history of Coho salmon (*Oncorhynchus kisutch*), *in* Groot, C., and Margolis, L., eds., Pacific salmon life histories: Vancouver, British Columbia, UBC Press, p. 397–445.

Saunders, M.W., and McFarlane, G.A., 1993, Age and length of maturity of the female spiny dogfish, *Squalus acanthias*, in the Strait of Georgia, British Columbia, Canada: Environmental Biology of Fishes, v. 38, p. 49–57.

Schmidt, D.R., Griffiths, W.B., Beaubien, D.K., and Herlugson, C.J., 1991, Movement of young-of-the-year Arctic ciscoes across the Beaufort Sea coast, 1985–1988: American Fisheries Society Symposium, no. 11, p. 132–144. Schmidt, D.R., Griffiths, W.B., and Martin, L.R., 1987, Importance of anadromous fish overwintering habitat in the Sagavanirktok River Delta, Alaska: Anchorage, Alaska, Report by Ecological Research Associates for Standard Alaska Production Company and North Slope Borough, 71 p.

Schmidt, D.R., Griffiths, W.R., and Martin, L.R., 1989, Overwintering biology of anadromous fish in the Sagavanirktok River Delta, Alaska: Biological Papers of the University of Alaska, v. 24, p. 55–74.

Scholander, P.F., Flagg, W., Walters, V., and Irving, L., 1953, Climatic adaptations in Arctic and tropical poikilotherms: Physiological Zoology, v. 26, no. 1, p. 67–92.

Scott, G.K., Fletcher, G.L., and Davies, P.L., 1986, Fish antifreeze proteins—Recent gene evolution: Canadian Journal of Fisheries and Aquatic Science, v. 43, no. 5, p. 1,028–1,034.

Scott, G.K., Hayes, P.H., Fletcher, G.L., and Davies, P.L., 1988, Wolffish antifreeze protein genes are primarily organized as tandem repeats that each contain two genes in inverted orientation: Molecular and Cellular Biology, v. 8, no. 9, p. 3,670–3,675.

Scott, W.B., and Scott, M.G., 1988, Atlantic fishes of Canada: Toronto, University of Toronto Press, 730 p.

Seitz, A.C., Loher, T., and Nielsen, J.L., 2008, Seasonal movements and environmental conditions experienced by Pacific halibut along the Aleutian Islands, examined by pop-up satellite tags: International Pacific Halibut Commission Scientific Report, no. 85, 24 p.

Seitz, A.C., Norcross, B.L., and Nielsen, J.L., 2003, Pip-up archival transmitting (PAT) tags—A method to investigate the migration and behavior of Pacific halibut *Hippoglossus stenolepsis* in the Gulf of Alaska: Alaska Fishery Research Bulletin, v. 10, no. 2, p. 124–135.

Seitz, A.C., Norcross, B.L., Wilson, D., and Nielsen, J.L., 2005, Identifying spawning behavior in Pacific halibut, *Hippoglossus stenolepis*, using electronic tags: Environmental Biology of Fishes, v. 73, p. 445–451.

Sekerak, A.D., Stallard, N., and Griffiths, W.B., 1992, Distribution of fish and fish harvests in the nearshore Beaufort Sea and Mackenzie Delta during ice-covered periods, October–June: Environmental Studies Research Funds Report, LGS Ltd., no. 117, 524 p.

Shchetinnikov, A.S., 1983, Nutrition of *Acantholumpenus mackayi* (Stichaeidae) in the Gulf of Terpenium (Sakhalin Island): Journal of Ichthyology, v. 23, no. 6, p. 155–158.

Shears, M.A., Kao, M.H., Scott, G.K., Davies, P.L., and Fletcher, G.L., 1993, Distribution of type III antifreeze in the Zoarcoidei: Molecular Marine Biology and Biotechnology, v. 2, p. 104–111.

Sheiko, B.A., and Fedorov, V.V., 2000, Class cephalaspidomorphi-lampreys, Class chondrichthyescartilaginous fishes, Class holocephali-chimaeras, Class osteichthyes-bony fishes, *in* Artuykhin, Yu. B, and Sheyko, B.A., eds., Catalog of vertebrates of Kamchatka and adjacent waters: Russia, Rossiyskaya Akademiya Nauk, p. 7–69. [In Russian.]

Sheiko, B.A., and Mecklenburg, C.W., 2004, Family Agonidae Swainson 1839—Poachers: California Academy of Sciences Annotated Checklists of Fishes, no. 30, 27 p.

Shestakov, A.V., and Nazarkin, M.V., 2006, On the biology of the white-spotted greenling *Hexagrammos stelleri* and the masked greenling *H. octogrammus* (Hexagrammidae) from Taui Bay of the Sea of Okhotsk: Journal of Ichthyology, v. 46, no. 8, p. 677–680.

Shuntov, V.P., 1970, Seasonal distribution of black and arrowtoothed halibuts in the Bering Sea: *in* Moiseev, P.A., ed., Soviet fisheries investigations in the northeastern Pacific: Jerusalem, Israel Program for Scientific Translations 1968, p. 397–408.

Sick, K., 1965, Hemoglobin polymorphism of cod in the North Sea and the North Atlantic Ocean: Heriditas, v. 54, no. 1, p. 49–68.

Siefert, D.L., Incze, L.S., and Ortner, P.B., 1988, Vertical distribution of zooplankton, including ichthyoplankton, in Shelikof Strait, Alaska—Data from fisheries-oceanography coordinated investigations (FOCI) cruise in May 1986: National Marine Fisheries Service, NWAFC Processed Report, v. 88, no. 28, 232 p.

Sigler, M.F., Renner, M., Danielson, S.L., Eisner, L.B., Lauth, R.R., Kuletz, K.J., Logerwell, E.A., and Hunt, G.L., Jr., 2011, Fluxes, fins, and feathers—Relationships among the Bering, Chukchi, and Beaufort Seas in a time of climate change: Oceanography, v. 24, no. 3, p. 250–265.

Simenstad, C.A., Isakson, J.S., and Nakatani, R.E., 1977, Marine fish communities, *in* Merritt, M.L., and Fuller, R.G., eds., The environment of Amchitka Island, Alaska: National Technical Information Center, Energy Research and Development Administration TID-26712, p. 451–492.

Sinclair, E.H., and Zeppelin, T.K., 2002, Seasonal and spatial differences in diet in the western stock of Steller sea lions (*Eumetopias jubatus*): Journal of Mammology, v. 83, no. 4, p. 973–990. Slagstad, D., Ellingsen, I.H., and Wassmann, P., 2011, Evaluating primary and secondary production in an Arctic Ocean void of summer sea ice—An experimental simulation approach: Progressive Oceanography, v. 90, no. 1–4, p. 117–131.

Smetacek V., and Nicol, S., 2005, Polar ecosystems in a changing world: Nature, v. 437, p. 362–368.

Smith, K.R., Somerton, D.A., Yang, M.-S., and Nichol, D.G., 2004, Distribution and biology of prowfish (*Zaprora silenus*) in the northeast Pacific: Fishery Bulletin, v. 102, no. 1, p. 168–178.

Smith, M.A., 2010, Arctic marine synthesis—Atlas of the Chukchi and Beaufort Seas: Anchorage, Alaska, Audubon Alaska and Oceana.

Smith, R.E.H., Anning, J., Clement, P., and Cota, G., 1988, Abundance and production of ice algae in Resolute Passage, Canadian Arctic: Marine Ecology Progress Series, v. 48, p. 251–263.

Smith, R.L., Vallarino, M., Barbour, E., Fitzpatrick, E., and Barber, W.E., 1997, Population biology of the Bering flounder in the Northeastern Chukchi Sea: American Fisheries Society Symposium, v. 19, p. 127–132.

Smith, R.L., Barber, W.E., Vallarino, M., Gillispie, J., and Ritchie, A., 1997a, Population biology of the Arctic staghorn sculpin in the northeastern Chukchi Sea: American Fisheries Society Symposium 19, p. 133–139.

Smith, R.L., Vallarino, M., Barbour, E., Fitzpatrick, E., and Barber, W.E., 1997b, Population biology of the Bering flounder in the northeastern Chukchi Sea: American Fisheries Society Symposium, v. 19, p. 127–132.

Smith, R.T., 1936, Report on the Puget Sound otter trawl investigations: Washington Department of Fisheries, Biological Report No. 36B, 120 p.

Smith, S.L., and Schnack-Schiel, S.B., 1990, Polar zooplankton polar oceanography part B—Chemistry biology and geology: San Diego, Academic Press, p. 527–598.

Sohn, D., Ciannelli, L., and Duffy-Anderson, J.T., 2010, Distribution and drift pathways of Greenland halibut (*Reinhardtius hippoglossoides*) during early life stages in the eastern Bering Sea and Aleutian Islands: Fisheries Oceanography, v. 19, no. 5, p. 339–353.

Sokolovskaya, T.G., Sokolovskii, A.S., and Sobolevskii, E.I., 1998, A list of fishes of Peter the Great Bay (the Sea of Japan): Journal of Ichthyology, v. 38, no. 1, p. 1–11.

Sokolovskii, A.S., and Sokolovskaya, T.G., 1999, The early ontogenesis of *Enophrys diceraus* (Cottidae) from the waters of Peter the Great Bay (the Sea of Japan): Journal of Ichthyology, v. 39, no. 3, p. 268–272.

Sokolovskii, A., and Sokolovskaya, T., 2009, Distribution of ichthyoplankton in the eastern Peter the Great Bay, Sea of Japan, in June and July of 2007: Russian Journal of Marine Biology, v. 35, no. 3, p. 250–254.

Somero, G.N., 1992, Adapting to water stress—Convergence on common solutions, *in* Somero, G.N., Osmond, C.B., and Bolis, C.L., eds., Water and life, comparative analysis of water relationships at the organismic, cellular and molecular levels: Berlin, Springer-Verlag, p. 3–18.

Sorenson, T.F., and Ramlov, H., 2001, Variations in antifreeze activity and serum inorganic ions in the eelpout Zoarces viviparous—Antifreeze activity in the embryonic state: Comparative Biochemistry and Physiology Part A— Molecular and Integrative Physiology, v. 130, no. 1, p. 123–132.

Spalding, M.D., Fox, H.E., Allen, G.R., Davidson, N., Ferdaña, Z.A., Finlayson, M., Halpern, B.S., Martin, K.D., Mcmanus, E., Molnar, J., Recchia, C.A., and Robertson, J., 2007, Marine ecosystems of the world—A bioregionalization of coastal and shelf areas: BioScience, v. 57, no. 7, p. 573–583.

Speer, L., and Laughlin, T.L., 2011, IUNC/NRDC workshop to identify areas of ecological and biological significance or vulnerability in the Arctic marine environment: IUNC/ NRDC Workshop Report, La Jolla, Calif., November 2–4, 2010, 16 p.

Springer, A.M., Murphy, E.C., Rosenau, D.G., Peter McRoy, C., and Cooper, B.A., 1986, The paradox of pelagic food webs in the northern Bering Sea-I—Seabird food habits: Continental Shelf Research, v. 7, no. 8, p. 895–911.

St.-Pierre, G., 1984, Spawning locations and season for Pacific halibut: Seattle, Washington, International Pacific Halibut Commission Scientific Report no. 70, 46 p.

Stabeno, P.J., Kachel, N.B., Moore, S.E., Napp, J.M., Sigler, M., Yamaguchi, A., and Zerbini, A.N., 2012b, Comparison of warm and cold years on the southeastern Bering Sea shelf and some implications for the ecosystem, Deep-Sea Research II: Tropical Studies in Oceanography. v. 65–70, p. 31–45.

Stabeno, P., and others, 2012a, A comparison of the physics of the northern and southern shelves of the eastern Bering Sea and some implications for the ecosystem, Deep Sea Research Part II: Topical Studies in Oceanography, v. 65–70, p. 14–30.

Stark, J.W., 2007, Geographic and seasonal variations in maturation and growth of female Pacific cod (*Gadus macrocephalus*) in the Gulf of Alaska and Bering Sea: Fishery Bulletin, v. 105, no. 3, p. 396–407.

Stauffer, G., comp., 2004, NOAA protocols for groundfish bottom trawl surveys of the Nation's fishery resources: U.S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Memorandum NMFS-F/SPO-65, 205 p.

Stearns, S.C., 1977, The evolution of life history traits—A critique of the theory and a review of the data: Annual Review of Ecology and Systematics, v. 8, p. 145–171.

Steffensen, J.F., 2002, Metabolic cold adaptation of polar fish based on measurements of aerobic oxygen consumption— Fact or artefact? Artefact!: Comparative Biochemistry and Physiology Part A—Molecular and Integrative Physiology, v. 132, no. 4, p. 789–795.

Steffensen, J.F., 2005, Respiratory systems and metabolic rates, *in* Farrell, A.P., and Steffensen, J.F., eds., The physiology of polar fishes: Amsterdam, Elsevier, p. 203–238.

Steffensen, J.F., Bushnell, P.G., and Schurmann, H., 1994, Oxygen consumption in four species of teleosts from Greenland—No evidence of metabolic cold adaptation: Polar Biology, v. 14, no. 1, p. 49–54.

Stehmann, M., and Bürkel, D.L., 1984, Rajidae, *in* Whitehead, P.J.P., and others, eds., Fishes of the North-eastern Atlantic and the Mediterranean: Paris, Unesco, p. 163–196.

Stephenson, S.A., 2006, A review of the occurrence of Pacific Salmon (*Oncorhynchus* spp.) in the Canadian Western Arctic: Arctic, v. 59, no. 1, p. 37–46.

Stewart, D.B., Ratynski, R.A., Bernier, L.M.J., and Ramsey, D.J., 1993, A fishery development strategy for the Canadian Beaufort Sea-Amundsen Gulf area: Canadian Technical Report Fisheries and Aquatic Sciences 1910, 135 p.

Stewart, H., 1977, Indian fishing—Early methods on the northwest coast: Seattle, University of Washington Press, 182 p.

Stockwell, D., Horner, R., Saitoh, S., and Hill, V., 2008, Phytoplankton, *in* Hopcroft, R., Bluhm, B., and Gradinger, R., eds., Arctic Ocean synthesis—Analysis of climate change impacts in the Chukchi and Beaufort Seas with strategies for future research: Institute of Marine Sciences, University of Alaska Fairbanks, p. 33–37.

Stoner, A.W., and Abookire, A.A., 2002, Sediment preferences and size-specific distribution of young-of-the-year Pacific halibut in an Alaska nursery: Journal of Fish Biology, v. 61, no. 3, p. 540–559.

Stout, H.A., Gastafson, R.G., Lenarz, W.H., McCain, B.B., VanDoornik, D.M., Builder, T.L., and Methot, R.D., 2001, Status review of Pacific herring (*Clupea pallasi*) in Puget Sound, Washington: National Oceanic and Atmospheric Administration Technical Memorandum NMFS-NWFSC-45, 175 p.

Suryan, R.M., Sab, V.S., Wallace, B.P., Hatch, S.A., Frederiksen, M., and Wanless, S., 2009, Environmental forcing on life history strategies—Evidence for multitrophic level responses at ocean basin scales: Progress in Oceanography, v. 81, no. 1–4, p. 214–222.

Swanson, J.A., and Hoppe, A.D., 2004, The coordination of signaling during Fc receptor-mediated phagocytosis: Journal of Leukocyte Biology, v. 76, no. 6, p. 1,093–1,103.

Tack, S.L., 1970, The summer distribution and standing stock of the fishes of Izembek Lagoon, Alaska: Fairbanks, University of Alaska, Master's thesis.

Tallman, R.F., Abrahams, M.V., and Chudobiak, D.H., 2002, Migration and life history alternatives in a high latitude species, the broad whitefish, *Coregonus nasus* Pallas: Ecology of Freshwater Fish, v. 11, no. 2, p. 101–111.

Tarasyuk, S.N., 2002, Survival of yellowfin sole (*Limanda aspera Pallas*) in the northern part of the Tatar Strait (Sea of Japan) during the second half of the 20th century, in Batchelder, H.P., and others, eds., PICES-Globe International Program on Climate Change and Carrying Capacity—Report of 2110 BASS/MODEL, MONITOR and REX workshops, and The 2002 MODEL/REX workshop: Sidney, British Columbia, North Pacific Marine Science Organization (PICES).

Thompson, G., Ianelli, J., Dorn, M., Nichol, D., Gaichas, S., and Aydin, K., 2007, Assessment of the Pacific Cod stock in the eastern Bering Sea and Aleutian Islands Area, chap. 2 of Plan team for groundfish fisheries of the Bering Sea/ Aleutian Islands (comps.): Anchorage, Alaska, North Pacific Fishery Management Council, Stock assessment and fishery evaluation report for groundfish resources of the Bering Sea/Aleutian Islands regions, p. 209–328.

Thorsteinson, F.V., Sample, T., and Thorsteinson, L.K., 1984, Fishery resources, chap. 6 *of* Jarvela, L.E., ed., The Navarin Basin environment and possible consequences of planned offshore oil and gas development: Juneau, Alaska, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, OCSEAP, 157 p.

Thorsteinson, L.K., 1996, National Biological Service OCS environmental studies program—research and monitoring in Arctic Alaska, *in* Newbury, T., ed., Proceedings of the 1995 Arctic Synthesis Meeting, October 23–25, 1995, Anchorage, Alaska: U.S. Department of the Interior, Minerals Management Service, Alaska OCS Region, OCS Study MMS 95-0065, 206 p. plus appendixes. Thorsteinson, L.K., Becker, P.R., and Hale, D.A., 1989, The Yukon Delta synthesis of information: U.S. Department of Commerce, National Oceanic and Atmospheric Administration, OCS Study MMS 89-0081, 89 p.

Thorsteinson, L.K., Jarvela, L.E., and Hale, D.A., 1990, Arctic fish habitat use investigations—Nearshore studies in the Alaskan Beaufort Sea, summer 1988: Final Report, Alaska Office, Ocean Assessments Division, National Oceanic and Atmospheric Administration, Research Unit 682, 125 p.

Thorsteinson, L.K., Jarvela, L.E., and Hale, D.A., 1991, Arctic fish habitat use investigations—Nearshore studies in the Alaskan Beaufort Sea, summer 1990: Anchorage, Alaska, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, OCSEAP Final Report to U.S. Department of the Interior, Mineral Management Service, 166 p.

Tokranov, A.M., 1981, Distribution of sculpins (Pisces, Cottidae) on the west Kamchatka shelf in summer: Zoologicheskii Zhurnal, v. 60, no. 2, p. 229–237.

Tokranov, A.M., 1987a, Feeding of giant sculpin Myoxocephalus polyacanthocephalus Pallas and plain sculpin M. jaok (Cuvier) (Cottidae) in coastal waters of Kamchatka: Journal of Ichthyology, v. 27, p. 104–114.

Tokranov, A.M., 1987b, Species composition and distribution of sea poachers (Pisces, Agonidae) in nearshore waters of Kamchatka: Zoologicheskii Zhurnal, v. 66, p. 385–392.

Tokranov, A.M., 1991, Diet of the Pacific cod, *Gadus morhua macrocephalus*, and its position in the food chain in Kamchatkan coastal waters: Journal of Ichthyology, v. 31, no. 5, p. 84–98.

Tokranov, A.M., 1995, The size and sex structure of sculpins of the genus Triglops (Cottidae) in the coastal waters of Kamchatka: Journal of Ichthyology, v. 35, no. 4, p. 140–144.

Tokranov, A.M., 1999, Some features of biology of the prowfish *Zaprora silenus* (Zaproridae) in the Pacific waters of the northern Kuril Islands and southeastern Kamchatka: Journal of Ichthyology, v. 39, no. 6, p. 475–478.

Tokranov, A.M., 2007, Distribution and some biological features of the Pacific sand lance *Ammodytes hexapterus* (Ammodytidae) in waters off Kamchatka in the Sea of Okhotsk: Journal of Ichthyology, v. 47, no. 4, p. 288–295.

Tokranov, A.M., and Maksimenkov, V.V., 1994, Feeding of the starry flounder, *Platichthys stellatus*, in the Bol'shaya River estuary (Western Kamchatka): Journal of Ichthyology, v. 34, no. 1, p. 76–83. Tokranov, A.M., and Orlov, A.M., 2005, Some features of the biology of *Icelus spatula* (Cottidae) in Pacific waters off the northern Kuril Islands: Journal of Ichthyology, v. 45, no. 3, p. 229–236.

Tokranov, A.M., Orlov, A.M., and Biryukov, I.A., 2004, Distribution and size-weight composition of some rare fish species in Pacific waters off the northern Kurile Islands and southeast Kamchatka: Journal of Ichthyology, v. 44, no. 3, p. 208–216.

Tokranov, A.M., and Polutov, V.I., 1984, Fish distribution in the Kronitsky Bay in relation to abiotic factors: Zoologicheskii Zhurnal, v. 63, p. 1,363–1,373.

Tokranov, A.M., and Zavarina, S.V., 1992, Size-age structure and sex ratios of Alaska plaice, *Pleuronectes quadrituberculatus*, on the Western Kamchatka Shelf: Journal of Ichthyology, v. 32, no. 7, p. 118–127.

Treble, M.A., Campana, S.E., Wastle, R.J., Jones, C.M., and Boje, J., 2008, Growth analysis and age validation of a deepwater Arctic fish, the Greenland halibut (*Reinhardtius hippoglossoides*): Canadian Journal of Fisheries and Aquatic Sciences, v. 65, no. 6, p. 1,047–1,059.

Trevor, H., 2013, GAM—Generalized Additive Models: R package version 1.09, accessed April 15, 2014, at http:// CRAN.R-project.org/package=gam.

Turner, L.M., 1886, Contributions to the natural history of Alaska—Arctic series of publications, no. 2: Washington, D.C., U.S. Government Printing Office, 226 p.

Tyler, A.V., 1972, Food resource division among northern, marine, demersal fishes: Journal of the Fisheries Research Board of Canada, v. 29, no. 7, p. 997–1,003.

Underwood, T.J., Gordon, J.A., Millard, M.J., Thorpe, L.A., and Osborne, B.M., 1995, Characteristics of selected fish populations of Arctic National Wildlife Refuge coastal waters, final report, 1988–91: Fairbanks, Alaska, U.S. Fish and Wildlife Service, Fairbanks Fishery Resource Office, Alaska Fisheries Technical Report, no. 28, 590 p.

United Nations Environment Programme, 2013, What future for the Arctic?: Nairobi, Kenya, Background paper for the Arctic Side Event at the Governing Council/Global Environmental Ministerial Forum, 12 p.

Vadstein, O., Mo, T.A., and Bergh, Ø., 2004, Microbial interactions, prophylaxes and diseases, *in* Moksness, E., Kjørsvik, E., and Olsen, Y., eds., Culture of cold-water marine fish: Victoria, Australia, Blackwell Publishing, p. 28–72.

Valerio, P.F., Kao, M.H., and Fletcher, G.L., 1990, Thermal hysteresis activity in the skin of the cunner, *Tautogolabrus adspersus*: Canadian Journal of Zoology, v. 68, no. 5, p. 1,065–1,067.

Valerio, P.F., Kao, M.H., and Fletcher, G.L., 1992, Fish skin— An effective barrier to ice crystal propagation: Journal of Experimental Biology, v. 164, p. 135–151.

VanGerwen-Toyne, M., Walker-Larsen, J., and Tallman, R.F., 2008, Monitoring spawning populations of migratory coregonids in the Peel River, NT—The Peel River fish study 1998–2002: Canadian Manuscript Report of Fisheries and Aquatic Sciences no. 2851, 56 p.

Van Guelpen, L., 1986, Hookear sculpins (genus Artediellus) of North American Atlantic—Taxonomy, morphological variability, distribution, and aspects of life history: Canadian Journal of Zoology, v. 64, no. 3, p. 677–690.

Van Voorhies, W.V., Raymond, J.A., and DeVries, A.L., 1978, Glycoproteins as biological antifreeze agents in the cod, *Gadus ogac* (Richardson): Physiological Zoology, v. 51, no. 4, p. 347–353.

Verde, C., Carratore, V., Riccio, A., Tamburrini, M., Parisi, E., and di Prisco, G., 2002, The functionally distinct hemoglobins of the Arctic spotted wolffish *Anarhichas minor*: Journal of Biological Chemistry, v. 277, no. 39, p. 36,312–36,320.

Verde, C., de Rosa, M.C., Giordanao, D., Mosca, D., De Pascale, D., Raiola, L., Cocca, E., Carratore, V., Giardina, B., and Di Prisco, G., 2005, Structure, function and molecular adaptations of haemoglobins of the polar cartilaginous fish *Bathyraja eatonii* and *Raja hyperborea*: Biochemical Journal, v. 389, no. 2, p. 297–306.

Verde, C., Parisi, E., and di Prisco, G., 2006, The evolution of thermal adaptation in polar fish: Gene, v. 385, p. 137–145.

Vermeij, G.J., 1991, Marine extinctions and their implications for conservation and biogeography, *in* The Unity of Evolutionary Biology, Proceedings of the Fourth International Congress of Systematic and Evolutionary Biology, v. 1, p. 143–148.

Viavant, T., 2001, Eastern North Slope Dolly Varden spawning and over-wintering assessment feasibility: Anchorage, Alaska, U.S. Fish and Wildlife Service, Office of Subsistence Management, Fisherie Information Services Division, Federal Subsistence Fisheries Resource Monitoring Program, Final Report for Study 00-002, 25 p.

Viavant, T., 2009, Aerial monitoring of Dolly Varden overwintering abundance in the Anaktuvuk, Ivishak, Canning, and Hulahula rivers, 2006-2008, Alaska Department of Fish and Game, Division of Sport Fish, Research and Technical Services, Fishery Data Series No. 09-21, 9 p.

Vollen, T., and Albert, O.T., 2008, Pelagic behavior of adult Greenland halibut (*Reinhardtius hippoglossoides*): Fishery Bulletin, v. 106, no. 4, p. 457–470.

von Biela, V.R., Zimmerman, C.E., and Moulton, L.L., 2011, Long-term increases in young-of-the-year growth of Arctic cisco *Coregonus autumnalis* and environmental influences: Journal of Fish Biology, v. 78, no. 1, p. 39–56.

Walkusz, W., Paulic, J.E., Williams, W.J., Kwasniewski, S., and Papst, M.H., 2011, Distribution and diet of larval and juvenile Arctic cod (*Boreogadus saida*) in the shallow Canadian Beaufort Sea: Journal of Marine Systems, v. 84, p. 78–84.

Wang, M., and Overland, J.E., 2009, A sea ice free summer Arctic within 30 years?: Geophysical Research Letters, v. 36, no. 7, L07502.

Wassmann, P., Duarte, C.M., Agust, S., and Sejr, M.K., 2011, Footprints of climate change in the Arctic marine ecosystem: Global Change Biology, v. 17, no. 2, p. 1,235–1,249.

Weingartner, T.J.,1997, A review of the physical oceanography of the northeastern Chukchi Sea, *in* Reynolds, J., ed., Fish ecology in Arctic North America: Bethesda, Md., Special Publication (No. 19) of the American Fisheries Society, p. 40–59.

Weingartner, T., Shimada, K., McLaughlin, F., and Proshutinsky, A., 2008, Physical oceanography, *in* Hopcroft, R., Bluhm, B., and Gradinger, R., eds., Arctic Ocean synthesis—Analysis of climate change impacts in the Chukchi and Beaufort Seas with strategies for future research: Institute of Marine Science, University of Fairbanks, Alaska, p. 6–17.

Weslawski, J.M., Kwaśniewski, S., Stempniewicz, L., and Blachowiak-Samolyk, K., 2006, Biodiversity and energy transfer to top trophic levels in two contrasting Arctic fjords: Polish Polar Research, v. 27, no. 3, p. 259–278.

Welch, H.E., Crawford, R.E., and Hop, H., 1993, Occurrence of Arctic cod (*Boreogadus saida*) schools and their vulnerability to predation in the Canadian High Arctic: Arctic, v. 46, no. 4, p. 331–339.

Wells, R.M.G., 2005, Blood-gas transport and hemoglobin function in polar fishes—Does low temperature explain physiological characters?, *in* Farrell, A.P. and Steffensen, J.F., eds., The physiology of polar fishes: Amsterdam, Elsevier, p. 281–316.

Wernberg T., Smale, D.A., and Thomsen, M.S., 2012, A decade of climate change experiments on marine organisms—Procedures, patterns, and problems: Global Change Biology, v. 18, no. 5, p. 1,491–1,498.

West, R.L., and Fruge, D.J., 1989, A review of coastal plain fish surveys and the results of 1986 fish surveys of selected coastal lakes and streams Arctic National Wildlife Refuge, Alaska: Alaska Fisheries Technical Report, no. 4, 22 p. Wheeler, A., 1992, A list of common and scientific names of fishes of the British Isles, Fisheries Society of the British Isles, Academic Press, London.

White House, 2013, National strategy for the Arctic Region: Washington D.C., Office of the President, 13 p. https:// www.whitehouse.gov/sites/default/files/docs/nat_arctic_ strategy.pdf.

Whiteley, N.M., Christiansen, J.S., and Egginton, S., 2006, Polar cod, *Boreogadus saida* (Gadidae), show an intermediate stress response between Antarctic and temperate fishes: Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology, v. 145, no. 4, p. 493–501.

Wienerroither, R., Johannesen, E., Langøy, H., Børve Eriksen, K., de Lange Wenneck, T., Høines, Å., Bjelland, O., Aglen, A., Prokhorova, T., Murashko, P., Prozorkevich, D., Konstantin Byrkjedal, I., Langhelle Drevetnyak, G., and Smirnov, O., 2011, Atlas of the Barents Sea fishes: IMR/ PINRO Joint Report Series 1-2011, ISSN 1502-8828, 274 p.

Wiese, F.K., Merrick, R., Auad, G., Williams, D., Stabeno, P., and Kelly, B., 2013, Strawman conceptual ecosystem model for the Chukchi and Beaufort Seas, appendix B *in* Developing a conceptual model of the Arctic marine ecosystem: Interagency Arctic Research Policy Committee workshop, April 30–May 2, 2013, Washington, D.C., p. B1–B29.

Wiese, F., Auad, G., Williams, D., Merrick, R., Jeffries, M., Kelly, B., Thornhill, A., Thorsteinson, L., and D. Dickson, D., 2015, Framing Arctic marine research initiatives: A Framework for coordinated marine ecosystem research in the U.S. Chukchi and Beaufort Seas, North Pacific Research Board, Anchorage, AK., 34 p., http://www. iarpccollaborations.org/uploads/cms/documents/framing_ arctic_marine_research_initiatives_report_p2b_lowres.pdf.

Wildermuth, D.A., 1983, Length-weight regression analysis for thirty-eight species of sport caught marine fishes: Western Department of Fisheries Progress Report, no. 189, 7 p.

Wilderbuer, T.K., Walters, G.E., and Bakkala, R.G., 1992, Yellowfin sole, *Pleuronectes asper*, of the eastern Bering Sea—Biological characteristics, history of exploitation, and management: Marine Fisheries Review, v. 54, no. 4, p. 1–18.

Wilimovsky, N.J., 1974, Fishes of the Bering Sea—The state of existing knowledge and requirements for future effective efforts, *in* Hood, D.W., and Kelly, E.J., eds., Oceanography of the Bering Sea with emphasis on renewable resources: Proceedings of the International Symposium, Bering Sea Study, Hakodate, Japan, January 31–February 4, 1972, Institute Marine Science, University of Alaska, Fairbanks, Occasional Publication No. 2., p. 243–256. Williams, B.K., Nichols, J.D., and Conroy, M.H., 2002, Analysis and management of animal populations: San Diego, Academic Press, 817 p.

Willson, M.F., Armstrong, R.H., Robards, M.D., and Piatt, J.F., 1999, Sand lance as cornerstone prey for predator populations, *in* Robards, M.D., and others, eds., Sand land—A review of biology and predator relations and annotated bibliography: U.S. Forest Service, Pacific Northwest Research Station, Research Paper PNW-RP-521.

Wilson, W.J., and Ormseth, O.A., 2009, A new management plan for Arctic waters of the United States: Fisheries, v. 34, no. 11, p. 555–558.

Winemiller, K.O., 2004, Life history strategies, population regulation, and implications for fisheries management: Canadian Journal of Fisheries and Aquatic Sciences, v. 62, no. 4, p. 872–885.

Winemiller, K.O, 2005, Life history strategies, population regulation, and implications for fisheries management: Canadian Journal of Fisheries and Aquatic Sciences, v. 62, p. 872–885.

Winemiller, K.O., and Rose, K.A., 1992, Patterns of life-history diversification in North American fishes— Implications for population regulation: Canadian Journal of Fisheries and Aquatic Sciences, v. 49, no. 10, p. 2,196–2,218.

Wing, B.L., 1985, Salmon stomach contents from the Alaska troll logbook program 1977–84: U.S. Department of Commerce, National Oceanic and Atmospheric Administration National Marine Fisheries Service, Technical Memorandum NMFS F/NWC-91, 43 p.

Winters, G.H., 1970, Biological changes in coastal capelin from the over-wintering to the spawning condition: Journal of the Fisheries Board of Canada, v. 27, no. 12, p. 2,215–2,224.

Wiswar, D.W., 1994, Summer distribution of Arctic fishes in the 1002 Area of the Arctic National Wildlife Refuge, Alaska, 1991, with special emphasis on selected lakes, tundra streams, and the Sadlerochit River drainage: U.S. Fish and Wildlife Service, Alaska Fisheries Technical Report Number 27, 47 p.

Wiswar, D.W., and Frugé, D.J., 2006, Fisheries investigations in western Camden Bay, Arctic National Wildlife Refuge, Alaska, 1987: U.S. Fish and Wildlife Service, Alaska Fisheries Data Series 2006-10, 49 p.

Wiswar, D.W., West, R.L., and Winkleman, W.N., 1995, Fisheries investigation in Oruktalik Lagoon, Arctic National Wildlife Refuge, Alaska, 1986: U.S. Fish and Wildlife Service, Alaska Fisheries Technical Report, no. 30, 38 p. Wohlschlag, D.E., 1960, Metabolism of an Arctic fish and the phenomenon of cold adaptation: Ecology, v. 41, no. 2, p. 287–292.

Wolotira, R.J., Jr., Sample, T.M., and Morin, M., Jr., 1977, Demersal fish and shellfish resources of Norton Sound, the southeastern Chukchi Sea, and adjacent waters in the baseline year 1976: Seattle, Wash., Northwest and Alaska Fisheries Center, NMFS, National Oceanic and Atmospheric Administration, Processed Report, 69 p.

Wood, G., 2008, Thresholds and criteria for evaluating and communicating impact significance in environmental studies—'See no evil, hear no evil, speak no evil'?: Environmental Impact Assessment Review, v. 28, no. 1, p. 22–38.

Wood, S.N., 2003, Thin plate regression splines: Journal of the Royal Statistical Society Series B, Statistical Methodology, v., 65, no. 1, p. 95–114.

Wood, S.N., 2006, Low-rank scale-invariant tensor product smooths for generalized additive mixed models: Biometrics, v. 62, no. 4, p. 1,025–1,036.

Wood, S.N., 2011, Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models: Journal of the Royal Statistical Society Series B, v. 73, no. 1, p. 3–36.

Woodgate, R.A., K.M. Stafford, K.M., and Prahl, F.G., 2015. A synthesis of year-round interdisciplinary mooring measurements in the Bering Strait (1990-1994) and the RUSALCA years (2004–2011), Oceanography, v. 28, no. 3, p. 45–67.

Wu, Y., and Fletcher, G.L., 2000, Efficacy of antifreeze protein types in protecting liposome membrane integrity depends on phospholipid class: Biochimica et Biophysica Acta (BBA), General Subjects, v. 1524, no. 1, p. 11–16.

Wu, Y., Kao, M.H., Goddard, S.V., Banoub, J., and Fletcher, G.L., 2001, Antifreeze glycoproteins—Relationship between molecular weight, thermal hysteresis and the inhibition of leakage from liposomes during thermotropic phase transition: Comparative Biochemistry and Physiology Part B—Biochemistry and Molecular Biology, v. 128, no. 2, p. 265–273.

Woodson, D., Dello, K., Flint, L., Hamilton, R., Neilson, R., and Winton, J., 2011, Climate change effects in the Klamath Basin, chap. 6 *of* Thorsteinson, L., VanderKooi, S., and Duffy, W., eds., Proceedings of the Klamath Basin Science Conference, Medford, Oregon, February 1–5, 2010: U.S. Geological Survey Open-File Report 2011-1196, p. 123– 149. [Also available at http://pubs.er.usgs.gov/publication/ ofr20111196/.]

Wyllie Echeverria, T., 1996, Sea-ice conditions and the distribution of walleye pollock (*Theragra chalcogramma*) on the Bering and Chukchi Shelf, *in* Beamish, R.J., ed., Climate change and northern fish populations: Canadian Special Publication Fisheries and Aquatic Sciences 121, p. 131–136.

Yager, T., Kirchman, D., Sherr, E., Sherr, B., and Lovejoy, C., 2008, Microbes, in Hopcroft, R., Bluhm, B., and Gradinger, R., eds., Arctic Ocean synthesis—Analysis of climate change impacts in the Chukchi and Beaufort Seas with strategies for future research: Institute of Marine Sciences, University of Alaska, Fairbanks, p. 38–44.

Yang, M.-S., 1996, Diets of the important groundfishes in the Aleutian Islands in summer 1991: U.S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Memorandum NMFS-AFSC-60, Alaska Fisheries Science Center, 105 p.

Yang, M.-S., 2003, Food habits of the important groundfishes in the Aleutian Islands in 1994 and 1997: National Marine Fisheries Service, AFSC Processed Report 2003-07, 233 p.

Yang, M.-S., Dodd, K., Hibpshman, R., and Whitehouse, A., 2006, Food habits of groundfishes in the Gulf of Alaska in 1999 and 2001: U.S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Memorandum, NMFS-AFSC-164, 189 p. Yang, M.-S., and Nelson, M.W., 2000, Food habits of the commercially important groundfishes in the Gulf of Alaska in 1990, 1993, and 1996: Seattle, Washington, U.S. Department of Commerce, National Oceanic and Atmospheric Administration, NOAA Technical Memorandum NMFS-AFSC-112, 174 p.

Yoshihara, H.T., 1973, Annual progress report for monitoring and evaluation of Arctic waters with emphasis on the North Slope: Alaska Department of Fish Game, Division of Sport Fisheries, 49 p.

Zenger, H.H., Jr., 2004, Data report—2002 Aleutian Islands bottom trawl survey: U.S. Department of Commerce, National Oceanic and Atmospheric Administration Technical Memorandum NMFS-AFSC-143, 247 p.

Zhang, C.I., Wilderbuer, T.K., and Walters, G.E., 1998, Biological characteristics and fishery assessment of Alaska plaice, *Pleuronectes quadritubercultus*, in the eastern Bering Sea: Marine Fisheries Review, v. 60, no. 4, p. 16–27.

Zimmerman, C.E., Ramsey, A.M., Turner, S.M., Mueter, F.J., and Murphy, S.M., 2013, Genetics, recruitment, and migration patterns of Arctic cisco (*Coregonus autumnalis*) in the Colville River, Alaska, and Mackenzie River, Canada: Polar Biology, v. 36, no. 11, p. 1,543–1,555.