


Outer Continental Shelf

Estimated Oil and Gas Reserves Gulf of Mexico OCS Region December 31, 2017









Outer Continental Shelf

Estimated Oil and Gas Reserves Gulf of Mexico OCS Region December 31, 2017

Authors

Grant L. Burgess Kellie K. Cross Eric G. Kazanis

The Office of Resource Evaluation Reserves Section

Published by

U.S. Department of the Interior Bureau of Ocean Energy Management Gulf of Mexico OCS Region

New Orleans May 2019

TABLE OF CONTENTS

	Page
FIGURES	iii
TABLES	iv
ABBREVIATIONS AND ACRONYMS	V
ABSTRACT	v i
INTRODUCTION	1
BACKGROUND	
Methods Used for Estimating Reserves.	
RESERVES AND RELATED DATA BY PLANNING AREA	4
FIELD-SIZE DISTRIBUTION	6
RESERVOIR-SIZE DISTRIBUTION	11
DRILLING AND PRODUCTION TRENDS	
DEVELOPMENT BY ASSESSMENT UNIT	15
SUMMARY AND CONCLUSIONS	
Comparison of Reserves	
CONTRIBUTING PERSONNEL	25
REFERENCES	26
APPENDIX A _ Definitions of Field Resource and Reserves Terms	27

FIGURES

	P	age
Figure 1.	BOEM GOM Production and Reserves.	1
Figure 2.	BOEM resource classification framework.	2
Figure 3.	BOEM GOM OCS Planning Areas and Protraction Areas.	5
Figure 4.	Field-size Distribution of all GOM Fields by Planning Area	7
Figure 5.	Field-size Distribution of GOM Oil Fields by Planning Area.	7
Figure 6.	Field-size Distribution of GOM Gas Fields by Planning Area	7
Figure 7.	Cumulative percent total reserves versus rank order of field size	8
Figure 8.	Largest 20 fields, with associated water depths, ranked by Reserves and compared to Original Reserves	9
Figure 9.	Reservoir-size distribution, combination reservoirs	.11
Figure 10.	Reservoir-size distribution, oil reservoirs	.12
Figure 11.	Reservoir-size distribution, gas reservoirs	12
Figure 12.	Number of exploratory wells drilled by water depth	.13
	Number of development wells drilled by water depth	
Figure 14.	Original Reserves categorized by water depth and reservoir discovery year	14
Figure 15.	Annual oil and gas production	14
Figure 16A	. Pleistocene Shelf Development	.16
Figure 16B	. Pleistocene Slope Development	.16
Figure 17A	Pliocene Shelf Development	.17
Figure 17B	. Pliocene Slope Development	.17
Figure 18A	. Upper Miocene Shelf Development.	18
Figure 18B	. Upper Miocene Slope Development	18
Figure 19A	. Middle Miocene Shelf Development	19
Figure 19B	. Middle Miocene Slope Development	.19
Figure 20A	. Lower Miocene Shelf Development	20
Figure 20B	. Lower Miocene Slope Development	.20
Figure 21.	Lower Tertiary Slope Development.	21
Figure 22.	James Play Development	.21
Figure 23.	Norphlet Play Development	22

TABLES

Table 1.	Estimated oil and gas reserves by area, December 31, 2017	4
Table 2.	Description of deposit-size classes.	6
Table 3.	Field and reserves distribution by water depth	8
Table 4.	Fields by rank order, based on Original BOE reserves, top 50 fields	10
Table 5.	Summary and comparison of GOM oil and gas reserves as of December 31, 2016, and December 31, 2017.	23
Table 6.	Oil and gas reserves and cumulative production at end of year, 1975-2017	24

ABBREVIATIONS AND ACRONYMS

AL Alabama

Bbbl Billion barrels

Bbl barrels

BBO billion barrels of oil

BBOE billion barrels of oil equivalent

Bcf billion cubic feet

BOE barrels of oil equivalent BOEM Bureau of Ocean Energy

Management

DOI U.S. Department of the Interior

⁰F degrees Fahrenheit

FL Florida ft feet

GOM Gulf of Mexico

GOMR Gulf of Mexico Region

GOR gas oil ratio
LA Louisiana
MMbbl million barrels

MMBOE million barrels of oil equivalent

MMcf million cubic feet

MMS Minerals Management Service

MS Mississippi N north

OAP Offshore Atlas Project
OCS Outer Continental Shelf

ONRR Office of Natural Resources Revenue psia pounds per square inch absolute P/Z pressure/gas compressibility factor

RE Resource Evaluation

SCF/STB standard cubic feet per stock tank barrel

SPE-PRMS Society of Petroleum Engineers Petroleum Resources

Management System

Tcf trillion cubic feet

TX Texas

U.S. United States

USGS United States Geological Survey

ABSTRACT

This publication presents the Bureau of Ocean Energy Management's (BOEM) estimates of oil and gas reserves in the Gulf of Mexico Outer Continental Shelf. As of December 31, 2017, it is estimated that the *Original Reserves* are 24.65 billion barrels of oil and 195.2 trillion cubic feet of gas from 1,319 fields. *Original Reserves* are the total of the *Cumulative Production* and the *Reserves*. This number includes one new discovery, MC768, and three fields that moved from *Resources* to *Reserves* during 2017: GC627, MC427 and MC794. It also includes 830 fields that have produced and expired. *Cumulative Production* from all fields accounts for 20.78 billion barrels of oil and 188.9 trillion cubic feet of gas.

Reserves are estimated to be 3.87 billion barrels of oil and 6.3 trillion cubic feet of gas. These reserves are recoverable from 489 active fields. Reserves in this report are proved plus probable (2P) reserves estimates. The reserves must be discovered, recoverable, commercial and remaining. Reserves, starting with the 2011 report, now include Reserves Justified for Development.

The estimates of reserves for this report represent the combined efforts of engineers, geoscientists, paleontologists, petrophysicists, and other personnel of the BOEM Gulf of Mexico Region, Office of Resource Evaluation, in New Orleans, Louisiana. Reserves estimates are derived for individual reservoirs from geologic and engineering calculations. For any field spanning State and Federal waters, reserves are estimated for the Federal portion only.

INTRODUCTION

This report supersedes the <u>Estimated Oil and Gas Reserves</u>, <u>Gulf of Mexico OCS Region</u>, <u>December 31</u>, <u>2016</u> (Burgess et al., 2018). It presents estimated Original Reserves, Cumulative Production, and Reserves as of December 31, 2017, for the Gulf of Mexico (GOM). **Figure 1** represents the percentages of Cumulative Production and Reserves in the GOM. Contingent and Undiscovered Resources are not included in this report.

As of December 31, 2017, the 1,319 oil and gas fields in the federally regulated part of the Gulf of Mexico Outer Continental Shelf (GOM OCS) contained Original Reserves estimated to be 24.65 billion barrels of oil (BBO) and 195.2 trillion cubic feet (Tcf) of gas. Cumulative Production from the fields accounts for 20.78 BBO and 188.9 Tcf of gas. Reserves are estimated to be 3.87 BBO and 6.3 Tcf of gas for the 489 active fields. Oil Reserves have increased 8.4 percent and the gas Reserves have decreased 7.4 percent since the 2016 report.




Figure 1. BOEM GOM Production and Reserves

BACKGROUND

Classification of Resources and Reserves

The BOEM resource classification framework is shown in Figure 2. Definitions for each resource class are presented in Appendix A. At the point in time a discovery is made, the identified accumulation of hydrocarbons is classified as a Contingent Resource, since a development project has not yet been identified. When the lessee makes a formal commitment to develop and produce the accumulation, it is classified as a Reserves Justified for Development. During the period when infrastructure is being constructed and installed, the accumulation is classified as Undeveloped Reserves. After the equipment is in place the accumulation is classified as Developed Non-Producing Reserves, and when production of the accumulation has begun, the status becomes Developed Producing Reserves. If an accumulation goes off production, for a year or more, for any reason, the classification changes back to Developed Non-Producing. Reserves in this report are proved plus probable (2P) reserves estimates. This is based on the classifications recommended in *Petroleum Resource* Management System (2007) which account for the range of uncertainty associated with reserve/resource estimation. For example, a 1P estimate would include only proved reserves, while a 3P estimate would incorporate proved, probable, and possible reserves. The reserves must be discovered, recoverable, commercial and remaining. Reserves, starting with the 2011 report, now include Reserves Justified for Development. All hydrocarbons produced and sold are included in the Cumulative Production category. Should a project be abandoned, at any phase of development, any estimates of remaining hydrocarbon volumes could be reclassified to Contingent Resources.




Figure 2. BOEM resource classification framework.

Methods Used for Estimating Reserves

The Reserves inventory component of the Resource Evaluation (RE) Program incorporates new producible leases into fields and develops independent estimates of recoverable amounts of oil and gas contained within discovered fields. The RE Program also develops independent estimates of natural gas and oil in previously discovered OCS fields by conducting field reserve studies and reviews of fields, sands, and reservoirs. The Program periodically revises the estimates of natural gas and oil volumes to reflect new discoveries, development, and annual production. This report, *Estimated Oil and Gas Reserves, Gulf of Mexico OCS Region, December 31, 2017*, is based on field studies completed at the reservoir and sand levels. All of the reservoir level data have been linked to the sand, pool, play, chronozone, and series level to support the Offshore Atlas Project (OAP).

Additional reports address GOM reserves and undiscovered resources on the OCS. Minerals Management Service (MMS) OCS Report, *Atlas of Gulf of Mexico Gas and Oil Sands as of January 1, 1999* (Bascle et al., 2001) provides a detailed geologic reporting of oil and gas reserves. A brief summary of the Atlas is available on the BOEM's Web site at https://www.boem.gov/BOEM-Newsroom/Library/Publications/2001/2001-086.aspx and current Atlas data associated with the 2017 Estimated Oil and Gas Report are available at https://www.data.boem.gov/Main/GandG.aspx. The BOEM Report, 2016a National Assessment of Undiscovered Oil and Gas Resources of the U.S. Outer Continental Shelf, summarizes the results of the Bureau of Ocean Energy Management 2016 assessment of the undiscovered oil and gas resources for the U.S. Outer Continental Shelf. For more information visit BOEM's Web site at https://www.boem.gov/National-Assessment-2016/.

Reserve estimates from geological and engineering analyses have been completed for the 1,319 fields. The accuracy of the reserve estimate improves as additional reservoir data becomes. Well logs, well file data, seismic data, and production data are periodically analyzed to improve the accuracy of the reserve estimate. As a field is depleted and/or abandoned, the Original Reserves of productive reservoirs are assigned a value equal to the amount produced and any unrecovered reserve volumes may be converted to Contingent Resources. Currently, there are 830 expired, depleted fields.

Methods used for estimating reserves can be categorized into three groups: analog, volumetric, and performance. Reserve estimates in this report are based primarily on volumetric and performance methods. Reserve estimates are reported deterministically, providing a single "best estimate" based on known geological, engineering, and economic data.

Production data are the metered volumes of raw liquids and gas reported to BOEM (from ONRR, Office of Natural Resources Revenue) by Federal OCS unit and lease operators. Metered volumes from production platforms and/or leases are allocated to individual wells and reservoirs on the basis of periodic well test gauges. These procedures introduce approximations in both production and remaining reserves volumes.

Oil and gas volume measurements and reserves are corrected to reference standard conditions of 60°F and one atmosphere (14.73 pounds per square inch absolute [psia]). Prior to September 1998, gas was reported at 15.025 psia. BOEM has converted all historical gas production volumes to the 14.73 pressure base.

RESERVES AND RELATED DATA BY PLANNING AREA

The GOM OCS is divided into three planning areas for administrative purposes (**Figure 3**). Each planning area is subdivided into protractions, which in turn are divided into numbered blocks. Fields in the GOM are identified by the protraction area name and block number of discovery – for example, East Cameron Block 271 (EC 271) Field. As the field is developed, the limits may expand into adjacent blocks and planning areas. These adjacent blocks are then identified as part of the original field and are added to that field. Statistics in this report are presented as area totals compiled under each field name. For example, all of the data associated with EC 271 Field are included in the East Cameron totals, although part of the field extends into the adjacent area of Vermilion. There are four exceptions: Tiger Shoal and Lighthouse Point, included in South Marsh Island; Coon Point, included in Ship Shoal; and Bay Marchand, included in South Timbalier.

As of December 31, 2017, there were 489 fields active in the federally regulated part of the GOM. A list, updated quarterly, of the active and expired fields can be found in the <u>OCS Operations Field Directory</u>. Included are the 830 expired, depleted and/or abandoned fields that produced 23.8 percent of the total cumulative GOM oil and gas production (by barrels oil equivalent (BOE)). One hundred thirty three fields expired, relinquished, or terminated without production. These fields may be included in the <u>Indicated Hydrocarbon List</u>. Reserves data are presented as area totals in **Table 1**.

Table 1. Estimated oil and gas reserves by area, December 31, 2017.

		Number	of fields						Cumulative	9				
Area(s) (Fig. 3)	Active	Active nonprod	Expired depleted	Expired nonprod	Ori	Original Reserves			Production hrough 201		Reserves			
(-3)	prod				Oil (MMbbl)	Gas (Bcf)	BOE (MMbbl)	Oil (MMbbl)	Gas (Bcf)	BOE (MMbbl)	Oil (MMbbl)	Gas (Bcf)	BOE (MMbbl)	
Western Planning Area					,	(- /	,	(,	(-)	, ,,,	(,	(- /	, ,,,	
Alaminos Canyon	4	0	1	5	360	644	474	254	435	331	106	209	143	
Brazos	3	0	35	3	10	3,754	678	10	3,730	674	0	24	4	
East Breaks	10	0	11	5	274	2,227	670	264	2,162	649	10	65	21	
Galveston	3	1	46	3	67	2,229	464	66	2,213	460	1	16	4	
Garden Banks	1	0	6	2	39	330	98	37	326	95	2	4	3	
High Island and Sabine Pass	23	7	99	10	426	15,531	3,189	420	15,471	3,173	6	60	16	
Keathley Canyon	0	0	0	1	0	0	0	0	0	0	0	0	0	
Matagorda Island	3	0	26	2	24	5,274	962	24	5,261	960	0	13	2	
Mustang Island	2	0	27	5	8	1,794	328	8	1,784	326	0	10	2	
N.& S.Padre Island	0	0	19	0	0	625	112	0	625	112	0	0	0	
Port Isabel	0	0	0	1	0	0	0	0	0	0	0	0	0	
West Cameron and Sabine Pass	2	1	23	1	36	2,932	559	35	2,920	555	1	12	4	
Western Planning Area Subtotal	51	9	293	38	1,244	35,340	7,534	1,118	34,927	7,335	126	413	199	
Central Planning Area										,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,				
Atwater Valley	1	1	4	6	65	609	173	39	590	144	26	19	29	
Chandeleur	1	0	13	0	0	385	69	0	385	69	0	0	0	
Desoto Canyon	3	0	3	4	15	517	106	7	509	97	8	8	9	
Destin Dome	0	0	0	1	0	0	0	0	0	0	0	0	0	
East Cameron	12	2	53	0	368	11,026	2,329	357	10,962	2,307	11	64	22	
Eugene Island	36	0	53	4	1,784	20,651	5,458	1,721	20,290	5,331	63	361	127	
Ewing Bank	12	2	4	2	440	822	586	369	713	496	71	109	90	
Garden Banks	14	1	17	5	866	4,446	1,658	804	4,195	1,551	62	251	107	
Grand Isle	9	2	12	1	1,033	5,146	1,948	1,003	5,020	1,896	30	126	52	
Green Canyon	32	2	13	26	3,738	4,394	4,520	2,460	3,523	3,087	1,278	871	1,433	
Keathley Canyon	2	0	0	3	164	560	262	72	331	130	92	229	132	
Lloyd Ridge	0	0	4	1	0	330	59	0	330	59	0	0	0	
Main Pass and Breton Sound	29	6	57	4	1,206	7,156	2,479	1,181	7,003	2,427	25	153	52	
Mississippi Canyon	39	6	18	11	4,754	11,753	6,847	3,377	9,565	5,080	1,377	2,188	1,767	
Mobile	8	1	25	2	0	2,433	433	0	2,353	419	0	80	14	
Pensacola	0	0	1	0	0	8	1	0	8	1	0	0	0	
Ship Shoal	36	4	29	3	1,516	12,951	3,821	1,466	12,661	3,719	50	290	102	
South Marsh Island	27	7	17	0	1,000	15,126	3,692	967	14,840	3,608	33	286	84	
South Pass	8	0	5	1	1,120	4,590	1,938	1,107	4,525	1,913	13	65	25	
South Pelto	5	1	3	0	162	1,194	374	159	1,173	367	3	21	7	
South Timbalier	23	2	38	3	1,632	10,481	3,498	1,607	10,332	3,446	25	149	52	
Vermilion	26	5	54	1	599	16,857	3,599	584	16,675	3,551	15	182	48	
Viosca Knoll	17	1	35	8	658	3,751	1,324	620	3,639	1,267	38	112	57	
Walker Ridge	6	1	0	5	649	145	675	143	30	148	506	115	527	
West Cameron and Sabine Pass	20	5	69	0	200	18,698	3,527	195	18,518	3,490	5	180	37	
West Delta	10	4	10	3	1,439	5,830	2,477	1,425	5,755	2,449	14	75	28	
Central Planning Area Subtotal	376	53	537	94	23,408	159,859	51,853	19,663	153,925	47,052	3,745	5,934	4,801	
Eastern Planning Area					,	,	,	,	,	,	-,	.,		
Destin Dome	0	0	0	1	0	0	0	0	0	0	0	0	0	
Eastern Planning Area Subtotal***	0	0	0	1	0	0	0	0	0	0	0	0	0	
GOM Total:	427	62	830	133	24,652	195,199	59,387	20,781	188,852	54,387	3,871	6,347	5,000	
		1,319				,	,	.,	,	. ,	-,-	-,-	-,	




Figure 3. BOEM GOM OCS Planning Areas and Protraction Areas.

FIELD-SIZE DISTRIBUTION

Field Reserve volumes are expressed in terms of BOE. Gas reserves are converted to BOE and added to the liquid reserves for the convenience of comparison. The conversion factor of 5,620 standard cubic feet of gas equals 1 BOE is based on the average heating values of domestic hydrocarbons. A geometric progression, developed by the United States Geological Survey (USGS) (Attanasi, 1998), was selected for field-size (deposit-size) distribution ranges (**Table 2**).

In this report, fields are classified as either oil or gas; some fields do produce both products, making a field type classification difficult. The classification is made on a case-by-case basis by analysis of the field's reservoirs and their fluid distributions.

Table 2. Description of deposit-size classes.

Class	Deposit-size range*	Class	Deposit-size range*	Class	Deposit-size range*					
1	0.031 - 0.062	10	16 - 32	18	4,096 - 8,192					
2	0.062 - 0.125	11	32 -64	19	8,192 - 16,384					
3	0.125 - 0.25	12	64 - 128	20	16,384 - 32,768					
4	0.25 - 0.50	13	128 - 256	21	32,768 - 65,536					
5	0.50 - 1.00	14	256 - 512	22	65,536 - 131,072					
6	1 - 2	15	512 - 1,024	23	131,072 - 262,144					
7	2 - 4	16	1,024 - 2,048	24	262,144 - 524,288					
8	4 - 8	17	2,048 - 4,096	25	524,288 - 1,048,576					
9	8 - 16	*N	*Million Barrels of Oil Equivalent (MMBOE)							

The field-size distribution based on Original Reserves (in BOE) for 1,319 fields is shown in **Figure 4**, along with the planning area distributions. Of the 1,319 oil and gas fields, there are 266 oil fields represented in **Figure 5** and 1,053 gas fields shown in **Figure 6**. These figures also display the planning area distributions.

Analysis of the 1,319 oil and gas fields indicates that the GOM is historically a gas-prone basin. The GOR, based on original reserves of the 266 oil fields, is 2,357 SCF/STB. The yield (condensate divided by gas), based on original reserves for the 1,053 gas fields, is 26.1 barrels (Bbl) of condensate per million cubic feet (MMcf) of gas.




Figure 4. Field-size Distribution of all GOM Fields by Planning Area




Figure 5. Field-size Distribution of GOM Oil Fields by Planning Area




Figure 6. Field-size Distribution of GOM Gas Fields by Planning Area

Figure 7 shows the cumulative percent distribution of Original Reserves in billion barrels of oil equivalent (BBOE), by field size rank. All 1,319 fields in the GOM OCS are included in this figure. A phenomenon often observed in hydrocarbon-producing basins is a rapid drop-off in size from that of largest known field to smallest. Twenty-five percent of the Original Reserves are contained in the 27 largest fields. Fifty percent of the Original Reserves are contained in the 91 largest fields. Ninety percent of the Original Reserves are contained in the 438 largest fields.




Figure 7. Cumulative percent total reserves versus rank order of field size.

Table 3 shows the distribution of the number of fields and reserves by water depth. A field's water depth is determined by averaging the water depth where the wells are drilled in the field. Reserves and production, reported in MMBOE, are associated with the 1,319 fields. Reserves located in greater than or equal to 1,500 ft of water accounts for 83 percent of the total GOM Reserves.

Table 3. Field and reserves distribution by water depth.

Water Depth Range (Feet)	Number of Fields	Cumulative Production (MMBOE)	Reserves (MMBOE)
< 500	1,082	41,608	697
500 - 999	54	1,276	33
1,000 - 1,499	28	1,446	120
1,500 - 4,999	100	7,045	2,398
5,000 - 7,499	37	2,414	1,505
>= 7,500	18	598	247
Totals:	1,319	54,387	5,000

Figure 8 shows the largest 20 fields ranked in order by Reserves. All 20 of the fields lie in water depths of greater than or equal to 1,500 ft and account for 64.3 percent of the Reserves in the GOM. Of the 237 fields in water depths greater than 500 ft, 133 are producing, 96 are depleted or expired, and 8 have yet to produce.




Figure 8. Largest 20 fields, with associated water depths, ranked by Reserves and compared to Original Reserves.

Table 4 ranks the 50 largest fields based on Original Reserves expressed in BOE. Rank, Field name, Field Nickname, Discovery year, Water depth, Field Classification, Field type, Field GOR, Original Reserves, Cumulative Production through 2017 and Reserves are presented. A complete listing of all 1,319 fields is available on the BOEM Web site at: https://www.data.boem.gov/Main/HtmlPage.aspx?page=estimated2017.

Table 4. Fields by rank order, based on Original BOE reserves, top 50 fields

 $Table \ 4. \ A \ listing \ of \ Gulf \ of \ Mexico \ fields \ by \ rank \ order, \ based \ on \ Original \ BOE \ reserves, \ top \ 50 \ fields.$

Field class: P (PDP - Developed Producing, PDN - Developed Non-Producing and PU - Undeveloped); J (RJD- Reserves Justified for Development) Field type: O - Oil; G - Gas

F	Field				Cumulative Production through 2017			Reserves							
Rank n	ame	Nickname	year	depth c	lass type	GOR (SCF/STB)	Oil (MMbbl)	Gas (Bcf)	BOE (MMbbl)	Oil (MMbbl)	Gas (Bcf)	BOE (MMbbl)	Oil (MMbbl)	Gas (Bcf)	BOE (MMbbl)
1 MC	2807	MARS-URSA	1989	3,341 P	0	1,359	1871.2	2542.7	2323.7	1352.9	1740.9	1662.7	518.3	801.8	661.0
2 El3	330		1971	248 P	0	4,081	466.6	1904.5	805.4	455.9	1889.3	792.0	10.7	15.2	13.4
3 W [D030		1949	48 P	0	1,650	594.5	983.2	769.5	590.7	973.3	763.9	3.8	9.9	5.6
4 GI0	043		1956	140 P	0	4,365	401.9	1743.5	712.1	379.8	1654.0	674.1	22.1	89.5	38.0
5 TS	000		1958	13 P	G	79,565	46.8	3725.0	709.6	45.3	3634.0	691.9	1.5	91.0	17.7
6 GC	C640	TAHITI/CAESAR/TONGA	2002	4,339 P	0	605	616.7	373.0	683.1	322.9	208.1	359.9	293.8	164.9	323.2
7 BM	<i>I</i> 002		1949	50 P	0	1,057	547.2	578.6	650.2	544.2	574.5	646.5	3.0	4.1	3.7
8 VR	R014		1956	26 P	G	65,255	47.9	3126.5	604.2	47.9	3126.5	604.2	0	0.0	0
9 GC	C743	ATLANTIS	1998	6,303 P	0	748	494.8	369.9	560.6	323.6	213.6	361.6	171.2	156.3	199.0
10 MP	P041		1956	42 P	0	5,719	272.8	1560.0	550.3	271.4	1556.6	548.3	1.4	3.4	2.0
11 VR	2039		1948	38 P	G	79,958	32.6	2606.9	496.5	32.2	2603.8	495.5	0.4	3.1	1.0
12 GC	C654	SHENZI	2002	4,303 P	0	394	462.5	182.3	494.9	289.8	114.8	310.2	172.7	67.5	184.7
13 SS	3208		1960	102 P	0	6,110	234.0	1433.9	489.1	226.4	1399.1	475.3	7.6	34.8	13.8
14 GB	3426	AUGER	1987	2,847 P	0	3,503	289.0	1012.3	469.2	268.0	937.4	434.8	21.0	74.9	34.4
15 GC	2826	MAD DOG	1998	4,799 P	0	412	393.3	162.1	422.2	189.5	56.2	199.5	203.8	105.9	222.7
16 W	D073		1962	177 P	0	2,501	279.2	702.8	404.2	277.3	689.7	400.0	1.9	13.1	4.2
17 El2	238		1964	147 P	G	16,126	102.6	1612.7	389.5	94.7	1537.1	368.2	7.9	75.6	21.3
18 GIO	016		1948	54 P	0	1,298	310.1	402.0	381.6	307.3	396.4	377.8	2.8	5.6	3.8
19 SP	2061		1967	220 P	0	1,931	276.0	532.9	370.9	272.2	529.3	366.4	3.8	3.6	4.5
20 MC	2778	THUNDER HORSE	1999	6,095 P	0	741	322.9	239.1	365.4	145.0	109.3	164.4	177.9	129.8	201.0
21 SP	089		1969	421 P	0	4,425	198.3	877.3	354.3	196.5	873.7	351.9	1.8	3.6	2.4
22 ST	172		1962	98 P	G	158,376	12.0	1898.9	349.9	12.0	1898.9	349.9	0.0	0.0	0.0
23 MC	084	KING/HORN MT.	1993	5,315 P	0	1,064	290.3	308.9	345.2	236.7	261.6	283.2	53.6	47.3	62.0
24 W	C180		1961	48 P	G	139,651	13.2	1845.8	341.7	13	1846	342	0.0	0.0	0.0
25 MC	2776	N.THUNDER HORSE	2000	5,672 P	0	964	290.0	279.6	339.7	244.7	237.2	286.9	45.3	42.4	52.8
26 ST	021		1957	46 P	0	1,652	259.8	428.9	336.1	258.8	426.6	334.7	1.0	2.3	1.4
27 SS	169		1960	63 P	0	5,280	172.5	911.0	334.7	168.8	897.3	328.5	3.7	13.7	6.2
28 MC	C194	COGNAC	1975	1,022 P	0	4,166	185.1	770.9	322.3	182.0	761.8	317.6	3.1	9.1	4.7
29 El2	292		1964	214 P	G	61,864	26.8	1657.8	321.8	24.4	1650.6	318.1	2.4	7.2	3.7
30 ST	176		1963	127 P	G	13,936	92.3	1286.4	321.2	90.1	1272.3	316.5	2.2	14.1	4.7
31 EC	271		1971	172 P	G	17,775	76.3	1355.4	317.4	72.6	1343.8	311.7	3.7	11.6	5.7
32 EC	064		1957	50 P	G	56,112	28.7	1609.0	315.0	27.4	1606.4	313.2	1.3	2.6	1.8
33 SM	<i>1</i> 048		1961	100 P	G	51,549	30.3	1566.3	309.1	29.3	1558.7	306.7	1.0	7.6	2.4
34 SS	3176		1956	101 P	G	19,163	69.5	1332.8	306.7	68.8	1317.5	303.3	0.7	15.3	3.4
35 AC	857	GREAT WHITE	2002	7,915 P	0	2,017	218.0	439.6	296.2	138.4	256.6	184.0	79.6	183.0	112.2
36 W	C587		1971	210 P	G	118,356	13.4	1581.1	294.7	13.4	1581.1	294.7	0.0	0.0	0.0
37 SP	2027	EAST BAY	1954	64 P	0	5,166	153.1	791.6	294.0	152.2	783.9	291.7	0.9	7.7	2.3
38 ST	135		1956	129 P	0	3,699	171.8	635.4	284.9	169.8	626.5	281.3	2.0	8.9	3.6
39 W [D079		1966	123 P	0	3,881	168.3	650.2	284.1	165.3	639.8	279.2	3.0	10.4	4.9
40 El2	296		1971	214 P	G	71,230	20.6	1464.0	281.1	20.6	1464.0	281.1	0.0	0.0	0.0
41 W	C192		1954	57 P	G	60,168	23.7	1420.5	276.4	23.4	1416.3	275.4	0.3	4.2	1.0
42 GB	3171	SALSA	1984	1,212 P	G	3,961	161.5	639.7	275.3	143.7	573.2	245.7	17.8	66.5	29.6
43 HI5	573A		1973	341 P	0	7,443	118.3	880.6	275.0	116.3	877.6	272.5	2.0	3.0	2.5
44 Ml6	623		1980	83 P	G	102,244	13.8	1412.2	265.1	13.8	1410.1	264.7	0.0	2.1	0.4
45 GI0	047		1955	88 P	0	3,834	156.6	602.1	263.7	154.2	593.6	259.8	2.4	8.5	3.9
46 VK	956	RAM-POWELL	1985	3,238 P	0	9,114	100.2	912.6	262.5	99.1	905.0	260.1	1.1	7.6	2.4
47 SP	2078		1972	202 P	G	11,038	85.0	937.8	251.9	83.0	931.8	248.8	2.0	6.0	3.1
48 GC	244	TROIKA	1994	2,795 P	0	1,897	182.5	346.2	244.2	180.1	343.5	241.3	2.4	2.7	2.9
49 WF	R678	SAINT MALO	2003	6,953 P	0	245	233.4	57.2	243.5	61.0	15.1	63.6	172.4	42.1	179.9
50 SM	<i>I</i> 023		1960	82 P	G	39,519	29.8	1176.3	239.1	29.8	1176.3	239.1	0.0	0.0	0.0

RESERVOIR-SIZE DISTRIBUTION

The size distributions of the reservoirs are shown in **Figures 9, 10, and 11**. The size ranges are based on Original Reserves and are presented on a geometrically progressing horizontal scale. These sizes correspond with the USGS deposit-size ranges shown in **Table 2,** with a modification to subdivide small reservoirs into finer distributions. In **Figure 9,** the Original Reserves are presented in million barrels of Oil Equivalent (MMBOE). For the combination reservoirs (saturated oil rims with associated gas caps), shown in **Figure 9,** gas is converted to BOE and added to the liquid reserves. **Figures 10 and 11** are presented in million barrels of Oil (MMBbl) and billion cubic feet (Bcf), respectively. The number of reservoirs in each size grouping, shown as percentages of the total, is presented on a linear vertical scale.

Figure 9 shows the reservoir-size distribution, on the basis of Original BOE, for 2,374 combination reservoirs. The median is 0.9 MMBOE and the mean is 3.0 MMBOE. The GOR, based on Original Reserves, for the oil portion of the reservoirs is 1,225 SCF/STB, and the yield, based on Original Reserves, for the gas cap is 20.2 Bbl of condensate per MMcf of gas.




Figure 9. Reservoir-size distribution, combination reservoirs.

Figure 10 shows the reservoir-size distribution, on the basis of Original Oil reserves, for 8,957 undersaturated oil reservoirs. The median is 0.3 MMbbl, the mean is 2.1 MMbbl, and the GOR, based on Original Oil reserves, is 1,185 SCF/STB. **Figure 11** shows the reservoir-size distribution, on the basis of Original Gas reserves, for 18,734 gas reservoirs. The median is 2.0 Bcf of gas, the mean is 8.3 Bcf, and the yield, based on Original Reserves, is 12.3 Bbl of condensate per MMcf of gas.




Figure 10. Reservoir-size distribution, oil reservoirs.




Figure 11. Reservoir-size distribution, gas reservoirs.

DRILLING AND PRODUCTION TRENDS

Figure 12 presents the number of exploratory wells drilled each year by water depth category. The total footage drilled in 2017 was 2.02 million feet, compared to 2.03 million feet in 2016.




Figure 12. Number of exploratory wells drilled by water depth.

Figure 13 presents the number of development wells drilled each year by water depth category. The total footage drilled in 2017 was 1.19 million feet, compared to 0.89 million feet in 2016.




Figure 13. Number of development wells drilled by water depth.

Original Reserves in BBOE for water depth categories by reservoir discovery year are presented in Figure 14.




Figure 14. Original Reserves categorized by water depth and reservoir discovery year.

Annual production in the GOM is shown in **Figure 15**. The oil plot includes condensate and the gas plot includes casinghead gas. Annual production for oil and gas is presented as a total, in shallow water (less than 1,000 ft), and in deepwater (greater than 1,000 ft). From 2016 to 2017, annual oil production increased 4.6 percent to 612 MMbbl and annual gas production decreased 11.7 percent to 1.1 Tcf. The mean daily production in the GOM during 2017 was 1.61 MMbbl of crude oil, 0.10 MMbbl of gas condensate, 1.74 Bcf of casinghead gas, and 1.21 Bcf of gas-well gas. The mean GOR of oil wells was 1,081 SCF/STB, and the mean yield from gas wells was 54.0 Bbl of condensate per MMcf of gas.




Figure 15. Annual oil and gas production.

DEVELOPMENT BY ASSESSMENT UNIT

Graphical displays of reservoir and production data within assessment units and plays are provided in this section. The assessment units represent a group of geologically related hydrocarbon accumulations; the term Assessment Unit can refer to groupings of chronozone(s) and/or geologic play(s). The data from each reservoir within an assessment unit or play were combined to create graphs displaying: the total reserves volume discovered each year, the number of reservoirs discovered within the unit, the production from the reservoirs in the unit, and the average size of each reservoir in that unit.

Assessment units are based on current water depths (shelf <=200m, slope >200m), and relative geologic age of Cenozoic sediments in the U.S. Gulf of Mexico OCS. Using these criteria, Cenozoic sediments are further divided into 12 assessment units as shown below; however, only 11 of these units have figures associated with them since the Lower Tertiary Shelf unit lacks reserves or production.

Pleistocene Shelf Pleistocene Slope Pliocene Shelf Pliocene Slope

Upper Miocene Shelf
Middle Miocene Shelf
Lower Miocene Shelf
Lower Tertiary Shelf
Upper Miocene Slope
Middle Miocene Slope
Lower Miocene Slope
Lower Tertiary Slope

Unlike the aggregated assessment units of Cenozoic sediments, the Mesozoic sediments of the U.S. Gulf of Mexico OCS were differentiated by specific rock units or plays. Although 19 Mesozoic rock units and plays are identified in, <u>Assessment of Technically and Economically Recoverable Hydrocarbon Resources of the Gulf of Mexico Outer Continental Shelf as of January 1, 2014</u>, only two are included in this report: the James Play and the Norphlet Play. These two plays are included because there are reserves and production associated with them.

Figures 16 through 23 depict reservoir and production data for the 11 Cenozoic assessment units described above, and the 2 Mesozoic plays. These data exhibit the lag time from reserves discovery to production, and show exploration and development moving from shallow-water to deepwater. Shallow-water Cenozoic data exhibit significant production decline rates; however, the development of discoveries in deepwater Cenozoic sediments have offset these declines.

In Mesozoic sediments, reserves and production data exist for only 2 assessment units. These data show production rates declining in both the James Play and the Norphlet Play (Figures 22 and 23); however, additional opportunities are expected in these plays and in other Mesozoic assessment units. Expected ranges of resources to be discovered in these, and other GOM assessment units, are reported in <u>Assessment of Technically and Economically Recoverable Hydrocarbon Resources of the Gulf of Mexico Outer Continental Shelf as of January 1, 2014.</u>

Figures 16A and 16B show the decline in volume of reserves discovered, number of reservoirs discovered, and production for the shallow-water and deepwater Pleistocene assessment units. The largest total reserves discovered (MMBOE) in a single year in the Pleistocene occurred in 1974, which included two large shallow water reservoirs, one in the WC617 Field and one in the HI343A Field, containing 155 MMBOE. Both reservoirs are now depleted. The Pleistocene assessment units exhibit the greatest decline in production rates among 13 assessment units and plays displayed in this section. The data indicate this is associated with significant decreases in reserves discovered in both shallow-water and deepwater.




Figure 16A. Pleistocene Shelf Development




Figure 16B. Pleistocene Slope Development

Over the last two decades, the shallow-water Pliocene assessment unit, while maintaining a consistent average reservoir size, has seen an overall decline in total reserves discovered, number of reservoirs discovered, and production (Figure 17A). In deepwater, Pliocene production rates have been considerably higher than in shallow-water (Figure 17B). In spite of a decreasing production trend over the last decade, deepwater rates have remained within 100-150 MMBOE since 2008; some of this can be attributed to production from the KC875 Field which came online in 2015.




Figure 17A. Pliocene Shelf Development




Figure 17B. Pliocene Slope Development

For the Upper Miocene in shallow-water, both the production and number of reservoirs discovered have decreased since the year 2000 (Figure 18A). The average reservoir size has remained below 1.0 MMBOE. In deepwater, the average reservoir size and number of reservoirs discovered have remained consistent; however; production has been decreasing very slightly over the last decade (Figure 18B). A large discovery in 2014 in the MC768 field caused the average reservoir size to spike in that year.




Figure 18A. Upper Miocene Shelf Development




Figure 18B. Upper Miocene Slope Development

Other than an increase in the average reservoir size in 2015 in the MP273 Field, the total reserves discovered (MMBOE), as well as average reservoir size and production in the shallow-water Middle Miocene have all remained consistently low over the last decade (Figure 19A). In contrast, the Middle Miocene in deepwater, while seeing its maximum total reserves discovered in 2002 with the addition of 506 MMBOE in the GC640 Field, has experienced a renewed rise in production since 2013 (Figure 19B).




Figure 19A. Middle Miocene Shelf Development




Figure 19B. Middle Miocene Slope Development

The shallow-water Lower Miocene's reserves discovered (MMBOE) peaked in 1982 and 1983 (Figure 20A). A peak in production occurred in 1990, followed by an overall decline to date.




Figure 20A. Lower Miocene Shelf Development

The initial discovery of deepwater Lower Miocene reserves in 1998 yielded both the second greatest average reservoir size and third largest total reserves discovered in a single year for the assessment unit (Figure 20B). The year 2002 saw a large average reservoir size, just under 57 MMBOE. In 2002, the greatest total reserves for this play were discovered, including 363 MMBOE in a single reservoir, in the GC654 Field. While production for the Lower Miocene peaked in 2009, and had a slight decline in the two years that followed, it has been increasing consistently since 2011.




Figure 20B. Lower Miocene Slope Development

The Lower Tertiary deepwater play was discovered in 2002. This included discovery of a 136 MMBOE reservoir in the AC857 Field. The largest average reservoir size of 50 MMBOE in the play (Figure 21) occurred in 2006 with the discovery of a single reservoir in the AC857 Field. The discovery of the largest total reserves in a single year for the Lower Tertiary occurred in 2008, which included the addition of 142 MMBOE from two reservoirs in the WR678 Field. Production in this play began in 2010 and has been increasing steadily to date.




Figure 21. Lower Tertiary Slope Development

The first discovery in the James play came in 1993. In 1997, four fields yielded the largest number of reservoirs, as well as the greatest total reserves discovered (MMBOE) for the play (Figure 22). The year 1997 also yielded the largest average reservoir size. Maximum production (MMBOE) in this play peaked in 2002 with a subsequent rapid decline.




Figure 22. James Play Development

The initial Norphlet discovery in the MO823 Field in 1983 resulted in the greatest total reserves discovered for this play, along with the largest single reservoir size (Figure 23). First production began in 1991 with discoveries continuing through the mid-nineties. Peak production in this play was reached in 1997 and, while declining through the first decade of the century, has remained fairly steady to date. There have been several deepwater Norphlet discoveries to date, but these have yet to produce.




Figure 23. Norphlet Play Development

SUMMARY AND CONCLUSIONS

A summary of the Reserve estimates for 2017 and a comparison with estimates from the previous year's report (December 31, 2016) are shown in **Table 5**. Three oil and one gas field were added for this report. They are tabulated and summarized as increases to Original Reserves. One of the fields added was discovered in 2004 (MC427), two in 2014 (GC627 and MC768) and one, discovered in 2015 (MC794).

Comparison of Reserves

A net change in the Original Reserve estimates is a result of combining the discoveries and revisions. Reserve estimates may increase or decrease with additional information (e.g., additional wells are drilled, leases are added or expire, and/or reservoirs are depleted). Re-evaluations of existing field studies are conducted using field development and/or production history to capture the changes in reserve estimates. Changes in Original Reserves are presented in **Table 5**. Based on periodic reviews and revisions of field studies conducted since the 2016 report, the reserves revisions have resulted in a slight increase in Original Reserves.

The table also demonstrates that the volumes from fields added in 2017 and the field revisions slightly exceeded production, resulting in a slight increase in Reserves. The Reserves increased 8.4 percent for oil and decreased 7.4 percent for gas, since the 2016 report.

Table 5. Summary and comparison of GOM oil and gas reserves as of December 31, 2016 and December 31, 2017.

	Oil	Gas	BOE
	(Bbbl)	(Tcf)	(Bbbl)
Original Reserves:			
Previous estimate, as of 12/31/2016*	23.73	194.6	58.37
Fields Added in 2017	0.20	0.2	0.23
Revisions	0.72	0.4	0.79
Estimate, as of 12/31/2017 (this report)	24.65	195.2	59.39
Cumulative production:			
Previous estimate, as of 12/31/2016*	20.16	187.8	53.58
Revisions	0.01	0.0	0.01
Production during 2017	0.61	1.1	0.80
Estimate, as of 12/31/2017 (this report)	20.78	188.9	54.39
Reserves:			
Previous estimate, as of 12/31/2016*	3.57	6.8	4.79
Fields Added in 2017	0.20	0.2	0.23
Revisions	0.71	0.4	0.78
Production during 2017	-0.61	-1.1	-0.80
Estimate, as of 12/31/2017 (this report)	3.87	6.3	5.00

Table 6 presents all previous reserve estimates by year. Because of adjustments and corrections to production data submitted by Gulf of Mexico OCS operators, the difference between historical cumulative production for successive years does not always equal the annual production for the latter year.

Table 6. Oil and gas reserves and cumulative production at end of year, 1975-2017.

"Oil" includes crude oil and condensate; "gas" includes associated and nonassociated gas. Reserves estimated as of December 31 each year.

Year	Number of fields	Origi	inal Rese	rves		cal Cumul		Reserves			
	included	Oil	Gas	BOE	Oil	Gas	BOE	Oil	Gas	BOE	
		(Bbbl)	(Tcf)	(Bbbl)	(Bbbl)	(Tcf)	(Bbbl)	(Bbbl)	(Tcf)	(Bbbl)	
1975	255	6.61	59.9	17.27	3.82	27.2	8.66	2.79	32.7	8.61	
1976	306	6.86	65.5	18.51	4.12	30.8	9.60	2.74	34.7	8.91	
1977	334	7.18	69.2	19.49	4.47	35.0	10.70	2.71	34.2	8.80	
1978	385	7.52	76.2	21.08	4.76	39.0	11.70	2.76	37.2	9.38	
1979 ⁽¹⁾	417	7.71	82.2	22.34	4.83	44.2	12.69	2.88	38.0	9.64	
1980	435	8.04	88.9	23.86	4.99	48.7	13.66	3.05	40.2	10.20	
1981	461	8.17	93.4	24.79	5.27	53.6	14.81	2.90	39.8	9.98	
1982	484	8.56	98.1	26.02	5.58	58.3	15.95	2.98	39.8	10.06	
1983	521	9.31	106.2	28.21	5.90	62.5	17.02	3.41	43.7	11.19	
1984	551	9.91	111.6	29.77	6.24	67.1	18.18	3.67	44.5	11.59	
1985	575	10.63	116.7	31.40	6.58	71.1	19.23	4.05	45.6	12.16	
1986	645	10.81	121.0	32.34	6.93	75.2	20.31	3.88	45.8	12.03	
1987	704	10.76	122.1	32.49	7.26	79.7	21.44	3.50	42.4	11.04	
1988	678	10.95	126.7	33.49	7.56	84.3	22.56	3.39	42.4	10.93	
1989	739	10.87	129.1	33.84	7.84	88.9	23.66	3.03	40.2	10.18	
1990	782	10.64	129.9	33.75	8.11	93.8	24.80	2.53	36.1	8.95	
1991	819	10.74	130.5	33.96	8.41	98.5	25.94	2.33	32.0	8.02	
1992	835	11.08	132.7	34.69	8.71	103.2	27.07	2.37	29.5	7.62	
1993	849	11.15	136.8	35.49	9.01	107.7	28.17	2.14	29.1	7.32	
1994	876	11.86	141.9	37.11	9.34	112.6	29.38	2.52	29.3	7.73	
1995	899	12.01	144.9	37.79	9.68	117.4	30.57	2.33	27.5	7.22	
1996	920	12.79	151.9	39.82	10.05	122.5	31.85	2.74	29.4	7.97	
1997	957	13.67	158.4	41.86	10.46	127.6	33.17	3.21	30.8	8.69	
1998	984	14.27	162.7	43.22	10.91	132.7	34.52	3.36	30.0	8.70	
1999	1,003	14.38	161.3	43.08	11.40	137.7	35.90	2.98	23.6	7.18	
2000	1,050	14.93	167.3	44.70	11.93	142.7	37.32	3.00	24.6	7.38	
2001	1,086	16.51	172.0	47.11	12.48	147.7	38.77	4.03	24.3	8.35	
2002	1,112	18.75	176.8	50.21	13.05	152.3	40.15	5.71	24.6	10.09	
2003	1,141	18.48	178.2	50.19	13.61	156.7	41.49	4.87	21.5	8.70	
2004	1,172	18.96	178.4	50.70	14.14	160.7	42.73	4.82	17.7	7.97	
2005	1,196	19.80	181.8	52.15	14.61	163.9	43.77	5.19	17.9	8.38	
2006	1,229	20.30	183.6	52.97	15.08	166.7	44.74	5.22	16.9	8.23	
2007	1,251	20.43	184.6	53.28	15.55	169.5	45.71	4.88	15.1	7.57	
2008	1,270	21.24	188.4	54.76	15.96	171.8	46.53	5.28	16.6	8.23	
2009 (2)	1,278	21.20	190.2	55.03	16.53	176.8	47.99	4.67	13.3	7.04	
2010	1,282	21.50	191.1	55.50	17.11	179.3	49.01	4.39	11.8	6.49	
2011 (3)	1,292	21.91	192.4	56.15	17.59	181.1	49.81	4.32	11.3	6.34	
2012	1,297	22.11	193.0	56.46	18.06	182.6	50.56	4.05	10.4	5.90	
2013	1,300	22.19	193.0	56.53	18.52	184.0	51.25	3.67	9.0	5.28	
2014	1,306	22.37	193.4	56.79	19.03	185.2	51.99	3.34	8.2	4.80	
2015	1,312	23.06	193.8	57.56	19.58	186.5	52.78	3.48	7.3	4.78	
2016	1,315	23.73	194.6	58.37	20.16	187.8	53.58	3.57	6.8	4.79 5.00	
2017	1,319	24.65	195.2	59.39	20.78	188.9	54.39	3.87	6.3	5.00	

⁽¹⁾ Gas plant liquids dropped from system

⁽²⁾ Conversion of historical gas production to 14.73 pressure base.

⁽³⁾ Includes Reserves Justified for Development

Conclusions

As of December 31, 2017, the 1,319 oil and gas fields in the federally regulated part of the Gulf of Mexico Outer Continental Shelf (GOM OCS) contained Original Reserves estimated to be 24.65 billion barrels of oil (BBO) and 195.2 trillion cubic feet (Tcf) of gas. Cumulative Production from the fields accounts for 20.78 BBO and 188.9 Tcf of gas. Reserves are estimated to be 3.87 BBO and 6.3 Tcf of gas for the 489 active fields. Oil Reserves have increased 8.4 percent and the gas Reserves have decreased 7.4 percent since the 2016 report.

CONTRIBUTING PERSONNEL

This report includes contributions from the following Gulf of Mexico Region, Office of Resource Evaluation, personnel.

Holly A. Karrigan

Gregory D. Klocek

Scott M. Palazzo

Mary A. Rather

Jack L. Shanks

Chee W. Yu

REFERENCES

- Attanasi, E.D., 1998, Economics and the National Assessment of United States Oil and Gas Resources, U.S. Geological Survey Circular 1145, United States Government Printing Office, Washington, D.C., Table A-4, p. 29.
- Bascle, B.J., L.D. Nixon, and K.M. Ross, 2001, *Atlas of Gulf of Mexico Gas and Oil Sands as of January 1, 1999*, U.S. Department of the Interior, Minerals Management Service, Gulf of Mexico OCS Regional Office, Office of Resource Evaluation, OCS Report MMS 2001-086, New Orleans, 342 p.
- [BOEM] Bureau of Ocean Energy Management. 2017: 2016a National Assessment of Undiscovered Oil and Gas Resources of the U.S. Outer Continental Shelf, US Department of the Interior, Bureau of Ocean Energy Management. OCS Report BOEM 2017-085.

 Web site: https://www.boem.gov/National-Assessment-2016/
- [BOEM] Bureau of Ocean Energy Management. 2017: Assessment of Technically and Economically Recoverable Hydrocarbon Resources of the Gulf of Mexico Outer Continental Shelf as of January 1, 2014, US Department of the Interior, Bureau of Ocean Energy Management. OCS Report BOEM 2017-005.

 Web site: https://www.boem.gov/BOEM-2017-005/
- Burgess, G. L., E.G. Kazanis, and K.K. Cross, 2018, *Estimated Oil and Gas Reserves, Gulf of Mexico OCS Region, December 31, 2016*, U. S. Department of the Interior, Bureau of Ocean Energy Management, Gulf of Mexico OCS Region, OCS Report BOEM 2018-034, New Orleans, 30p. Web site: https://www.boem.gov/BOEM-2018-034/
- Office of the Federal Register, National Archives and Records Administration, 2018, *Code of Federal Regulations*, 30 CFR, Mineral Resources, U.S. Government Printing Office, Washington, D.C. Web site: https://www.govinfo.gov/content/pkg/CFR-2018-title30-vol2/pdf/CFR-2018-title30-vol2.pdf
- Society of Petroleum Engineers (SPE), American Association of Petroleum Geologists (AAPG), World Petroleum Council (WPC), and Society of Petroleum Evaluation Engineers (SPEE), 2007, Petroleum Resource Management System, 49 p. Web site:

 https://www.spe.org/industry/docs/Petroleum Resources Management System 2007.pdf
- Witrock, R. B., 2017, Biostratigraphic chart of the Gulf of Mexico offshore region, Jurassic to Quaternary, U. S. Department of the Interior, Bureau of Ocean Energy Management, New Orleans, Louisiana. Web site: https://www.data.boem.gov/Paleo/Files/biochart.pdf

APPENDIX A: Definitions of Field, Resource and Reserves Terms

The following definitions as used in this report have been modified from SPE-PRMS and other sources where necessary to conform to requirements of the BOEM Reserves Inventory Program.

Field

A *Field* is an area consisting of a single reservoir or multiple reservoirs all grouped on, or related to, the same general geologic structural feature and/or stratigraphic trapping condition. There may be two or more reservoirs in a field that are separated vertically by impervious strata, laterally by local geologic barriers, or by both. The area may include one OCS lease, a portion of an OCS lease, or a group of OCS leases with one or more wells that have been approved as producible by BOEM (see New Producible Lease). A field is usually named after the area and block on which the discovery well is located. Field names and/or field boundaries may be changed when additional geologic and/or production data initiate such a change. Using geological criteria, BOEM designates a new producible lease as a new field or assigns it to an existing field. http://www.boem.gov/BOEM-Newsroom/Offshore-Stats-and-Facts/Gulf-of-Mexico-Region/Field-Naming-Handbook---March-1996.aspx.

New Producible Lease

A lease that contains at least one well which an operator has requested a well producibility determination, and BOEM has determined that well meets the criteria of producible hydrocarbons defined by 30 CFR 550.115 or 30 CFR 550.116, or a lease that has begun producing.

Project

A *Project* represents the link between the petroleum accumulation and the decision-making process, including budget allocation. A project, for BOEM's classification of Resources and Reserves, is the Field (see also Field).

Resources

Resources encompass all quantities of petroleum (recoverable and unrecoverable) naturally occurring on or within the Earth's crust, discovered and undiscovered, plus those quantities already produced. Further, it includes all types of petroleum whether currently considered conventional or unconventional.

Undiscovered Resources

Resources postulated, on the basis of geologic knowledge and theory, to exist outside of known fields or accumulations. Included also are resources from undiscovered pools within known fields to the extent that they occur within separate plays. BOEM assesses two types of undiscovered resources, *Undiscovered Technically Recoverable Resources (UTRR)* and *Undiscovered Economically Recoverable Resources (UERR)*.

Discovered Resources

Hydrocarbons whose location and quantity are known or estimated from specific geologic evidence are *Discovered Resources*. Included are *Contingent Resources* and *Reserves* depending upon economic, technical, contractual, or regulatory criteria.

Contingent Resources

Those quantities of petroleum estimated, as of a given date, to be potentially recoverable from known accumulations by application of development projects but which are not currently considered to be commercially recoverable due to one or more contingencies.

Reserves

Reserves are those quantities of petroleum anticipated to be commercially recoverable by application of development projects to known accumulations from a given date forward under defined conditions. Reserves must further satisfy four criteria: They must be discovered, recoverable, commercial, and remaining (as of a given date) based on the development project(s) applied. Reserves are further sub-classified based on economic certainty.

Original Reserves Original Reserves are the total of the Cumulative Production and Reserves, as of a specified date.

Proved plus Probable Reserves (2P) The sum of the estimated proved reserves and any additional probable reserves (2P). *Proved Reserves* are commonly defined as those quantities of petroleum which, by analysis of geoscience and engineering data, can be estimated with reasonable certainty to be commercially recoverable, from a given date forward, from known reservoirs and under defined economic conditions, operating methods, and government regulations. *Probable Reserves* are commonly defined as those additional reserves which analysis of geoscience and engineering data indicate are less likely to be recovered than proved reserves but more certain to be recovered than possible reserves.

Reserves
Justified for
Development

The lowest level of reserves certainty. Implementation of the development project is justified on the basis of reasonable forecast commercial conditions at the time of reporting and that there are reasonable expectations that all necessary approvals/contracts will be obtained.

Undeveloped Reserves *Undeveloped Reserves* are those *Reserves* that are expected to be recovered from future wells and facilities, including future improved recovery projects which are anticipated with a high degree of certainty in reservoirs which have previously shown favorable response to improved recovery projects.

Developed Reserves Developed Reserves can be expected to be recovered through existing wells and facilities and by existing operating methods. Improved recovery reserves can be considered as Developed Reserves only after an improved recovery project has been installed and favorable response has occurred or is expected with a reasonable degree of certainty. Developed reserves are expected to be recovered from existing wells, including reserves behind pipe. Improved recovery reserves are considered developed only after the necessary equipment has been installed, or when the costs to do so are relatively minor. Proved Developed Reserves may be sub-categorized as Producing or Non-producing.

Developed Non-producing Reserves Developed Non-producing Reserves are precluded from producing due to being shut-in or behind-pipe. Shut-in includes (1) completion intervals which are open at the time of the estimate, but which have not started producing, (2) wells which were shut-in for market conditions or pipeline connections, or (3) wells not capable of production for mechanical reasons. Behind-pipe refers to zones in existing wells which will require additional completion work or future re-completion prior to the start of production. In both cases, production can be initiated or restored with relatively low expenditure compared to the cost of drilling a new well.

Developed Producing Reserves Developed Producing Reserves are expected to be recovered from completion intervals that are open and producing at the time of the estimate. Improved recovery reserves are considered producing only after the improved recovery project is in operation.

Cumulative Production

Cumulative Production is the sum of all produced volumes of oil and gas prior to a specified date.

Unrecoverable The portion of discovered or undiscovered petroleum-initially-in-place quantities which are estimated, as of a given date, not to be recoverable. A portion of these quantities may become recoverable in the future as commercial circumstances change, technological developments occur, or additional data are acquired.

BOEMChronozone

A body of rock formed during the same time span, bounded by biostratigraphic or correlative seismic markers. Definition taken from BOEM Biostratigraphic Chart of the Gulf of Mexico Region

Assessment Unit

A group of geologically related hydrocarbon accumulations; the term Assessment Unit can refer to groupings of chronozone(s) and/or geologic play(s). Definition modified from the report: 2016a National Assessment of Undiscovered Oil and Gas Resources of the U.S. Outer Continental Shelf

Play

A group of pools that share a common history of hydrocarbon generation, migration, reservoir development, and entrapment. Definition from 2016 National Assessment of Undiscovered Oil and Gas Resources of the U.S. Outer Continental Shelf

Notice

This report, Estimated Oil and Gas Reserves, Gulf of Mexico OCS Region, December 31, 2017, has undergone numerous changes over the last few years. We are continually striving to provide meaningful information to the users of this document. Suggested changes, additions, or deletions to our data or statistical presentations are encouraged so that we can publish the most useful report possible. Please contact the Reserves Section Chief, Grant L. Burgess, at (504) 736-2948 at the Bureau of Ocean Energy Management, 1201 Elmwood Park Boulevard, MS GM773E, New Orleans, Louisiana 70123-2394, to communicate your ideas for consideration in our next report. An overview of the Reserves Inventory Program is available on BOEM's Website.

For free publication and digital data, visit the Gulf of Mexico Web site. The report can be accessed as an Acrobat .pdf (portable document format) file, which allows you to view, print, navigate, and search the document with the free downloadable Acrobat Reader. Digital data used to create the tables and figures presented in the document are also accessible as Excel 2016 spreadsheet files (.xlsx; using Microsoft's Excel spreadsheet viewer, a free file viewer for users without access to Excel). These files are made available in a zipped format, which can be unzipped with the downloadable WinZip program.

For information on this publication contact:

Bureau of Ocean Energy Management Gulf of Mexico OCS Region Attn: Public Information Unit (MS GM250I) 1201 Elmwood Park Boulevard New Orleans, Louisiana 70123-2394 1-800-200-GULF http://www.BOEM.gov

Matthew G. Wilson Regional Supervisor Resource Evaluation



The Department of the Interior Mission

As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering sound use of our land and water resources; protecting our fish, wildlife and biological diversity; preserving the environmental and cultural values of our national parks and historical places; and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to ensure that their development is in the best interests of all our people by encouraging stewardship and citizen participation in their care. The Department also has a major responsibility for American Indian reservation communities and for people who live in island communities.



The Bureau of Ocean Energy Management

The Bureau of Ocean Energy Management (BOEM) works to manage the exploration and development of the nation's offshore resources in a way that appropriately balances economic development, energy independence, and environmental protection through oil and gas leases, renewable energy development and environmental reviews and studies.