

Environmental Studies Program: Ongoing Study

Title	Cook Inlet Physical Oceanography Data Curation, Visualization, and Analyses
Administered by	Alaska Regional Office
BOEM Contact(s)	Dr. Heather Crowley; heather.crowley@boem.gov
Procurement Type(s)	Cooperative Agreement
Conducting Organization(s)	University of Alaska Fairbanks
Total BOEM Cost	\$500,000 (plus joint/in-kind funding)
Performance Period	FY 2022–2025
Final Report Due	July 2025
Date Revised	September 16, 2022
PICOC Summary	
<i><u>Problem</u></i>	Physical oceanographic information for Cook Inlet is not synthesized into a regional framework and is difficult to access. This dispersion of data, particularly for the most recent period, adds complexity for evaluating a changing baseline or for using it in general circulation model (GCM) verification and validation.
<i><u>Intervention</u></i>	This project will collate physical oceanographic data since 2000 in Cook Inlet and Shelikof Strait. Researchers will synthesize the data into a regional framework and identify information needs for sampling and improved modeling approaches to support future oceanographic modeling efforts.
<i><u>Comparison</u></i>	This study will compare areas to assess the patterns and mechanistic causes of spatiotemporal variability and examine the processes that affect the seasonal evolution of water mass exchange between Cook Inlet/Shelikof Strait and adjacent water masses.
<i><u>Outcome</u></i>	A comprehensive and understandable reference of Cook Inlet oceanography datasets and data visualization tools that are readily available to support future environmental analyses and modeling efforts.
<i><u>Context</u></i>	Cook Inlet Planning Area

BOEM Information Need(s): BOEM uses baseline physical oceanographic information, including their seasonal variation and how the baseline is changing, to inform National Environmental Policy Act analyses in Cook Inlet. Teasing apart impacts to the environment from multiple stressors, including a warming Gulf of Alaska, requires historical and up-to-date physical oceanographic measurements. These measurements are also necessary to verify and validate GCM output to assure quality products for use in BOEM’s oil spill trajectory analyses.

Background: Because Cook Inlet oceanographic data depict relatively short-term deployments focused on specific features of Cook Inlet’s complex oceanography, there is a need to: assemble the data and

make it accessible, synthesize it into a comprehensive understanding of the spatial and temporal circulation of the region, and plan for coordinated focused sampling where information needs are identified to improve ocean models (Johnson and Okkonen 2000; Two Crow 2006). Recently Johnson (2021) compiled surface and upper layer Lagrangian drifter data, collected mostly from spring through fall.

Accurate information on surface wind fields, ocean currents, and sea ice is important for oil spill trajectory simulations and the potential impacts on Cook Inlet physical, biological, or social resources from a large spill. It is particularly important to know locations and seasonal changes in oceanographic features that have substantial impact on oil transport. Prior GCM validation by Danielson et al. (2016) identified areas for improvement in Cook Inlet, including a bias towards summer conditions, inability to model high resolution features that are known to impact oil fate in the Inlet (e.g. convergence zones on the scale of ~100 m), and over-stratification of the water column by the model (sometimes by 10 psu). In addition, the Gulf of Alaska is warming substantially (Litzow et al. 2020) and the downstream influences on the oceanography of Cook Inlet and Shelikof Strait are not well documented.

Objectives:

- Improve access to and utility of existing oceanographic data in Cook Inlet.
- Enhance the understanding of the large-scale surface and subsurface circulation and density fields and their interannual variation with focus on these four primary areas:
 - a. Assess the patterns and mechanistic causes of spatiotemporal variability in Cook Inlet and Shelikof Strait subtidal currents.
 - b. Investigate the seasonal impact freshwater coastal discharge (river runoff and glacial ablation) and tides on estuarine circulation and lateral/vertical stratification.
 - c. Examine the processes that affect the seasonal evolution of water mass exchange between Cook Inlet/Shelikof Strait and adjacent water masses, such as those carried by the Alaskan Coastal Current, shelf waters, and Gulf of Alaska basin waters.
- Develop recommendations for additional oceanographic measurements and modeling approaches that can improve our ability to understand and predict Cook Inlet currents.

Methods: This partnership will identify and gather existing, relevant, and readily available physical oceanographic datasets for the Cook Inlet and Shelikof Strait. The datasets will be organized into a common framework for review and identification of specific information needs to guide planning for future fieldwork. Researchers will identify areas with low resolution or dated sampling to develop recommendations for additional measurements needed to provide stratification, freshwater forcing, and higher resolution surface and subsurface current data to enhance the ability to model 3-D currents. Researchers will conduct a field campaign aimed at collecting new measurements that better quantify the dynamics of known rip current zones in middle Cook Inlet. Information from all project components will be synthesized to discuss the physical oceanography of Cook Inlet and Shelikof Strait within a regional framework. Data products and associated metadata will be disseminated through the Cook Inlet Response Tool on AOS web portal.

Specific Research Question(s):

1. What is the current physical oceanographic baseline in Cook Inlet and Shelikof Strait?

2. Where and what types of additional data collections and algorithms will improve ocean model output in Cook Inlet?

Current Status: Awarded

Publications Completed: None

Affiliated WWW Sites: <http://www.boem.gov/akstudies/>

References:

Danielson SL, Hedström KS, Curchitser E. 2016. Cook Inlet Circulation Model Calculations. OCS Study BOEM 2015-050. Anchorage, AK: USDOJ BOEM, Alaska OCS Region. 71 pp.

Johnson MA 2021. Subtidal Surface Circulation in Lower Cook Inlet and Kachemak Bay, Alaska. Regional Studies in Marine Science: 101609. First published online 11 January 2021

Johnson MA, Okkonen SR. 2000. Proceedings Cook Inlet Oceanography Workshop. OCS Study MMS 2000-043. Fairbanks, AK: University of Alaska, Coastal Marine Institute and Oil Spill Recovery Institute. 103 pp.

Two Crow, ed. 2006. Cook Inlet Physical Oceanography Workshop Proceedings. Kenai, AK: AOOS, CIRCAC and Kachemak Bay Research Reserve. 172 pp.