

Table 4.2.4-11

A Summary of Northern Gannet Counts as Observed by ESS During Winter and Summer Aerial Surveys  
Conducted in 2002, 2003, and 2004, Nantucket Sound, Massachusetts

Survey Area	Mar-Apr 2002 Total No/Mean Per Survey (n=5)	May-Aug 2002 Total No/Mean Per Survey (n=6)	Sept 2002- Feb 2003 Total No/Mean Per Survey (n=11)	Mar-Jun 2003 Total No/Mean Per Survey (n=6)	Jun- Aug. 2003 Total No/Mean Per Survey (n=6)	Sept 2003-Feb 2004 Total No/Mean Per Survey (n=12)	Total No.	Percent
Horseshoe Shoal	21/4.2	1/0.2	10/0.9	97/16.2	0/0.0	11/0.9	140	10
Monomoy- Handkerchief Shoal	13/2.6	12/2.0	0/0.0	7/1.2	0/0.0	3/0.2	35	3
Tuckernuck Shoal	32/6.4	0/0.0	17/1.5	101/16.8	0/0.0	9/0.8	159	11
Outside Shoal Areas	294/58.8	76/12.7	160/14.5	268/44.7	0/0.0	283/23.6	1,081	76
Totals	360/72.0	89/14.8	187/17.0	473/78.8	0/0.0	306/25.5	1,415	100