

Table 4.2.7-10

Comparison of MasDMF and NOAA Fisheries Summary Statistics for the Leading Finfish Products in Nantucket Sound: 1998-2007

MassDMF	AVG	S.D.	Cum. % of Total Catch	Est. Avg. Value	Gear	NOAA Fisheries	AVG	S.D.	Cum. % of Total Catch	Est. Avg. Value
Bass, Black Sea	265,937	103,916	20%	\$0.600	<u>P</u> ,W	Squid, general	379,585	252,358	50%	\$0.300
Mackerel, Atlantic	254,775	221,148	39%	\$0.060	<u>W</u> ,G	Fluke	109,224	64,937	64%	\$0.200
Squid, general	242,793	219,441	57%	\$0.200	W	Mackerel, Atlantic	105,976	97,527	78%	\$0.030
Fluke*	229,106	89,938	75%	\$0.500	<u>I</u> ,W,G	Bass, Black Sea	76,303	28,052	88%	\$0.200
Scup	121,957	44,792	84%	\$0.100	W,P	Scup	44,126	31,336	94%	\$0.040
Bass, Striped	98,667	59,678	91%	\$0.200	P	Bluefish	12,072	17,693	95%	\$0.007
Menhaden	45,173	52,024	95%	\$0.005	W	Menhaden	11,286	22,839	97%	\$0.001
Bluefish	39,233	16,117	98%	\$0.020	<u>W</u> ,G	Butterfish	5,637	6,312	98%	\$0.003
Butterfish	9,505	13,105	98%	\$0.006	W	Flounder, Winter	5,130	7,269	98%	\$0.008
Bonito	7,225	10,807	99%	\$0.020	W	Whiting, King	4,153	6,289	99%	\$0.004

Notes:
Annual average catches in pounds and estimated average value in \$m.
*MassDMF Fluke catches are trawl catches averaged over 2006-2007 only.
MassDMF Gear Codes: P=pot; W=weir; G=gillnet; T=trawl